

# Probability theory

## Lecture 6: Variance and covariance

Maksim Zhukovskii

MIPT

## Definitions

- ▶ **Variance** of  $\xi$  is  $\text{Var}\xi = E(\xi - E\xi)^2$ .

## Definitions

- ▶ **Variance** of  $\xi$  is  $\text{Var}\xi = E(\xi - E\xi)^2$ .
- ▶ **Covariance** of  $\xi, \eta$  is  
$$\text{cov}(\xi, \eta) = E((\xi - E\xi)(\eta - E\eta)).$$

## Definitions

- ▶ **Variance** of  $\xi$  is  $\text{Var}\xi = E(\xi - E\xi)^2$ .
- ▶ **Covariance** of  $\xi, \eta$  is  
$$\text{cov}(\xi, \eta) = E((\xi - E\xi)(\eta - E\eta)).$$

Clearly,  $\text{Var}\xi = \text{cov}(\xi, \xi)$ .

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

Proof.

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

Proof.  $\text{Var}\xi = E(\xi - E\xi)^2 =$

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

**Proof.**  $\text{Var}\xi = E(\xi - E\xi)^2 =$   
 $E[\xi^2 - 2\xi E\xi + (E\xi)^2] =$



## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

**Proof.**  $\text{Var}\xi = E(\xi - E\xi)^2 =$   
 $E[\xi^2 - 2\xi E\xi + (E\xi)^2] =$   
 $E\xi^2 - 2(E\xi)^2 + (E\xi)^2 =$

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

**Proof.**  $\text{Var}\xi = E(\xi - E\xi)^2 =$   
 $E[\xi^2 - 2\xi E\xi + (E\xi)^2] =$   
 $E\xi^2 - 2(E\xi)^2 + (E\xi)^2 = E\xi^2 - (E\xi)^2.$

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

Proof.  $\text{Var}\xi = E(\xi - E\xi)^2 =$   
 $E[\xi^2 - 2\xi E\xi + (E\xi)^2] =$   
 $E\xi^2 - 2(E\xi)^2 + (E\xi)^2 = E\xi^2 - (E\xi)^2.$

2.  $\text{Var}\xi \geq 0.$

## Properties of variance

1.  $\text{Var}\xi = E\xi^2 - (E\xi)^2.$

Proof.  $\text{Var}\xi = E(\xi - E\xi)^2 =$   
 $E[\xi^2 - 2\xi E\xi + (E\xi)^2] =$   
 $E\xi^2 - 2(E\xi)^2 + (E\xi)^2 = E\xi^2 - (E\xi)^2.$

2.  $\text{Var}\xi \geq 0.$

Since  $(\xi - E\xi)^2 \geq 0.$

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

Proof.

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

Proof.  $\text{cov}(\xi, \eta) = E(\xi - E\xi)(\eta - E\eta)$

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

**Proof.**  $\text{cov}(\xi, \eta) = E(\xi - E\xi)(\eta - E\eta)$   
 $= E[\xi\eta - \xi E\eta - \eta E\xi + E\xi E\eta] =$



## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

**Proof.**  $\text{cov}(\xi, \eta) = E(\xi - E\xi)(\eta - E\eta)$   
 $= E[\xi\eta - \xi E\eta - \eta E\xi + E\xi E\eta] =$   
 $E\xi\eta - 2E\xi E\eta + E\xi E\eta =$

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

**Proof.**  $\text{cov}(\xi, \eta) = E(\xi - E\xi)(\eta - E\eta)$   
 $= E[\xi\eta - \xi E\eta - \eta E\xi + E\xi E\eta] =$   
 $E\xi\eta - 2E\xi E\eta + E\xi E\eta = E\xi\eta - E\xi E\eta.$

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

**Proof.**  $\text{cov}(\xi, \eta) = E(\xi - E\xi)(\eta - E\eta)$   
 $= E[\xi\eta - \xi E\eta - \eta E\xi + E\xi E\eta] =$   
 $E\xi\eta - 2E\xi E\eta + E\xi E\eta = E\xi\eta - E\xi E\eta.$

2. If  $\xi, \eta$  are independent, then  
 $\text{cov}(\xi, \eta) = 0.$

## Properties of covariance

1.  $\text{cov}(\xi, \eta) = E(\xi\eta) - E\xi E\eta.$

Proof.  $\text{cov}(\xi, \eta) = E(\xi - E\xi)(\eta - E\eta)$   
 $= E[\xi\eta - \xi E\eta - \eta E\xi + E\xi E\eta] =$   
 $E\xi\eta - 2E\xi E\eta + E\xi E\eta = E\xi\eta - E\xi E\eta.$

2. If  $\xi, \eta$  are independent, then  
 $\text{cov}(\xi, \eta) = 0.$

Since  $E\xi\eta = E\xi E\eta.$

## Properties of covariance

3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

## Properties of covariance

3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

4. Covariance is **bilinear**:

$$\begin{aligned} \text{cov}(a_1\xi_1 + a_2\xi_2, \eta) = \\ a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta). \end{aligned}$$

## Properties of covariance

3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

4. Covariance is **bilinear**:

$$\begin{aligned} \text{cov}(a_1\xi_1 + a_2\xi_2, \eta) = \\ a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta). \end{aligned}$$

Proof.

## Properties of covariance

3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

4. Covariance is **bilinear**:

$$\begin{aligned} \text{cov}(a_1\xi_1 + a_2\xi_2, \eta) = \\ a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta). \end{aligned}$$

**Proof.**  $\text{cov}(a_1\xi_1 + a_2\xi_2, \eta) =$



## Properties of covariance

### 3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

### 4. Covariance is **bilinear**:

$$\begin{aligned} \text{cov}(a_1\xi_1 + a_2\xi_2, \eta) = \\ a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta). \end{aligned}$$

**Proof.**  $\text{cov}(a_1\xi_1 + a_2\xi_2, \eta) =$   
 $E(a_1\xi_1\eta + a_2\xi_2\eta) - E(a_1\xi_1 + a_2\xi_2)E\eta =$

## Properties of covariance

### 3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

### 4. Covariance is **bilinear**:

$$\begin{aligned} \text{cov}(a_1\xi_1 + a_2\xi_2, \eta) = \\ a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta). \end{aligned}$$

**Proof.**  $\text{cov}(a_1\xi_1 + a_2\xi_2, \eta) =$   
 $E(a_1\xi_1\eta + a_2\xi_2\eta) - E(a_1\xi_1 + a_2\xi_2)E\eta =$   
 $a_1E\xi_1\eta + a_2E\xi_2\eta - a_1E\xi_1E\eta - a_2E\xi_2E\eta$

## Properties of covariance

### 3. Covariance is **symmetric**:

$$\text{cov}(\xi, \eta) = \text{cov}(\eta, \xi).$$

### 4. Covariance is **bilinear**:

$$\begin{aligned}\text{cov}(a_1\xi_1 + a_2\xi_2, \eta) = \\ a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta).\end{aligned}$$

**Proof.**  $\text{cov}(a_1\xi_1 + a_2\xi_2, \eta) =$   
 $E(a_1\xi_1\eta + a_2\xi_2\eta) - E(a_1\xi_1 + a_2\xi_2)E\eta =$   
 $a_1E\xi_1\eta + a_2E\xi_2\eta - a_1E\xi_1E\eta - a_2E\xi_2E\eta$   
 $= a_1\text{cov}(\xi_1, \eta) + a_2\text{cov}(\xi_2, \eta).$

## Variance of a sum of random variables

### Claim

$$\begin{aligned} \text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j). \end{aligned}$$

## Variance of a sum of random variables

### Claim

$$\begin{aligned} \text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j). \end{aligned}$$

Proof.

## Variance of a sum of random variables

### Claim

$$\begin{aligned} \text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j). \end{aligned}$$

**Proof.**  $\text{Var}(\xi_1 + \dots + \xi_n) =$

## Variance of a sum of random variables

### Claim

$$\begin{aligned} \text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j). \end{aligned}$$

**Proof.**  $\text{Var}(\xi_1 + \dots + \xi_n) =$   
 $\text{cov}(\xi_1 + \dots + \xi_n, \xi_1 + \dots + \xi_n) =$

## Variance of a sum of random variables

### Claim

$$\begin{aligned} \text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j). \end{aligned}$$

**Proof.**  $\text{Var}(\xi_1 + \dots + \xi_n) =$   
 $\text{cov}(\xi_1 + \dots + \xi_n, \xi_1 + \dots + \xi_n) =$   
 $\sum_{i=1}^n \text{cov}(\xi_i, \xi_i) + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j) =$



## Variance of a sum of random variables

### Claim

$$\begin{aligned} \text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j). \end{aligned}$$

**Proof.**  $\text{Var}(\xi_1 + \dots + \xi_n) =$   
 $\text{cov}(\xi_1 + \dots + \xi_n, \xi_1 + \dots + \xi_n) =$   
 $\sum_{i=1}^n \text{cov}(\xi_i, \xi_i) + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j) =$   
 $\text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j).$

## Variance of a sum of random variables

### Claim

$$\begin{aligned}\text{Var}(\xi_1 + \dots + \xi_n) = \\ \text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j).\end{aligned}$$

**Proof.**  $\text{Var}(\xi_1 + \dots + \xi_n) =$   
 $\text{cov}(\xi_1 + \dots + \xi_n, \xi_1 + \dots + \xi_n) =$   
 $\sum_{i=1}^n \text{cov}(\xi_i, \xi_i) + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j) =$   
 $\text{Var}\xi_1 + \dots + \text{Var}\xi_n + \sum_{i \neq j} \text{cov}(\xi_i, \xi_j).$

If  $\xi_1, \dots, \xi_n$  are pairwise independent, then  
 $\text{Var}(\xi_1 + \dots + \xi_n) = \text{Var}(\xi_1) + \dots + \text{Var}(\xi_n).$

## Variance of a binomial random variable

$$\xi \sim \text{Bin}(n, p).$$

## Variance of a binomial random variable

$\xi \sim \text{Bin}(n, p)$ .

$$E\xi^2 = \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k} =$$

## Variance of a binomial random variable

$\xi \sim \text{Bin}(n, p)$ .

$$\begin{aligned} E\xi^2 &= \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + \\ &\sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} = \end{aligned}$$

## Variance of a binomial random variable

$\xi \sim \text{Bin}(n, p)$ .

$$E\xi^2 = \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k} =$$

$$\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} +$$

$$\sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} =$$

$$\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + np =$$

## Variance of a binomial random variable

$\xi \sim \text{Bin}(n, p)$ .

$$\begin{aligned} E\xi^2 &= \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + \\ &\sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + np = \\ &n(n-1)p^2 \sum_{k=2}^n \binom{n-2}{k-2} p^{k-2} (1-p)^{n-k} + np = \end{aligned}$$

## Variance of a binomial random variable

$\xi \sim \text{Bin}(n, p)$ .

$$\begin{aligned} E\xi^2 &= \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + \\ &\sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + np = \\ &n(n-1)p^2 \sum_{k=2}^n \binom{n-2}{k-2} p^{k-2} (1-p)^{n-k} + np = \\ &n(n-1)p^2 \sum_{j=0}^{n-2} \binom{n-2}{j} p^j (1-p)^{n-2-j} + np = \\ &n(n-1)p^2 + np. \end{aligned}$$



## Variance of a binomial random variable

$\xi \sim \text{Bin}(n, p)$ .

$$\begin{aligned} E\xi^2 &= \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + \\ &\sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} = \\ &\sum_{k=1}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} + np = \\ &n(n-1)p^2 \sum_{k=2}^n \binom{n-2}{k-2} p^{k-2} (1-p)^{n-k} + np = \\ &n(n-1)p^2 \sum_{j=0}^{n-2} \binom{n-2}{j} p^j (1-p)^{n-2-j} + np = \\ &n(n-1)p^2 + np. \end{aligned}$$

$$\text{Var}\xi = n^2 p^2 - np^2 + np - n^2 p^2 = np(1-p).$$

## Variance of a binomial random variable

$$\xi \sim \text{Bin}(n, p).$$

## Variance of a binomial random variable

$$\xi \sim \text{Bin}(n, p).$$

Let  $\xi_1, \dots, \xi_n$  are independent Bernoulli random variables.

## Variance of a binomial random variable

$$\xi \sim \text{Bin}(n, p).$$

Let  $\xi_1, \dots, \xi_n$  are independent Bernoulli random variables.

Then  $\xi \stackrel{d}{=} \xi_1 + \dots + \xi_n$ .

## Variance of a binomial random variable

$$\xi \sim \text{Bin}(n, p).$$

Let  $\xi_1, \dots, \xi_n$  are independent Bernoulli random variables.

Then  $\xi \stackrel{d}{=} \xi_1 + \dots + \xi_n$ .

$$\begin{aligned} \text{Var}\xi &= n\text{Var}\xi_1 = n(\text{E}\xi_1^2 - (\text{E}\xi_1)^2) = \\ &= np(1 - p). \end{aligned}$$

## Other examples

- ▶  $\xi \sim U\{1, \dots, n\}$

## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\text{Var}\xi = \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 =$$

## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\begin{aligned}\text{Var}\xi &= \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 = \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.\end{aligned}$$



## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\begin{aligned}\text{Var}\xi &= \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 = \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.\end{aligned}$$

►  $\xi \sim \mathcal{N}(0, 1)$

## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\begin{aligned}\text{Var}\xi &= \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 = \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.\end{aligned}$$

►  $\xi \sim \mathcal{N}(0, 1)$

$$\text{Var}\xi = \int_{-\infty}^{\infty} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} dx =$$

## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\begin{aligned}\text{Var}\xi &= \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 = \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.\end{aligned}$$

►  $\xi \sim \mathcal{N}(0, 1)$

$$\begin{aligned}\text{Var}\xi &= \int_{-\infty}^{\infty} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} dx = \\ &= - \int_{-\infty}^{\infty} \frac{x}{\sqrt{2\pi}} de^{-x^2/2} =\end{aligned}$$

## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\begin{aligned}\text{Var}\xi &= \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 = \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.\end{aligned}$$

►  $\xi \sim \mathcal{N}(0, 1)$

$$\begin{aligned}\text{Var}\xi &= \int_{-\infty}^{\infty} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} dx = \\ &= \int_{-\infty}^{\infty} \frac{x}{\sqrt{2\pi}} d e^{-x^2/2} = \\ &= -\frac{x}{\sqrt{2\pi}} e^{-x^2/2} \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx =\end{aligned}$$

## Other examples

►  $\xi \sim U\{1, \dots, n\}$

$$\begin{aligned}\text{Var}\xi &= \frac{1}{n} \sum_{i=1}^n i^2 - \left(\frac{n+1}{2}\right)^2 = \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.\end{aligned}$$

►  $\xi \sim \mathcal{N}(0, 1)$

$$\begin{aligned}\text{Var}\xi &= \int_{-\infty}^{\infty} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} dx = \\ &= \int_{-\infty}^{\infty} \frac{x}{\sqrt{2\pi}} d e^{-x^2/2} = \\ &= -\frac{x}{\sqrt{2\pi}} e^{-x^2/2} \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 1.\end{aligned}$$

# Cauchy–Bunyakovsky–Schwarz inequality

## Theorem

*Let  $E\xi^2 < \infty$ ,  $E\eta^2 < \infty$ .*

# Cauchy–Bunyakovsky–Schwarz inequality

## Theorem

*Let  $E\xi^2 < \infty$ ,  $E\eta^2 < \infty$ .*

*Then  $E|\xi\eta| < \infty$  and  $(E|\xi\eta|)^2 \leq E\xi^2 E\eta^2$ .*

## The proof

- ▶ Let  $E\xi^2 = 0$ .



## The proof

► Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1$$

## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1$$

## The proof

► Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow \\ E|\xi\eta| = 0.$$

## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow \\ E|\xi\eta| = 0.$$

- ▶ Let  $E\xi^2 > 0$ ,  $E\eta^2 > 0$ .

## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow \\ E|\xi\eta| = 0.$$

- ▶ Let  $E\xi^2 > 0$ ,  $E\eta^2 > 0$ .

$$\text{Denote } \xi_0 = \xi / \sqrt{E\xi^2}, \eta_0 = \eta / \sqrt{E\eta^2}.$$

## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow \\ E|\xi\eta| = 0.$$

- ▶ Let  $E\xi^2 > 0$ ,  $E\eta^2 > 0$ .

$$\text{Denote } \xi_0 = \xi / \sqrt{E\xi^2}, \eta_0 = \eta / \sqrt{E\eta^2}.$$

$$2|\xi_0\eta_0| \leq \xi_0^2 + \eta_0^2$$

## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow E|\xi\eta| = 0.$$

- ▶ Let  $E\xi^2 > 0$ ,  $E\eta^2 > 0$ .

$$\text{Denote } \xi_0 = \xi / \sqrt{E\xi^2}, \eta_0 = \eta / \sqrt{E\eta^2}.$$

$$2|\xi_0\eta_0| \leq \xi_0^2 + \eta_0^2 \Rightarrow$$

$$E(2|\xi_0\eta_0|) \leq E(\xi_0^2 + \eta_0^2) = 2$$

## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow E|\xi\eta| = 0.$$

- ▶ Let  $E\xi^2 > 0$ ,  $E\eta^2 > 0$ .

$$\text{Denote } \xi_0 = \xi / \sqrt{E\xi^2}, \eta_0 = \eta / \sqrt{E\eta^2}.$$

$$2|\xi_0\eta_0| \leq \xi_0^2 + \eta_0^2 \Rightarrow$$

$$E(2|\xi_0\eta_0|) \leq E(\xi_0^2 + \eta_0^2) = 2 \Rightarrow$$

$$E|\xi_0\eta_0| \leq 1$$



## The proof

- ▶ Let  $E\xi^2 = 0$ .

$$P(\xi = 0) = 1 \Rightarrow P(\xi\eta = 0) = 1 \Rightarrow E|\xi\eta| = 0.$$

- ▶ Let  $E\xi^2 > 0$ ,  $E\eta^2 > 0$ .

$$\text{Denote } \xi_0 = \xi / \sqrt{E\xi^2}, \eta_0 = \eta / \sqrt{E\eta^2}.$$

$$2|\xi_0\eta_0| \leq \xi_0^2 + \eta_0^2 \Rightarrow$$

$$E(2|\xi_0\eta_0|) \leq E(\xi_0^2 + \eta_0^2) = 2 \Rightarrow$$

$$E|\xi_0\eta_0| \leq 1 \Rightarrow E|\xi\eta| \leq \sqrt{E\xi^2 E\eta^2}.$$

# Markov inequality

## Theorem

*Let*  $E|\xi| < \infty$ .

# Markov inequality

## Theorem

Let  $E|\xi| < \infty$ .

Then, for every  $a > 0$ ,  $P(|\xi| \geq a) \leq \frac{E|\xi|}{a}$ .

## The proof

$$E|\xi| =$$

## The proof

$$E|\xi| = E(|\xi|I(|\xi| < a) + |\xi|I(|\xi| \geq a))$$

## The proof

$$E|\xi| = E(|\xi|I(|\xi| < a) + |\xi|I(|\xi| \geq a))$$

$$\geq aE(I(|\xi| \geq a))$$

## The proof

$$E|\xi| = E(|\xi|I(|\xi| < a) + |\xi|I(|\xi| \geq a))$$

$$\geq aE(I(|\xi| \geq a)) = aP(|\xi| \geq a).$$

## The proof

$$E|\xi| = E(|\xi|I(|\xi| < a) + |\xi|I(|\xi| \geq a))$$

$$\geq aE(I(|\xi| \geq a)) = aP(|\xi| \geq a).$$

Therefore,  $P(|\xi| \geq a) \leq \frac{E|\xi|}{a}$ .



# Chebyshev inequality

## Theorem

*Let  $E\xi^2 < \infty$ .*

# Chebyshev inequality

## Theorem

*Let  $E\xi^2 < \infty$ .*

*Then, for every  $a > 0$ ,*

$$P(|\xi - E\xi| \geq a) \leq \frac{\text{Var}\xi}{a^2}.$$

# Chebyshev inequality

## Theorem

Let  $E\xi^2 < \infty$ .

Then, for every  $a > 0$ ,

$$P(|\xi - E\xi| \geq a) \leq \frac{\text{Var}\xi}{a^2}.$$

**Proof.** Follows from Markov inequality by replacing  $\xi \rightarrow (\xi - E\xi)^2$ ,  $a \rightarrow a^2$ .

# Jensen's inequality

## Theorem

*Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be convex and  $E|\xi| < \infty$ .*

# Jensen's inequality

## Theorem

*Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be convex and  $E|\xi| < \infty$ .*

*Then  $g(E\xi) \leq Eg(\xi)$ .*

## The proof

For every  $x_0$ , there exists  $\lambda(x_0)$  such that  
 $g(x) \geq g(x_0) + \lambda(x_0)(x - x_0)$  for all  $x \in \mathbb{R}$ .

## The proof

For every  $x_0$ , there exists  $\lambda(x_0)$  such that  
 $g(x) \geq g(x_0) + \lambda(x_0)(x - x_0)$  for all  $x \in \mathbb{R}$ .

Set  $x_0 = E\xi$ ,  $x = \xi$ .

## The proof

For every  $x_0$ , there exists  $\lambda(x_0)$  such that  
 $g(x) \geq g(x_0) + \lambda(x_0)(x - x_0)$  for all  $x \in \mathbb{R}$ .

Set  $x_0 = E\xi$ ,  $x = \xi$ .

Then  $g(\xi) \geq g(E\xi) + \lambda(E\xi)(\xi - E\xi)$ .



## The proof

For every  $x_0$ , there exists  $\lambda(x_0)$  such that  
 $g(x) \geq g(x_0) + \lambda(x_0)(x - x_0)$  for all  $x \in \mathbb{R}$ .

Set  $x_0 = E\xi$ ,  $x = \xi$ .

Then  $g(\xi) \geq g(E\xi) + \lambda(E\xi)(\xi - E\xi)$ .

Therefore,  $Eg(\xi) \geq g(E\xi)$ .

# Law of large numbers

## Theorem

*Let  $\xi_1, \xi_2, \dots$  be pairwise independent identically distributed random variables.  
Let  $E\xi_1^2 < \infty$ .*

# Law of large numbers

## Theorem

*Let  $\xi_1, \xi_2, \dots$  be pairwise independent identically distributed random variables.*

*Let  $E\xi_1^2 < \infty$ .*

*Then, for all  $\varepsilon > 0$*

$$P\left(\frac{|S_n - ES_n|}{n} > \varepsilon\right) \rightarrow 0.$$

## The proof

$$\mathbb{P} \left( \frac{|S_n - \mathbb{E}S_n|}{n} > \varepsilon \right) \leq \frac{\text{Var} S_n}{n^2 \varepsilon^2} =$$

## The proof

$$\mathbb{P} \left( \frac{|S_n - \mathbb{E}S_n|}{n} > \varepsilon \right) \leq \frac{\text{Var}S_n}{n^2\varepsilon^2} =$$
$$\frac{\text{Var}\xi_1}{\varepsilon^2 n}$$

## The proof

$$\begin{aligned} \mathbb{P} \left( \frac{|S_n - \mathbb{E}S_n|}{n} > \varepsilon \right) &\leq \frac{\text{Var}S_n}{n^2\varepsilon^2} = \\ &\frac{\text{Var}\xi_1}{\varepsilon^2 n} \rightarrow 0, \quad n \rightarrow \infty. \end{aligned}$$

## A generalization

### Theorem

*Let  $\xi_1, \xi_2, \dots$  be a sequence of random variables such that*

- ▶ *for  $i \neq j$ ,  $\text{cov}(\xi_i, \xi_j) = 0$ ,*
- ▶ *there exists  $C > 0$  such that, for every  $i$ ,  $\text{Var}\xi_i < C$ .*

## A generalization

### Theorem

*Let  $\xi_1, \xi_2, \dots$  be a sequence of random variables such that*

- ▶ for  $i \neq j$ ,  $\text{cov}(\xi_i, \xi_j) = 0$ ,*
- ▶ there exists  $C > 0$  such that, for every  $i$ ,  $\text{Var}\xi_i < C$ .*

*Then, for all  $\varepsilon > 0, \delta > 0$*

$$P\left(\frac{|S_n - ES_n|}{n^{1/2+\delta}} > \varepsilon\right) \rightarrow 0.$$