

Probability theory

Lecture 4: Random variables and vectors

Maksim Zhukovskii

MIPT

Measurable functions

\mathcal{A} — σ -algebra on A , \mathcal{B} — σ -algebra on B

Measurable functions

\mathcal{A} — σ -algebra on A , \mathcal{B} — σ -algebra on B

- ▶ $f : A \rightarrow B$ is $(\mathcal{A}|\mathcal{B})$ -measurable, if,
for every $X \in \mathcal{B}$, $f^{-1}(X) \in \mathcal{A}$.

Measurable functions

\mathcal{A} — σ -algebra on A , \mathcal{B} — σ -algebra on B

- ▶ $f : A \rightarrow B$ is $(\mathcal{A}|\mathcal{B})$ -measurable, if,
for every $X \in \mathcal{B}$, $f^{-1}(X) \in \mathcal{A}$.

Theorem

Let $\mathcal{M} \subset \mathcal{B}$ such that $\sigma(\mathcal{M}) = \mathcal{B}$.

Measurable functions

\mathcal{A} — σ -algebra on A , \mathcal{B} — σ -algebra on B

- ▶ $f : A \rightarrow B$ is $(\mathcal{A}|\mathcal{B})$ -measurable, if,
for every $X \in \mathcal{B}$, $f^{-1}(X) \in \mathcal{A}$.

Theorem

Let $\mathcal{M} \subset \mathcal{B}$ such that $\sigma(\mathcal{M}) = \mathcal{B}$.

F is $(\mathcal{A}|\mathcal{B})$ -measurable

if and only if,

for every $X \in \mathcal{M}$, $f^{-1}(X) \in \mathcal{A}$.

The proof

Random variables and random vectors

(Ω, \mathcal{F}, P) — probability space,
 \mathcal{E} — σ -algebra on E .

Random variables and random vectors

(Ω, \mathcal{F}, P) — probability space,
 \mathcal{E} — σ -algebra on E .

- ▶ If $f : \Omega \rightarrow E$ is $(\mathcal{F}|\mathcal{E})$ -measurable, then it is called a random element.

Random variables and random vectors

(Ω, \mathcal{F}, P) — probability space,
 \mathcal{E} — σ -algebra on E .

- ▶ If $f : \Omega \rightarrow E$ is $(\mathcal{F}|\mathcal{E})$ -measurable, then it is called a random element.

$E = \mathbb{R}, \mathcal{E} = \mathcal{B}(\mathbb{R}) \Rightarrow$
 f — a random variable,

Random variables and random vectors

(Ω, \mathcal{F}, P) — probability space,
 \mathcal{E} — σ -algebra on E .

- ▶ If $f : \Omega \rightarrow E$ is $(\mathcal{F}|\mathcal{E})$ -measurable, then it is called a random element.

$$E = \mathbb{R}, \mathcal{E} = \mathcal{B}(\mathbb{R}) \Rightarrow$$

f — a random variable,

$$E = \mathbb{R}^n, \mathcal{E} = \mathcal{B}(\mathbb{R}^n)$$

$\Rightarrow f$ — a random vector.

Why do we need measurability?

Example: an indicator random variable

Let $A \in \mathcal{F}$.

Example: an indicator random variable

Let $A \in \mathcal{F}$.

Indicator of A

$$I_A : \Omega \rightarrow \{0, 1\},$$

$$I_A(\omega) = 1 \text{ if and only if } \omega \in A.$$

Functions of random variables

- ▶ $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ — Borel function,
if it is $(\mathcal{B}(\mathbb{R}^n) | \mathcal{B}(\mathbb{R}^k))$ -measurable

Functions of random variables

- ▶ $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ — Borel function,
if it is $(\mathcal{B}(\mathbb{R}^n) | \mathcal{B}(\mathbb{R}^k))$ -measurable

Theorem

*If ξ is an n -dimensional random vector,
 $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is a Borel function,*

Functions of random variables

- ▶ $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ — Borel function,
if it is $(\mathcal{B}(\mathbb{R}^n) | \mathcal{B}(\mathbb{R}^k))$ -measurable

Theorem

*If ξ is an n -dimensional random vector,
 $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is a Borel function,
then $f(\xi)$ is a random vector as well.*

The proof

Components of a random vector

Theorem

Let $\xi : \Omega \rightarrow \mathbb{R}^n$.

Components of a random vector

Theorem

Let $\xi : \Omega \rightarrow \mathbb{R}^n$.

$\xi = (\xi_1, \dots, \xi_n)$ is a random vector

if and only if

ξ_1, \dots, ξ_n are random variables.

The proof

Continuous functions

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is continuous (or has a finite number of discontinuities), then f is Borel.

Continuous functions

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is continuous (or has a finite number of discontinuities), then f is Borel.

Corollary

If ξ, η are random variables, then

$\xi + \eta, \xi - \eta, \xi\eta, (\xi/\eta)I(\eta \neq 0)$
are random variables as well.

Limits

Theorem

Let ξ_1, ξ_2, \dots be random variables.

Then $\overline{\lim}_{n \rightarrow \infty} \xi_n$, $\underline{\lim}_{n \rightarrow \infty} \xi_n$, $\sup_n \xi_n$, $\inf_n \xi_n$ are random variables as well.

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$



The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$



$$\exists \varepsilon > 0$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$



$$\exists \varepsilon > 0 \forall n$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$



$$\exists \varepsilon > 0 \forall n \exists k \geq n$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$



$$\exists \varepsilon > 0 \forall n \exists k \geq n \quad x_k > x + \varepsilon$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$

$$\Leftrightarrow$$

$$\exists \varepsilon > 0 \forall n \exists k \geq n \quad x_k > x + \varepsilon$$

$$\{\overline{\lim}_{n \rightarrow \infty} \xi_n > x\} =$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$

$$\Leftrightarrow$$

$$\exists \varepsilon > 0 \forall n \exists k \geq n \quad x_k > x + \varepsilon$$

$$\{\overline{\lim}_{n \rightarrow \infty} \xi_n > x\} =$$

$$\bigcup_{m=1}^{\infty} \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \{\xi_k > x + \frac{1}{m}\}$$

The proof

$$\overline{\lim}_{n \rightarrow \infty} \xi_n$$

$$\overline{\lim}_{n \rightarrow \infty} x_n > x$$

$$\Leftrightarrow$$

$$\exists \varepsilon > 0 \forall n \exists k \geq n \quad x_k > x + \varepsilon$$

$$\{\overline{\lim}_{n \rightarrow \infty} \xi_n > x\} =$$

$$\bigcup_{m=1}^{\infty} \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \{\xi_k > x + \frac{1}{m}\} \in \mathcal{F}$$

Probability distribution

Theorem

The function

$$P_\xi : \mathcal{B}(\mathbb{R}^n) \rightarrow [0, 1], \quad P_\xi(B) = P(\xi \in B),$$

is a probability on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$.

Probability distribution

- ▶ P_ξ is called **distribution** of ξ .

Probability distribution

- ▶ P_ξ is called **distribution of ξ** .
- ▶ The distribution function F_ξ of P_ξ is called **distribution function of ξ** .

Probability distribution

- ▶ P_ξ is called **distribution of ξ** .
- ▶ The distribution function F_ξ of P_ξ is called **distribution function of ξ** .
- ▶ If P_ξ is absolutely continuous, then its density p_ξ is called **density of ξ** , ξ is called **absolutely continuous**.

Probability distribution

- ▶ P_ξ is called **distribution of ξ** .
- ▶ The distribution function F_ξ of P_ξ is called **distribution function of ξ** .
- ▶ If P_ξ is absolutely continuous, then its density p_ξ is called **density of ξ** , ξ is called **absolutely continuous**.
- ▶ If P_ξ is discrete, then ξ is called **discrete** as well.

Notations

$$P(\{\omega : \xi(\omega) \in B\}) =: P(\xi \in B).$$

Notations

$$P(\{\omega : \xi(\omega) \in B\}) =: P(\xi \in B).$$

In particular,

Notations

$$P(\{\omega : \xi(\omega) \in B\}) =: P(\xi \in B).$$

In particular,

1. $P(\{\omega : \xi(\omega) < x\}) =: P(\xi < x),$

Notations

$$P(\{\omega : \xi(\omega) \in B\}) =: P(\xi \in B).$$

In particular,

1. $P(\{\omega : \xi(\omega) < x\}) =: P(\xi < x),$
2. $P(\{\omega : \xi(\omega) = x\}) =: P(\xi = x),$

Notations

$$P(\{\omega : \xi(\omega) \in B\}) =: P(\xi \in B).$$

In particular,

1. $P(\{\omega : \xi(\omega) < x\}) =: P(\xi < x),$

2. $P(\{\omega : \xi(\omega) = x\}) =: P(\xi = x),$

3. $F_\xi(x) = P(\{\omega : \xi(\omega) \leq x\}) =: P(\xi \leq x).$

Examples

1. Let ξ take values 0,1,2,3 with equal probabilities $1/4$.

Examples

1. Let ξ take values 0,1,2,3 with equal probabilities $1/4$.

$$F_{\xi}(x) = \begin{array}{ll} 0, & x < 0, \\ \frac{1}{4}, & 0 \leq x < 1, \\ \frac{1}{2}, & 1 \leq x < 2, \\ \frac{3}{4}, & 2 \leq x < 3, \\ 1, & x \geq 3. \end{array}$$

Examples

2. ξ — a number chosen from $[1, 4]$ uniformly at random.

Examples

2. ξ — a number chosen from $[1, 4]$ uniformly at random.

$$F_{\xi}(x) = \frac{x-1}{3}I(1 \leq x < 4) + I(x \geq 4),$$

Examples

2. ξ — a number chosen from $[1, 4]$ uniformly at random.

$$F_{\xi}(x) = \frac{x-1}{3}I(1 \leq x < 4) + I(x \geq 4),$$

$$p_{\xi}(x) = \frac{1}{3}I(1 \leq x \leq 4).$$

Independent random vectors

- ▶ ξ_1, ξ_2 are independent,

Independent random vectors

- ▶ ξ_1, ξ_2 are **independent**, if
for all $B_1 \in \mathcal{B}(\mathbb{R}^{k_1})$, $B_2 \in \mathcal{B}(\mathbb{R}^{k_2})$,
$$P(\xi_1 \in B_1, \xi_2 \in B_2) = P(\xi_1 \in B_1)P(\xi_2 \in B_2).$$

Independent random vectors

- ▶ ξ_1, ξ_2 are **independent**, if
for all $B_1 \in \mathcal{B}(\mathbb{R}^{k_1})$, $B_2 \in \mathcal{B}(\mathbb{R}^{k_2})$,
$$P(\xi_1 \in B_1, \xi_2 \in B_2) = P(\xi_1 \in B_1)P(\xi_2 \in B_2).$$
- ▶ ξ_1, \dots, ξ_n are **pairwise independent**,

Independent random vectors

- ▶ ξ_1, ξ_2 are **independent**, if
for all $B_1 \in \mathcal{B}(\mathbb{R}^{k_1})$, $B_2 \in \mathcal{B}(\mathbb{R}^{k_2})$,
$$P(\xi_1 \in B_1, \xi_2 \in B_2) = P(\xi_1 \in B_1)P(\xi_2 \in B_2).$$
- ▶ ξ_1, \dots, ξ_n are **pairwise independent**,
if, for every i, j , ξ_i, ξ_j are independent.

Independent random vectors

- ▶ ξ_1, \dots, ξ_n are mutually independent (simply independent),

Independent random vectors

- ▶ ξ_1, \dots, ξ_n are mutually independent (simply independent), if,
for all $B_1 \in \mathcal{B}(\mathbb{R}^{k_1}), \dots, B_n \in \mathcal{B}(\mathbb{R}^{k_n})$,
$$P(\xi_1 \in B_1, \dots, \xi_n \in B_n) =$$
$$P(\xi_1 \in B_1) \dots P(\xi_n \in B_n).$$

Independent random vectors

- ▶ ξ_1, \dots, ξ_n are mutually independent (simply independent), if,
for all $B_1 \in \mathcal{B}(\mathbb{R}^{k_1}), \dots, B_n \in \mathcal{B}(\mathbb{R}^{k_n})$,
$$P(\xi_1 \in B_1, \dots, \xi_n \in B_n) = P(\xi_1 \in B_1) \dots P(\xi_n \in B_n).$$
- ▶ Random vectors from $\{\xi_\alpha\}_{\alpha \in \mathcal{A}}$ are independent,

Independent random vectors

- ▶ ξ_1, \dots, ξ_n are mutually independent (simply independent), if,
for all $B_1 \in \mathcal{B}(\mathbb{R}^{k_1}), \dots, B_n \in \mathcal{B}(\mathbb{R}^{k_n})$,
$$P(\xi_1 \in B_1, \dots, \xi_n \in B_n) = P(\xi_1 \in B_1) \dots P(\xi_n \in B_n).$$
- ▶ Random vectors from $\{\xi_\alpha\}_{\alpha \in \mathcal{A}}$ are independent, if,
for every $n \in \mathbb{N}$ and any $t_1, \dots, t_n \in \mathcal{A}$,
the random vectors $\xi_{t_1}, \dots, \xi_{t_n}$ are independent.

Independent discrete random variables

Theorem

Discrete random variables ξ_1, \dots, ξ_n are independent

Independent discrete random variables

Theorem

Discrete random variables ξ_1, \dots, ξ_n are independent

if and only if,

Independent discrete random variables

Theorem

Discrete random variables ξ_1, \dots, ξ_n are independent

if and only if,

for all $x_1, \dots, x_n \in \mathbb{R}$,

$$P(\xi_1 = x_1, \dots, \xi_n = x_n) =$$

$$P(\xi_1 = x_1) \dots P(\xi_n = x_n).$$

The proof

Criterion of independence

Theorem

Random variables ξ_1, \dots, ξ_n are independent

Criterion of independence

Theorem

*Random variables ξ_1, \dots, ξ_n are independent
if and only if,*

Criterion of independence

Theorem

Random variables ξ_1, \dots, ξ_n are independent

if and only if,

for every $x = (x_1, \dots, x_n) \in \mathbb{R}^n$,

$$F_{(\xi_1, \dots, \xi_n)}(x) = \prod_{i=1}^n F_{\xi_i}(x_i).$$

Criterion of independence

Theorem

Random variables ξ_1, \dots, ξ_n are independent

if and only if,

for every $x = (x_1, \dots, x_n) \in \mathbb{R}^n$,

$$F_{(\xi_1, \dots, \xi_n)}(x) = \prod_{i=1}^n F_{\xi_i}(x_i).$$

The same is true for random **vectors**.

Functions of independent vectors

Theorem

Let

- ▶ $\xi = (\xi_1, \dots, \xi_{n_1})$ $\eta = (\eta_1, \dots, \eta_{n_2})$ *be independent random vectors,*
- ▶ $f : \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{k_1}$, $g : \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{k_2}$ *be Borel functions.*

Functions of independent vectors

Theorem

Let

- ▶ $\xi = (\xi_1, \dots, \xi_{n_1})$ $\eta = (\eta_1, \dots, \eta_{n_2})$ *be independent random vectors,*
- ▶ $f : \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{k_1}$, $g : \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{k_2}$ *be Borel functions.*

Then $f(\xi)$, $g(\eta)$ are independent.

Functions of independent vectors

Theorem

Let

- ▶ $\xi = (\xi_1, \dots, \xi_{n_1})$ $\eta = (\eta_1, \dots, \eta_{n_2})$ *be independent random vectors,*
- ▶ $f : \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{k_1}$, $g : \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{k_2}$ *be Borel functions.*

Then $f(\xi)$, $g(\eta)$ are independent.

The same is true for **several** vectors.

The proof

Convolution

Theorem

Let ξ, η be independent absolutely continuous random variables with densities p_ξ, p_η .

Convolution

Theorem

Let ξ, η be independent absolutely continuous random variables with densities p_ξ, p_η .

Then

$$F_{\xi+\eta}(x) = \int_{\mathbb{R}} F_\xi(x-u)p_\eta(u)du,$$

Convolution

Theorem

Let ξ, η be independent absolutely continuous random variables with densities p_ξ, p_η .

Then

$$F_{\xi+\eta}(x) = \int_{\mathbb{R}} F_{\xi}(x-u)p_{\eta}(u)du,$$

$$p_{\xi+\eta}(x) = \int_{\mathbb{R}} p_{\xi}(x-u)p_{\eta}(u)du.$$

The proof

An example