

# Probability theory

## Lecture 1: Probability space

Maksim Zhukovskii

MIPT

## Heads or tails?

The probability of 'Head' and the probability of 'Tail' equals  $1/2$ .



## Heads or tails?

The probability of 'Head' and the probability of 'Tail' equals  $1/2$ .



1. —  $H$  represents the event '**H**ead'  
—  $T$  represents the event '**T**ail'

## Heads or tails?

The probability of 'Head' and the probability of 'Tail' equals  $1/2$ .



1. —  $H$  represents the event '**H**ead'  
—  $T$  represents the event '**T**ail'  
 $P(H) = P(T) = 1/2$

## Heads or tails?

The probability of 'Head' and the probability of 'Tail' equals  $1/2$ .



1. —  $H$  represents the event '**H**ead'  
—  $T$  represents the event '**T**ail'

$$P(H) = P(T) = 1/2$$

2.  $X$  tails in  $n$  independent coin tosses

## Heads or tails?

The probability of 'Head' and the probability of 'Tail' equals  $1/2$ .



1. —  $H$  represents the event '**H**ead'  
—  $T$  represents the event '**T**ail'

$$P(H) = P(T) = 1/2$$

2.  $X$  tails in  $n$  independent coin tosses  
 $X/n \approx 1/2$

## Sample space and events

Informal definition:

*sample space* — the set of all possible outcomes of an experiment

## Sample space and events

Informal definition:

*sample space* — the set of all possible outcomes of an experiment

Formal definition:



## Sample space and events

### Informal definition:

*sample space* — the set of all possible outcomes of an experiment

### Formal definition:

- *sample space*  $\Omega$  — abstract set

## Sample space and events

### Informal definition:

*sample space* — the set of all possible outcomes of an experiment

### Formal definition:

- *sample space*  $\Omega$  — abstract set
- $\omega \in \Omega$  — *elementary event*

## Sample space and events

### Informal definition:

*sample space* — the set of all possible outcomes of an experiment

### Formal definition:

- *sample space*  $\Omega$  — abstract set
- $\omega \in \Omega$  — *elementary event*
- $A \subseteq \Omega$  — *an event*

## Examples

Tossing a coin:

$$\Omega = \{H, T\}$$

## Examples

Tossing a coin:

$$\Omega = \{H, T\}$$

Tossing a dice:

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$



## Examples

Tossing a coin:

$$\Omega = \{H, T\}$$



Tossing a dice:

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$

$$A = \{2, 4, 6\} \text{ — 'outcome is even'}$$

## Classical probability

Find the probability that a dice shows an even number.

## Classical probability

Find the probability that a dice shows an even number.

**Solution:**  $\frac{\text{the number of 'even positions'}}{\text{the number of all positions}} = \frac{3}{6} = \frac{1}{2}$



## Classical probability

Find the probability that a dice shows an even number.

**Solution:**  $\frac{\text{the number of 'even positions'}}{\text{the number of all positions}} = \frac{3}{6} = \frac{1}{2}$

- ▶  $\Omega = \{\omega_1, \dots, \omega_n\}$
- ▶  $A \subseteq \Omega \Rightarrow P(A) = \frac{|A|}{n}$ .

## Several combinatorial examples

There are 4 red and 5 blue balls in a box.  
3 balls are chosen without repetition.  
 $P(\text{all these balls are blue}) = ?$

## Several combinatorial examples

There are 4 red and 5 blue balls in a box.  
3 balls are chosen without repetition.  
 $P(\text{all these balls are blue}) = ?$

Solution:

$$\Omega = \{(x_1, x_2, x_3) : x_i \in [9] \text{ distinct}\},$$

## Several combinatorial examples

There are 4 red and 5 blue balls in a box.  
3 balls are chosen without repetition.  
 $P(\text{all these balls are blue}) = ?$

Solution:

$$\Omega = \{(x_1, x_2, x_3) : x_i \in [9] \text{ distinct}\},$$

$$A = \{(x_1, x_2, x_3) : x_i \in [9] \setminus [4]\} \subset \Omega.$$

## Several combinatorial examples

There are 4 red and 5 blue balls in a box.  
3 balls are chosen without repetition.  
 $P(\text{all these balls are blue}) = ?$

Solution:

$$\Omega = \{(x_1, x_2, x_3) : x_i \in [9] \text{ distinct}\},$$

$$A = \{(x_1, x_2, x_3) : x_i \in [9] \setminus [4]\} \subset \Omega.$$

$$\text{Then } P(A) = |A|/|\Omega| = \binom{5}{3} / \binom{9}{3} = \frac{5}{42}.$$

## Several combinatorial examples

Throwing  $n$  coins independently,  
what is the probability of having exactly  $k$   
tails?

## Several combinatorial examples

Throwing  $n$  coins independently,  
what is the probability of having exactly  $k$   
tails?

Solution:

$$\Omega = \{0, 1\}^n,$$

## Several combinatorial examples

Throwing  $n$  coins independently,  
what is the probability of having exactly  $k$   
tails?

Solution:

$$\Omega = \{0, 1\}^n,$$

$$A = \{(x_1, \dots, x_n) : \sum x_i = k\} \subset \Omega.$$



## Several combinatorial examples

Throwing  $n$  coins independently,  
what is the probability of having exactly  $k$   
tails?

Solution:

$$\Omega = \{0, 1\}^n,$$

$$A = \{(x_1, \dots, x_n) : \sum x_i = k\} \subset \Omega.$$

$$\text{Then } P(A) = |A|/|\Omega| = \binom{n}{k}/2^n.$$

## Probability space

*A probability space  $(\Omega, \mathcal{F}, P)$ :*

## Probability space

*A probability space  $(\Omega, \mathcal{F}, P)$ :*

- ▶  $\Omega$  — *a sample space* (an arbitrary set);

## Probability space

*A probability space  $(\Omega, \mathcal{F}, P)$ :*

- ▶  $\Omega$  — *a sample space (an arbitrary set);*
- ▶  $\mathcal{F} \subset 2^\Omega$  — *a set of events;*

## Probability space

*A probability space  $(\Omega, \mathcal{F}, P)$ :*

- ▶  $\Omega$  — *a sample space (an arbitrary set);*
- ▶  $\mathcal{F} \subset 2^\Omega$  — *a set of events;*
- ▶  $P : \mathcal{F} \rightarrow [0, 1]$  — *a probability measure (or, simply, probability).*

## Probability space: events

$\mathcal{F}$  —  $\sigma$ -algebra:

## Probability space: events

$\mathcal{F}$  —  $\sigma$ -algebra:

- ▶  $\Omega \in \mathcal{F}$ ,

## Probability space: events

$\mathcal{F}$  —  $\sigma$ -algebra:

- ▶  $\Omega \in \mathcal{F}$ ,
- ▶  $A \in \mathcal{F} \Rightarrow \overline{A} \in \mathcal{F}$ ,



## Probability space: events

$\mathcal{F}$  —  $\sigma$ -algebra:

- ▶  $\Omega \in \mathcal{F}$ ,
- ▶  $A \in \mathcal{F} \Rightarrow \overline{A} \in \mathcal{F}$ ,
- ▶  $A_1, A_2, \dots \in \mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}$ .

## Probability space: events

$\mathcal{F}$  —  $\sigma$ -algebra:

- ▶  $\Omega \in \mathcal{F}$ ,
- ▶  $A \in \mathcal{F} \Rightarrow \bar{A} \in \mathcal{F}$ ,
- ▶  $A_1, A_2, \dots \in \mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}$ .

$A$  is *an event* if  $A \in \mathcal{F}$

## Probability space: probability measure

$P : \mathcal{F} \rightarrow [0, 1]$  is a *probability measure*, if

## Probability space: probability measure

$P : \mathcal{F} \rightarrow [0, 1]$  is a *probability measure*, if

- ▶  $P(\Omega) = 1,$

## Probability space: probability measure

$P : \mathcal{F} \rightarrow [0, 1]$  is a *probability measure*, if

- ▶  $P(\Omega) = 1$ ,
- ▶  $A_1, A_2, \dots \in \mathcal{F}$  are disjoint  $\Rightarrow$   
 $P(A_1 \sqcup A_2 \sqcup \dots) = \sum_{i=1}^{\infty} P(A_i)$ .

## Special $\sigma$ -algebras

- ▶ There are two trivial  $\sigma$ -algebras:

## Special $\sigma$ -algebras

- ▶ There are two trivial  $\sigma$ -algebras:  
 $\mathcal{F}_0 = \{\emptyset, \Omega\},$

## Special $\sigma$ -algebras

- There are two trivial  $\sigma$ -algebras:

$$\mathcal{F}_0 = \{\emptyset, \Omega\}, \mathcal{F} = 2^\Omega$$



## Special $\sigma$ -algebras

- ▶ There are two trivial  $\sigma$ -algebras:

$$\mathcal{F}_0 = \{\emptyset, \Omega\}, \mathcal{F} = 2^\Omega$$

- ▶ Let  $A \subset \Omega$

## Special $\sigma$ -algebras

- ▶ There are two trivial  $\sigma$ -algebras:

$$\mathcal{F}_0 = \{\emptyset, \Omega\}, \mathcal{F} = 2^\Omega$$

- ▶ Let  $A \subset \Omega$

$$\mathcal{F}_A := \{\emptyset, \Omega, A, \overline{A}\}$$

## Minimum $\sigma$ -algebras

$$\Sigma \subset 2^\Omega$$

## Minimum $\sigma$ -algebras

$$\Sigma \subset 2^\Omega$$

$\sigma(\Sigma)$  — inclusion-minimum  $\sigma$ -algebra  
containing all sets from  $\Sigma$

## Minimum $\sigma$ -algebras

$$\Sigma \subset 2^\Omega$$

$\sigma(\Sigma)$  — inclusion-minimum  $\sigma$ -algebra  
containing all sets from  $\Sigma$

### Examples

►  $\sigma(\{A\}) = \mathcal{F}_A = \{\emptyset, \Omega, A, \bar{A}\}.$

## Minimum $\sigma$ -algebras

$$\Sigma \subset 2^\Omega$$

$\sigma(\Sigma)$  — inclusion-minimum  $\sigma$ -algebra  
containing all sets from  $\Sigma$

### Examples

- ▶  $\sigma(\{A\}) = \mathcal{F}_A = \{\emptyset, \Omega, A, \bar{A}\}.$
- ▶  $\Omega = A_1 \sqcup A_2 \sqcup \dots,$

## Minimum $\sigma$ -algebras

$$\Sigma \subset 2^\Omega$$

$\sigma(\Sigma)$  — inclusion-minimum  $\sigma$ -algebra  
containing all sets from  $\Sigma$

### Examples

- ▶  $\sigma(\{A\}) = \mathcal{F}_A = \{\emptyset, \Omega, A, \bar{A}\}.$
- ▶  $\Omega = A_1 \sqcup A_2 \sqcup \dots,$   
 $\sigma(\{A_1, A_2, \dots\}) = \{\sqcup_{j \in \mathcal{J}} A_j, \mathcal{J} \subseteq \mathbb{N}\}.$

## Minimum $\sigma$ -algebras

$$\Sigma \subset 2^\Omega$$

$\sigma(\Sigma)$  — inclusion-minimum  $\sigma$ -algebra containing all sets from  $\Sigma$

### Examples

- ▶  $\sigma(\{A\}) = \mathcal{F}_A = \{\emptyset, \Omega, A, \bar{A}\}.$
- ▶  $\Omega = A_1 \sqcup A_2 \sqcup \dots,$   
 $\sigma(\{A_1, A_2, \dots\}) = \{\sqcup_{j \in \mathcal{J}} A_j, \mathcal{J} \subseteq \mathbb{N}\}.$
- ▶ The **Borel  $\sigma$ -algebra** on  $\mathbb{R}$ :  
 $\mathcal{B}(\mathbb{R}) = \sigma(\{(a, b), -\infty \leq a < b \leq \infty\}).$



## Existence of $\sigma(\Sigma)$

*Proof.*

## Existence of $\sigma(\Sigma)$

*Proof.*

1. Existence of a  $\sigma$ -algebra:

## Existence of $\sigma(\Sigma)$

*Proof.*

1. Existence of a  $\sigma$ -algebra:

$$2^{\Omega} \supset \Sigma$$

## Existence of $\sigma(\Sigma)$

*Proof.*

1. Existence of a  $\sigma$ -algebra:

$$2^{\Omega} \supset \Sigma$$

2. Construction of the minimum  $\sigma$ -algebra:

## Existence of $\sigma(\Sigma)$

*Proof.*

1. Existence of a  $\sigma$ -algebra:

$$2^{\Omega} \supset \Sigma$$

2. Construction of the minimum  $\sigma$ -algebra:

$$\mathcal{F}_* := \bigcap_{\mathcal{F} \supseteq \Sigma \text{ — } \sigma\text{-algebra}} \mathcal{F}$$

## Existence of $\sigma(\Sigma)$

3.  $\mathcal{F}_*$  is  $\sigma$ -algebra:

## Existence of $\sigma(\Sigma)$

3.  $\mathcal{F}_*$  is  $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F}$

## Existence of $\sigma(\Sigma)$

3.  $\mathcal{F}_*$  is  $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;



## Existence of $\sigma(\Sigma)$

3.  $\mathcal{F}_*$  is  $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_*$

## Existence of $\sigma(\Sigma)$

3.  $\mathcal{F}_*$  is  $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F}$

## Existence of $\sigma(\Sigma)$

3.  $\mathcal{F}_*$  is  $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\overline{A}$

## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$

## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$
- $A_1, A_2, \dots \in \mathcal{F}_*$

## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$
- $A_1, A_2, \dots \in \mathcal{F}_* \Rightarrow A_1, A_2, \dots \in \mathcal{F}$   
for every  $\mathcal{F}$

## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$
- $A_1, A_2, \dots \in \mathcal{F}_* \Rightarrow A_1, A_2, \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}$   
for every  $\mathcal{F}$

## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$
- $A_1, A_2, \dots \in \mathcal{F}_* \Rightarrow A_1, A_2, \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}_*$



## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$
- $A_1, A_2, \dots \in \mathcal{F}_* \Rightarrow A_1, A_2, \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}_*$

### 4. $\mathcal{F}_*$ is minimum:

## Existence of $\sigma(\Sigma)$

### 3. $\mathcal{F}_*$ is $\sigma$ -algebra:

- $\Omega \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow \Omega \in \mathcal{F}_*$ ;
- $A \in \mathcal{F}_* \Rightarrow A \in \mathcal{F}$  for every  $\mathcal{F} \Rightarrow$   
the same is true for  $\bar{A} \Rightarrow \bar{A} \in \mathcal{F}_*$
- $A_1, A_2, \dots \in \mathcal{F}_* \Rightarrow A_1, A_2, \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}$   
for every  $\mathcal{F} \Rightarrow A_1 \cup A_2 \cup \dots \in \mathcal{F}_*$

### 4. $\mathcal{F}_*$ is minimum:

If  $\mathcal{F} \supseteq \Sigma$  —  $\sigma$ -algebra, then  $\mathcal{F}_* \subseteq \mathcal{F}$ .

## Special probability measures on finite $\Omega$

- ▶ **Classical probability:**

$$\Omega = \{\omega_1, \dots, \omega_n\}, \mathcal{F} = 2^\Omega, P(A) = \frac{|A|}{n}$$

## Special probability measures on finite $\Omega$

- ▶ **Classical probability:**

$$\Omega = \{\omega_1, \dots, \omega_n\}, \mathcal{F} = 2^\Omega, P(A) = \frac{|A|}{n}$$

In finite cases, it is sufficient to define  $P$  on unit sets

## Special probability measures on finite $\Omega$

- ▶ **Classical probability:**

$$\Omega = \{\omega_1, \dots, \omega_n\}, \mathcal{F} = 2^\Omega, P(A) = \frac{|A|}{n}$$

In finite cases, it is sufficient to define  $P$  on unit sets

- ▶ **Bernoulli scheme:**

$$\Omega = \{0, 1\}^n, \mathcal{F} = 2^\Omega, \\ P(\{(x_1, \dots, x_n)\}) = p^{\sum x_i} (1 - p)^{n - \sum x_i}$$

## Geometric probability

$\mu$  — the Lebesgue measure on  $\mathbb{R}^n$

## Geometric probability

$\mu$  — the Lebesgue measure on  $\mathbb{R}^n$

$$\mathcal{B}(\mathbb{R}^n) := \sigma(B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R}))$$

## Geometric probability

$\mu$  — the Lebesgue measure on  $\mathbb{R}^n$

$$\mathcal{B}(\mathbb{R}^n) := \sigma(B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R}))$$

$\Omega \in \mathcal{B}(\mathbb{R}^n)$  such that  $\mu(\Omega) < \infty$



## Geometric probability

$\mu$  — the Lebesgue measure on  $\mathbb{R}^n$

$$\mathcal{B}(\mathbb{R}^n) := \sigma(B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R}))$$

$\Omega \in \mathcal{B}(\mathbb{R}^n)$  such that  $\mu(\Omega) < \infty$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}^n) \cap \Omega$$

## Geometric probability

$\mu$  — the Lebesgue measure on  $\mathbb{R}^n$

$$\mathcal{B}(\mathbb{R}^n) := \sigma(B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R}))$$

$\Omega \in \mathcal{B}(\mathbb{R}^n)$  such that  $\mu(\Omega) < \infty$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}^n) \cap \Omega$$

$$P(A) = \frac{\mu(A)}{\mu(\Omega)}$$

## Chance of meeting in a bar

Alice and Bob decided to meet in a bar on a given day.

Each one has to come in the bar at a random time between 12:00 and 13:00 and wait for another one for 15 minutes.

What is the probability that they will meet?


## Chance of meeting in a bar

Solution:

$$\Omega = [0, 1]^2 \subset \mathbb{R}^2,$$

$A$  is colored black,

$$P(A) = \frac{\mu(A)}{\mu(\Omega)} = \frac{1 - 2 \cdot \frac{9}{32}}{1} = \frac{7}{16}.$$



## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Binomial random graph

—  $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$ ,

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Binomial random graph

- $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$ ,
- $\mathcal{F} = 2^\Omega$ ,

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Binomial random graph

- $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$ ,
- $\mathcal{F} = 2^\Omega$ ,
- $P(\{G = (V, E)\}) = p^{|E|}(1 - p)^{\binom{n}{2} - |E|}$ .



## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Binomial random graph

- $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$ ,
- $\mathcal{F} = 2^\Omega$ ,
- $P(\{G = (V, E)\}) = p^{|E|}(1 - p)^{\binom{n}{2} - |E|}$ .

Every edge appears *independently* with probability  $p$

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Uniform random graph:

—  $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$  with exactly  $m$  edges,

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Uniform random graph:

- $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$  with exactly  $m$  edges,
- $\mathcal{F} = 2^\Omega$ ,

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Uniform random graph:

- $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$  with exactly  $m$  edges,
- $\mathcal{F} = 2^\Omega$ ,
- $P(G) = 1/\binom{\binom{n}{2}}{m}$ .

## Random graphs

Consider the vertex set  $V = \{1, \dots, n\}$

### Uniform random graph:

- $\Omega$  — the set of all undirected graphs without loops and multiple edges on  $V$  with exactly  $m$  edges,
- $\mathcal{F} = 2^\Omega$ ,
- $P(G) = 1/\binom{n}{m}_2$ .

Classical probability

# Properties of probability

1.  $P(\emptyset) = 0$

# Properties of probability

1.  $P(\emptyset) = 0$

*Proof.*

$$\emptyset = \emptyset \sqcup \emptyset$$



# Properties of probability

1.  $P(\emptyset) = 0$

*Proof.*

$$\emptyset = \emptyset \sqcup \emptyset \Rightarrow$$

$$P(\emptyset) = P(\emptyset) + P(\emptyset)$$

# Properties of probability

1.  $P(\emptyset) = 0$

*Proof.*

$$\emptyset = \emptyset \sqcup \emptyset \Rightarrow$$

$$P(\emptyset) = P(\emptyset) + P(\emptyset) \Rightarrow$$

$$P(\emptyset) = 0 \quad \square$$

## Properties of probability

$$2. P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

## Properties of probability

$$2. P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

*Proof.*

$$A \cup B = (A \setminus B) \sqcup B$$

## Properties of probability

$$2. P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

*Proof.*

$$A \cup B = (A \setminus B) \sqcup B \Rightarrow$$

$$P(A \cup B) = P(A \setminus B) + P(B);$$

## Properties of probability

$$2. P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

*Proof.*

$$A \cup B = (A \setminus B) \sqcup B \Rightarrow$$

$$P(A \cup B) = P(A \setminus B) + P(B);$$

$$(A \setminus B) \sqcup (A \cap B) = A$$

## Properties of probability

$$2. P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

*Proof.*

$$A \cup B = (A \setminus B) \sqcup B \Rightarrow$$

$$P(A \cup B) = P(A \setminus B) + P(B);$$

$$(A \setminus B) \sqcup (A \cap B) = A \Rightarrow$$

$$P(A \setminus B) = P(A) - P(A \cap B) \quad \square$$

## Properties of probability

$$3. B \subset A \Rightarrow P(B) \leq P(A)$$



## Properties of probability

$$3. B \subset A \Rightarrow P(B) \leq P(A)$$

*Proof.*

$$A = (A \setminus B) \sqcup B$$

## Properties of probability

$$3. B \subset A \Rightarrow P(B) \leq P(A)$$

*Proof.*

$$A = (A \setminus B) \sqcup B \Rightarrow$$

$$P(A) = P(A \setminus B) + P(B) \geq P(B) \quad \square$$

## Properties of probability

$$4. P(A_1 \cup A_2 \cup \dots) \leq P(A_1) + P(A_2) + \dots$$

## Properties of probability

$$4. P(A_1 \cup A_2 \cup \dots) \leq P(A_1) + P(A_2) + \dots$$

*Proof.*

$$B_i = A_i \setminus (A_1 \cup \dots \cup A_{i-1}) \subseteq A_i$$

## Properties of probability

$$4. P(A_1 \cup A_2 \cup \dots) \leq P(A_1) + P(A_2) + \dots$$

*Proof.*

$$B_i = A_i \setminus (A_1 \cup \dots \cup A_{i-1}) \subseteq A_i$$

$$P(A_1 \cup A_2 \cup \dots) = P(B_1 \cup B_2 \cup \dots) =$$

## Properties of probability

$$4. P(A_1 \cup A_2 \cup \dots) \leq P(A_1) + P(A_2) + \dots$$

*Proof.*

$$B_i = A_i \setminus (A_1 \cup \dots \cup A_{i-1}) \subseteq A_i$$

$$P(A_1 \cup A_2 \cup \dots) = P(B_1 \cup B_2 \cup \dots) = P(B_1) + P(B_2) + \dots \leq$$

## Properties of probability

$$4. P(A_1 \cup A_2 \cup \dots) \leq P(A_1) + P(A_2) + \dots$$

*Proof.*

$$B_i = A_i \setminus (A_1 \cup \dots \cup A_{i-1}) \subseteq A_i$$

$$\begin{aligned} P(A_1 \cup A_2 \cup \dots) &= P(B_1 \cup B_2 \cup \dots) = \\ &P(B_1) + P(B_2) + \dots \leq \\ &P(A_1) + P(A_2) + \dots \quad \square \end{aligned}$$

## Continuity of probability

Let  $\mathcal{F}$  be a  $\sigma$ -algebra of subsets of  $\Omega$ ,  
 $P : \mathcal{F} \rightarrow [0, 1]$  be such that

- ▶  $P(\Omega) = 1$ ,
- ▶  $P(A \sqcup B) = P(A) + P(B)$  whenever  $A, B \in \mathcal{F}$  are disjoint.



## Continuity of probability

The following four properties are equivalent:

## Continuity of probability

The following four properties are equivalent:

1.  $P$  — probability;

## Continuity of probability

The following four properties are equivalent:

1.  $P$  — probability;
2. for every sequence  $A_n \in \mathcal{F}$  such that  $A_n \uparrow A$  ( $A_n \subseteq A_{n+1}$ ,  $A = \bigcup A_n$ ),  
 $P(A) = \lim_{n \rightarrow \infty} P(A_n)$ ;

## Continuity of probability

The following four properties are equivalent:

1.  $P$  — probability;
2. for every sequence  $A_n \in \mathcal{F}$  such that  $A_n \uparrow A$  ( $A_n \subseteq A_{n+1}$ ,  $A = \bigcup A_n$ ),  
 $P(A) = \lim_{n \rightarrow \infty} P(A_n)$ ;
3. for every sequence  $A_n \in \mathcal{F}$  such that  $A_n \downarrow A$  ( $A_n \supseteq A_{n+1}$ ,  $A = \bigcap A_n$ ),  
 $P(A) = \lim_{n \rightarrow \infty} P(A_n)$ ;

## Continuity of probability

The following four properties are equivalent:

1.  $P$  — probability;
2. for every sequence  $A_n \in \mathcal{F}$  such that  $A_n \uparrow A$  ( $A_n \subseteq A_{n+1}$ ,  $A = \bigcup A_n$ ),  
 $P(A) = \lim_{n \rightarrow \infty} P(A_n)$ ;
3. for every sequence  $A_n \in \mathcal{F}$  such that  $A_n \downarrow A$  ( $A_n \supseteq A_{n+1}$ ,  $A = \bigcap A_n$ ),  
 $P(A) = \lim_{n \rightarrow \infty} P(A_n)$ ;
4. for every sequence  $A_n \in \mathcal{F}$  such that  $A_n \downarrow \emptyset$ ,  $\lim_{n \rightarrow \infty} P(A_n) = 0$ .

## The proof

1) $\Rightarrow$ 2)

$$B_n = A_n \setminus A_{n-1}, A_0 = \emptyset.$$

## The proof

1) $\Rightarrow$ 2)

$$B_n = A_n \setminus A_{n-1}, A_0 = \emptyset.$$

$$B_n \in \mathcal{F} \text{ are disjoint and } \bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} B_i$$

## The proof

1) $\Rightarrow$ 2)

$$B_n = A_n \setminus A_{n-1}, A_0 = \emptyset.$$

$$B_n \in \mathcal{F} \text{ are disjoint and } \bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} B_i$$

$$P(A) = \sum_{i=1}^{\infty} P(B_i) = \sum_{i=1}^{\infty} P(A_i - A_{i-1}) =$$

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n (P(A_i) - P(A_{i-1})) = \lim_{n \rightarrow \infty} P(A_n).$$



## The proof

2) $\Rightarrow$ 3)

$$P\left(\bigcap_{n=1}^{\infty} A_n\right) = P\left(\overline{\bigcap_{n=1}^{\infty} \bar{A}_n}\right) = 1 - P\left(\bigcup_{n=1}^{\infty} \bar{A}_n\right);$$

## The proof

2) $\Rightarrow$ 3)

$$P\left(\bigcap_{n=1}^{\infty} A_n\right) = P\left(\overline{\bigcap_{n=1}^{\infty} \bar{A}_n}\right) = 1 - P\left(\bigcup_{n=1}^{\infty} \bar{A}_n\right);$$

$$P\left(\bigcup_{n=1}^{\infty} \bar{A}_n\right) = \lim_{n \rightarrow \infty} P(\bar{A}_n) = \\ \lim_{n \rightarrow \infty} (1 - P(A_n)) = 1 - \lim_{n \rightarrow \infty} P(A_n).$$

## The proof

3) $\Rightarrow$ 4) because 4) is special case of 3).

## The proof

4) $\Rightarrow$ 1)

$$A_1, A_2, \dots \in \mathcal{F}, \bigcup_{i=1}^{\infty} A_i = A,$$

## The proof

4) $\Rightarrow$ 1)

$$A_1, A_2, \dots \in \mathcal{F}, \quad \bigsqcup_{i=1}^{\infty} A_i = A, \quad B_n = \bigcup_{i=n}^{\infty} A_i$$

## The proof

4) $\Rightarrow$ 1)

$$A_1, A_2, \dots \in \mathcal{F}, \quad \bigsqcup_{i=1}^{\infty} A_i = A, \quad B_n = \bigcup_{i=n}^{\infty} A_i$$

$$\blacktriangleright B_n = A \setminus (A_1 \cup \dots \cup A_{n-1}) \in \mathcal{F},$$

$$\blacktriangleright B_n \supseteq B_{n+1}, \quad \bigcap_{i=1}^{\infty} B_i = \emptyset.$$

## The proof

4) $\Rightarrow$ 1)

$$A_1, A_2, \dots \in \mathcal{F}, \quad \bigsqcup_{i=1}^{\infty} A_i = A, \quad B_n = \bigcup_{i=n}^{\infty} A_i$$

$$\blacktriangleright B_n = A \setminus (A_1 \cup \dots \cup A_{n-1}) \in \mathcal{F},$$

$$\blacktriangleright B_n \supseteq B_{n+1}, \quad \bigcap_{i=1}^{\infty} B_i = \emptyset.$$

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} P(B_n) = \lim_{n \rightarrow \infty} \left( P(A) - P\left(\bigcup_{i=1}^{n-1} A_i\right) \right) \\ &= P(A) - \sum_{i=1}^{\infty} P(A_i). \end{aligned}$$