

# Probability theory

## Lecture 7: Convergences

Maksim Zhukovskii

MIPT

## Almost sure convergence

$\xi_1, \xi_2, \dots$

converges almost surely (with prob. 1) to  $\xi$ ,

## Almost sure convergence

$\xi_1, \xi_2, \dots$

converges almost surely (with prob. 1) to  $\xi$ ,

if

$$P(\xi_n \rightarrow \xi) = 1.$$

## Almost sure convergence

$\xi_1, \xi_2, \dots$

converges almost surely (with prob. 1) to  $\xi$ ,

if

$$P(\xi_n \rightarrow \xi) = 1.$$

$$\xi_n \xrightarrow{\text{a.s.}} \xi$$

## Convergence in probability

$\xi_1, \xi_2, \dots$

converges in probability to  $\xi$ ,

## Convergence in probability

$\xi_1, \xi_2, \dots$

converges in probability to  $\xi$ ,

if

for every  $\varepsilon > 0$ ,  $\lim_{n \rightarrow \infty} P(|\xi_n - \xi| \geq \varepsilon) = 0$ .

## Convergence in probability

$\xi_1, \xi_2, \dots$

converges in probability to  $\xi$ ,

if

for every  $\varepsilon > 0$ ,  $\lim_{n \rightarrow \infty} P(|\xi_n - \xi| \geq \varepsilon) = 0$ .

$$\xi_n \xrightarrow{P} \xi$$

## Convergence in probability

$\xi_1, \xi_2, \dots$

converges in probability to  $\xi$ ,

if

for every  $\varepsilon > 0$ ,  $\lim_{n \rightarrow \infty} P(|\xi_n - \xi| \geq \varepsilon) = 0$ .

$$\xi_n \xrightarrow{P} \xi$$

► the law of large numbers:

$$\frac{S_n - ES_n}{n} \xrightarrow{P} 0$$



## Convergence in $L^p$

Let  $p > 0$ .

## Convergence in $L^p$

Let  $p > 0$ .

$\xi_1, \xi_2, \dots$

converges in the  $L^p$ -norm to  $\xi$ ,

## Convergence in $L^p$

Let  $p > 0$ .

$\xi_1, \xi_2, \dots$

converges in the  $L^p$ -norm to  $\xi$ ,

if

$$E\xi^p, E\xi_i^p < \infty, \quad \lim_{n \rightarrow \infty} E|\xi_n - \xi|^p = 0.$$

## Convergence in $L^p$

Let  $p > 0$ .

$\xi_1, \xi_2, \dots$

converges in the  $L^p$ -norm to  $\xi$ ,

if

$$E\xi^p, E\xi_i^p < \infty, \quad \lim_{n \rightarrow \infty} E|\xi_n - \xi|^p = 0.$$

$$\xi_n \xrightarrow{L_p} \xi$$

## Convergence in $L^p$

Let  $p > 0$ .

$\xi_1, \xi_2, \dots$

converges in the  $L^p$ -norm to  $\xi$ ,

if

$$E\xi^p, E\xi_i^p < \infty, \quad \lim_{n \rightarrow \infty} E|\xi_n - \xi|^p = 0.$$

$$\xi_n \xrightarrow{L_p} \xi$$

If  $p = 1$ , we say that

$\xi_1, \xi_2, \dots$  converges in mean to  $\xi$ .

## Convergence in distribution

$\xi_1, \xi_2, \dots$  converges in distribution  
(converges weakly) to  $\xi$ ,

## Convergence in distribution

$\xi_1, \xi_2, \dots$  converges in distribution  
(converges weakly) to  $\xi$ ,

if

for every  $x \in \mathbb{R}$  at which  $F_\xi$  is continuous,  
$$\lim_{n \rightarrow \infty} F_{\xi_n}(x) = F_\xi(x).$$

## Convergence in distribution

$\xi_1, \xi_2, \dots$  converges in distribution  
(converges weakly) to  $\xi$ ,

if

for every  $x \in \mathbb{R}$  at which  $F_\xi$  is continuous,  
$$\lim_{n \rightarrow \infty} F_{\xi_n}(x) = F_\xi(x).$$

$$\xi_n \xrightarrow{d} \xi$$



## Convergence in distribution

$\xi_1, \xi_2, \dots$  converges in distribution  
(converges weakly) to  $\xi$ ,

if

for every  $x \in \mathbb{R}$  at which  $F_\xi$  is continuous,  
$$\lim_{n \rightarrow \infty} F_{\xi_n}(x) = F_\xi(x).$$

$$\xi_n \xrightarrow{d} \xi$$

Equivalent definition:

for every continuous bounded function  
 $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $\lim_{n \rightarrow \infty} \mathbb{E}f(\xi_n) = \mathbb{E}f(\xi).$

# Relations

## Theorem

$$1) \xi_n \xrightarrow{a.s.} \xi \text{ implies } \xi_n \xrightarrow{P} \xi;$$

# Relations

## Theorem

$$1) \xi_n \xrightarrow{a.s.} \xi \text{ implies } \xi_n \xrightarrow{P} \xi;$$

$$2) \xi_n \xrightarrow{P} \xi \text{ implies } \xi_n \xrightarrow{d} \xi;$$

## Relations

### Theorem

1)  $\xi_n \xrightarrow{a.s.} \xi$  implies  $\xi_n \xrightarrow{P} \xi$ ;

2)  $\xi_n \xrightarrow{P} \xi$  implies  $\xi_n \xrightarrow{d} \xi$ ;

3) for every  $p > 0$ ,  $\xi_n \xrightarrow{L_p} \xi$  implies  $\xi_n \xrightarrow{P} \xi$ .

# Relations

## Theorem

1)  $\xi_n \xrightarrow{a.s.} \xi$  implies  $\xi_n \xrightarrow{P} \xi$ ;

2)  $\xi_n \xrightarrow{P} \xi$  implies  $\xi_n \xrightarrow{d} \xi$ ;

3) for every  $p > 0$ ,  $\xi_n \xrightarrow{L_p} \xi$  implies  $\xi_n \xrightarrow{P} \xi$ .

Almost Sure  
Convergence

Mean Convergence

Convergence in  
Probability

Convergence in  
Distribution



## The proof of 1)

$$\text{a.s.} \Rightarrow P$$

## The proof of 1)

a.s.  $\Rightarrow$  P

Let  $\xi_n \xrightarrow{\text{a.s.}} \xi, \varepsilon > 0$ .

## The proof of 1)

a.s.  $\Rightarrow$  P

Let  $\xi_n \xrightarrow{\text{a.s.}} \xi$ ,  $\varepsilon > 0$ .

$$A_n = \{\omega : |\xi_n(\omega) - \xi(\omega)| < \varepsilon\},$$

$$B_n = \{\omega : \forall k \geq n \ |\xi_k(\omega) - \xi(\omega)| < \varepsilon\} \subset A_n.$$



## The proof of 1)

a.s.  $\Rightarrow$  P

Let  $\xi_n \xrightarrow{\text{a.s.}} \xi$ ,  $\varepsilon > 0$ .

$$A_n = \{\omega : |\xi_n(\omega) - \xi(\omega)| < \varepsilon\},$$

$$B_n = \{\omega : \forall k \geq n \ |\xi_k(\omega) - \xi(\omega)| < \varepsilon\} \subset A_n.$$

It remains to prove that  $P(B_n) \rightarrow 1$ ,  $n \rightarrow \infty$ .

## The proof of 1)

$$\text{a.s.} \Rightarrow P$$

## The proof of 1)

a.s.  $\Rightarrow$  P

$$A := \{\omega : \exists n \in \mathbb{N} \forall k \geq n |\xi_k(\omega) - \xi(\omega)| < \varepsilon\},$$

## The proof of 1)

a.s.  $\Rightarrow$  P

$$A := \{\omega : \exists n \in \mathbb{N} \forall k \geq n |\xi_k(\omega) - \xi(\omega)| < \varepsilon\},$$

$$\{\xi_n \rightarrow \xi\} \subset A,$$

## The proof of 1)

a.s.  $\Rightarrow$  P

$$A := \{\omega : \exists n \in \mathbb{N} \forall k \geq n |\xi_k(\omega) - \xi(\omega)| < \varepsilon\},$$

$$\{\xi_n \rightarrow \xi\} \subset A,$$

$$P(A) = 1,$$

## The proof of 1)

a.s.  $\Rightarrow$  P

$$A := \{\omega : \exists n \in \mathbb{N} \forall k \geq n |\xi_k(\omega) - \xi(\omega)| < \varepsilon\},$$

$$\{\xi_n \rightarrow \xi\} \subset A,$$

$$P(A) = 1,$$

Clearly,  $B_n \uparrow A$ .

## The proof of 1)

a.s.  $\Rightarrow$  P

$$A := \{\omega : \exists n \in \mathbb{N} \forall k \geq n |\xi_k(\omega) - \xi(\omega)| < \varepsilon\},$$

$$\{\xi_n \rightarrow \xi\} \subset A,$$

$$P(A) = 1,$$

Clearly,  $B_n \uparrow A$ .

Therefore,  $P(B_n) \rightarrow 1, n \rightarrow \infty$ .

## The proof of 2)

$$P \Rightarrow d$$



## The proof of 2)

$$P \Rightarrow d$$

$f : \mathbb{R} \rightarrow \mathbb{R}$  — continuous,  $|f| \leq C$ ,  $\varepsilon > 0$ .

## The proof of 2)

$$P \Rightarrow d$$

$f : \mathbb{R} \rightarrow \mathbb{R}$  — continuous,  $|f| \leq C$ ,  $\varepsilon > 0$ .

Let  $\tilde{\xi}_n = \xi I(\xi \leq n)$ .

## The proof of 2)

$$P \Rightarrow d$$

$f : \mathbb{R} \rightarrow \mathbb{R}$  — continuous,  $|f| \leq C$ ,  $\varepsilon > 0$ .

Let  $\tilde{\xi}_n = \xi I(\xi \leq n)$ .

Obviously,  $\tilde{\xi}_n \xrightarrow{\text{a.s.}} \xi$ .

## The proof of 2)

$$P \Rightarrow d$$

$f : \mathbb{R} \rightarrow \mathbb{R}$  — continuous,  $|f| \leq C$ ,  $\varepsilon > 0$ .

Let  $\tilde{\xi}_n = \xi I(\xi \leq n)$ .

Obviously,  $\tilde{\xi}_n \xrightarrow{\text{a.s.}} \xi$ . Therefore,  $\tilde{\xi}_n \xrightarrow{P} \xi$ .

## The proof of 2)

$$P \Rightarrow d$$

$f : \mathbb{R} \rightarrow \mathbb{R}$  — continuous,  $|f| \leq C$ ,  $\varepsilon > 0$ .

Let  $\tilde{\xi}_n = \xi I(\xi \leq n)$ .

Obviously,  $\tilde{\xi}_n \xrightarrow{\text{a.s.}} \xi$ . Therefore,  $\tilde{\xi}_n \xrightarrow{P} \xi$ .

There exists  $n_1 \in \mathbb{N}$  such that, for all  $n \geq n_1$ ,

$$P(|\tilde{\xi}_n - \xi| > 1) < \varepsilon/(6C).$$

## The proof of 2)

$$P \Rightarrow d$$

$f : \mathbb{R} \rightarrow \mathbb{R}$  — continuous,  $|f| \leq C$ ,  $\varepsilon > 0$ .

Let  $\tilde{\xi}_n = \xi I(\xi \leq n)$ .

Obviously,  $\tilde{\xi}_n \xrightarrow{\text{a.s.}} \xi$ . Therefore,  $\tilde{\xi}_n \xrightarrow{P} \xi$ .

There exists  $n_1 \in \mathbb{N}$  such that, for all  $n \geq n_1$ ,

$$P(|\tilde{\xi}_n - \xi| > 1) < \varepsilon/(6C).$$

Therefore,  $\forall n \geq n_1, P(\xi > n) < \varepsilon/(6C)$ .

## The proof of 2)

$$P \Rightarrow d$$

## The proof of 2)

$$P \Rightarrow d$$

$f$  is uniformly continuous on any  $[a, b]$ .



## The proof of 2)

$$P \Rightarrow d$$

$f$  is uniformly continuous on any  $[a, b]$ .

Therefore, there exists a  $\delta > 0$  such that,  
if  $x, y \in [-n_1 - \delta, n_1 + \delta]$ ,  $|x - y| < \delta$ ,  
then  $|f(x) - f(y)| < \varepsilon/3$ .

## The proof of 2)

$$P \Rightarrow d$$

$f$  is uniformly continuous on any  $[a, b]$ .  
Therefore, there exists a  $\delta > 0$  such that,  
if  $x, y \in [-n_1 - \delta, n_1 + \delta]$ ,  $|x - y| < \delta$ ,  
then  $|f(x) - f(y)| < \varepsilon/3$ .

Let  $n_2 \in \mathbb{N}$  be such that, for all  $n \geq n_2$ ,

$$P(|\xi_n - \xi| > \delta) < \varepsilon/(6C).$$

## The proof of 2)

$$P \Rightarrow d$$

## The proof of 2)

$$P \Rightarrow d$$

For  $n \geq \max\{n_1, n_2\}$ ,

$$|Ef(\xi_n) - Ef(\xi)| = |E(f(\xi_n) - f(\xi))| \leq$$

## The proof of 2)

$$P \Rightarrow d$$

For  $n \geq \max\{n_1, n_2\}$ ,

$$|Ef(\xi_n) - Ef(\xi)| = |E(f(\xi_n) - f(\xi))| \leq$$

$$E|f(\xi_n) - f(\xi)| =$$

## The proof of 2)

$$P \Rightarrow d$$

For  $n \geq \max\{n_1, n_2\}$ ,

$$\begin{aligned} |Ef(\xi_n) - Ef(\xi)| &= |E(f(\xi_n) - f(\xi))| \leq \\ E|f(\xi_n) - f(\xi)| &= E\left(|f(\xi_n) - f(\xi)|I(\xi > n_1) + \right. \\ &\quad \left. |f(\xi_n) - f(\xi)|I(\xi \leq n_1, |\xi - \xi_n| \geq \delta) + \right. \\ &\quad \left. |f(\xi_n) - f(\xi)|I(\xi \leq n_1, |\xi - \xi_n| < \delta)\right) < \end{aligned}$$

## The proof of 2)

$$P \Rightarrow d$$

For  $n \geq \max\{n_1, n_2\}$ ,

$$\begin{aligned} |Ef(\xi_n) - Ef(\xi)| &= |E(f(\xi_n) - f(\xi))| \leq \\ E|f(\xi_n) - f(\xi)| &= E\left(|f(\xi_n) - f(\xi)|I(\xi > n_1) + \right. \\ &\quad \left. |f(\xi_n) - f(\xi)|I(\xi \leq n_1, |\xi - \xi_n| \geq \delta) + \right. \\ &\quad \left. |f(\xi_n) - f(\xi)|I(\xi \leq n_1, |\xi - \xi_n| < \delta)\right) < \\ 2CP(\xi > n_1) + 2CP(|\xi - \xi_n| \geq \delta) + \varepsilon/3 &< \varepsilon. \end{aligned}$$

## The proof of 3)

$$L_p \Rightarrow P$$



## The proof of 3)

$$L_p \Rightarrow P$$

Let  $\varepsilon > 0$ .

## The proof of 3)

$$L_p \Rightarrow P$$

Let  $\varepsilon > 0$ .

$$P(|\xi_n - \xi| \geq \varepsilon) = P(|\xi_n - \xi|^p \geq \varepsilon^p) \leq$$

## The proof of 3)

$$L_p \Rightarrow P$$

Let  $\varepsilon > 0$ .

$$P(|\xi_n - \xi| \geq \varepsilon) = P(|\xi_n - \xi|^p \geq \varepsilon^p) \leq$$

$$\frac{E|\xi_n - \xi|^p}{\varepsilon^p} \rightarrow 0.$$

## Counterexamples

$d$  does not imply  $P$

## Counterexamples

$d$  does not imply P

Let  $\xi_1 = \xi_2 = \dots = -\xi$ ,

## Counterexamples

$d$  does not imply  $P$

Let  $\xi_1 = \xi_2 = \dots = -\xi$ ,  $\xi \sim U(\{-1, 1\})$ .

## Counterexamples

$d$  does not imply  $P$

Let  $\xi_1 = \xi_2 = \dots = -\xi$ ,  $\xi \sim U(\{-1, 1\})$ .

$\xi_n \xrightarrow{d} \xi$ , since, for any  $f$ ,  $Ef(\xi_n) = Ef(\xi)$ .

## Counterexamples

$d$  does not imply  $P$

Let  $\xi_1 = \xi_2 = \dots = -\xi$ ,  $\xi \sim U(\{-1, 1\})$ .

$\xi_n \xrightarrow{d} \xi$ , since, for any  $f$ ,  $Ef(\xi_n) = Ef(\xi)$ .

However,  $P(|\xi - \xi_n| \geq 2) = 1$  for  $n \in \mathbb{N}$ .



## Counterexamples

$d$  does not imply  $P$

Let  $\xi_1 = \xi_2 = \dots = -\xi$ ,  $\xi \sim U(\{-1, 1\})$ .  
 $\xi_n \xrightarrow{d} \xi$ , since, for any  $f$ ,  $Ef(\xi_n) = Ef(\xi)$ .

However,  $P(|\xi - \xi_n| \geq 2) = 1$  for  $n \in \mathbb{N}$ .  
 $\xi_n$  does not converge in probability to  $\xi$ .

## Counterexamples

P does not imply neither a.s. nor  $L_p$

## Counterexamples

$P$  does not imply neither a.s. nor  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

## Counterexamples

P does not imply neither a.s. nor  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\xi_1 = 2^{1/p} I_{[0,1)},$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\xi_1 = 2^{1/p} I_{[0,1)},$$

$$\xi_2 = 2^{2/p} I_{[0,1/2)}, \quad \xi_3 = 2^{3/p} I_{[1/2,1)},$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\xi_1 = 2^{1/p} I_{[0,1)},$$

$$\xi_2 = 2^{2/p} I_{[0,1/2)}, \quad \xi_3 = 2^{3/p} I_{[1/2,1)},$$

$$\xi_4 = 2^{4/p} I_{[0,1/4)}, \quad \xi_5 = 2^{5/p} I_{[1/4,1/2)},$$

$$\xi_6 = 2^{6/p} I_{[1/2,3/4)}, \quad \xi_7 = 2^{7/p} I_{[3/4,1)},$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\begin{aligned}\xi_1 &= 2^{1/p} I_{[0,1)}, \\ \xi_2 &= 2^{2/p} I_{[0,1/2)}, \quad \xi_3 = 2^{3/p} I_{[1/2,1)}, \\ \xi_4 &= 2^{4/p} I_{[0,1/4)}, \quad \xi_5 = 2^{5/p} I_{[1/4,1/2)}, \\ \xi_6 &= 2^{6/p} I_{[1/2,3/4)}, \quad \xi_7 = 2^{7/p} I_{[3/4,1)}, \quad \dots\end{aligned}$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$




# Counterexamples

P does not imply neither a.s. nor  $L_p$




# Counterexamples

P does not imply neither a.s. nor  $L_p$




# Counterexamples

P does not imply neither a.s. nor  $L_p$




# Counterexamples

P does not imply neither a.s. nor  $L_p$



# Counterexamples

P does not imply neither a.s. nor  $L_p$



## Counterexamples

P does not imply neither a.s. nor  $L_p$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

Let  $\varepsilon > 0$ ,  $\delta > 0$ .

## Counterexamples

P does not imply neither a.s. nor  $L_p$

Let  $\varepsilon > 0$ ,  $\delta > 0$ .

Find  $k$  such that  $\delta > \frac{1}{2^k}$ .



## Counterexamples

P does not imply neither a.s. nor  $L_p$

Let  $\varepsilon > 0$ ,  $\delta > 0$ .

Find  $k$  such that  $\delta > \frac{1}{2^k}$ .

For every  $n \geq 2^k$ ,  $P(|\xi_n| > \varepsilon) \leq \frac{1}{2^k} < \delta$ .

## Counterexamples

P does not imply neither a.s. nor  $L_p$

Let  $\varepsilon > 0$ ,  $\delta > 0$ .

Find  $k$  such that  $\delta > \frac{1}{2^k}$ .

For every  $n \geq 2^k$ ,  $P(|\xi_n| > \varepsilon) \leq \frac{1}{2^k} < \delta$ .

$$\xi_n \xrightarrow{P} 0$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

## Counterexamples

$P$  does not imply neither a.s. nor  $L_p$

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

## Counterexamples

P does not imply neither a.s. nor  $L_p$

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

$$E|\xi_n|^p \geq 2^{2^k-1} \text{ for every } n \geq 2^k.$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

$$E|\xi_n|^p \geq 2^{2^k-1} \text{ for every } n \geq 2^k.$$

$$E|\xi_n|^p \rightarrow \infty.$$

## Counterexamples

P does not imply neither a.s. nor  $L_p$

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

$$E|\xi_n|^p \geq 2^{2^k-1} \text{ for every } n \geq 2^k.$$

$$E|\xi_n|^p \rightarrow \infty.$$

$\xi_n$  does not converge to 0 in  $L_p$



## Counterexamples

$L_p$  does not imply a.s.

## Counterexamples

$L_p$  does not imply a.s.

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

## Counterexamples

$L_p$  does not imply a.s.

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$\xi_1 = I_{[0,1)}$ ,

## Counterexamples

$L_p$  does not imply a.s.

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\xi_1 = I_{[0,1)},$$

$$\xi_2 = I_{[0,1/2)}, \quad \xi_3 = I_{[1/2,1)},$$

## Counterexamples

$L_p$  does not imply a.s.

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\xi_1 = I_{[0,1)},$$

$$\xi_2 = I_{[0,1/2)}, \xi_3 = I_{[1/2,1)},$$

$$\xi_4 = I_{[0,1/4)}, \xi_5 = I_{[1/4,1/2)},$$

$$\xi_6 = I_{[1/2,3/4)}, \xi_7 = I_{[3/4,1)},$$

## Counterexamples

$L_p$  does not imply a.s.

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform

$$\xi_1 = I_{[0,1)},$$

$$\xi_2 = I_{[0,1/2)}, \xi_3 = I_{[1/2,1)},$$

$$\xi_4 = I_{[0,1/4)}, \xi_5 = I_{[1/4,1/2)},$$

$$\xi_6 = I_{[1/2,3/4)}, \xi_7 = I_{[3/4,1)}, \dots$$

## Counterexamples

$L_p$  does not imply a.s.

## Counterexamples

$L_p$  does not imply a.s.

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .



## Counterexamples

$L_p$  does not imply a.s.

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

## Counterexamples

$L_p$  does not imply a.s.

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

$$E|\xi_n|^p \leq 2^{-k} \text{ for every } n \geq 2^k.$$

## Counterexamples

$L_p$  does not imply a.s.

For  $\omega \in [0, 1)$ , there are infinitely many  $n$  such that  $\xi_n(\omega) > 1$ .

$\xi_n$  does not converge to 0 a.s.

$$E|\xi_n|^p \leq 2^{-k} \text{ for every } n \geq 2^k.$$

$$\xi_n \xrightarrow{L_p} 0$$

## Counterexamples

a.s. does not imply  $L_p$

## Counterexamples

a.s. does not imply  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform.

## Counterexamples

a.s. does not imply  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform.

$$\xi_n = 2^{(1/p)n} I_{[0, 1/n]}, \quad n \in \mathbb{N}.$$

## Counterexamples

a.s. does not imply  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform.

$$\xi_n = 2^{(1/p)n} I_{[0, 1/n]}, \quad n \in \mathbb{N}.$$

For  $\omega \in (0, 1]$ ,  $\xi_n(\omega) = 0$  for  $n$  large enough.

## Counterexamples

a.s. does not imply  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform.

$$\xi_n = 2^{(1/p)n} I_{[0, 1/n]}, \quad n \in \mathbb{N}.$$

For  $\omega \in (0, 1]$ ,  $\xi_n(\omega) = 0$  for  $n$  large enough.

$$\xi_n \xrightarrow{\text{a.s.}} 0$$



## Counterexamples

a.s. does not imply  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform.

$$\xi_n = 2^{(1/p)n} I_{[0, 1/n]}, \quad n \in \mathbb{N}.$$

For  $\omega \in (0, 1]$ ,  $\xi_n(\omega) = 0$  for  $n$  large enough.

$$\xi_n \xrightarrow{\text{a.s.}} 0$$

$$E|\xi_n|^p = \frac{2^n}{n} \text{ for every } n \in \mathbb{N}.$$

## Counterexamples

a.s. does not imply  $L_p$

$\Omega = [0, 1]$ ,  $\mathcal{F} = \mathcal{B}[0, 1]$ ,  $P$  — uniform.

$$\xi_n = 2^{(1/p)n} I_{[0, 1/n]}, \quad n \in \mathbb{N}.$$

For  $\omega \in (0, 1]$ ,  $\xi_n(\omega) = 0$  for  $n$  large enough.

$$\xi_n \xrightarrow{\text{a.s.}} 0$$

$$E|\xi_n|^p = \frac{2^n}{n} \text{ for every } n \in \mathbb{N}.$$

$$E|\xi_n|^p \rightarrow \infty$$

## Simple random walk

$\xi_1, \xi_2, \dots$  — i.i.d. r.v.  $\sim U(\{-1, 1\})$ .

## Simple random walk

$\xi_1, \xi_2, \dots$  — i.i.d. r.v.  $\sim U(\{-1, 1\})$ .

$$S_0 = 0, S_n = \xi_1 + \dots + \xi_n$$

$\{S_n, n \in \mathbb{N}\}$  — simple random walk (on  $\mathbb{Z}$ )

## Simple random walk

$\xi_1, \xi_2, \dots$  — i.i.d. r.v.  $\sim U(\{-1, 1\})$ .


$$S_0 = 0, S_n = \xi_1 + \dots + \xi_n$$

$\{S_n, n \in \mathbb{N}\}$  — simple random walk (on  $\mathbb{Z}$ )

For  $\omega \in \Omega$ ,

$\{S_n(\omega), n \in \mathbb{N}\}$  — a trajectory of the random walk.

# Trajectory



## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

- ▶ if  $|x| > n$ ,



## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

- ▶ if  $|x| > n$ , then  $N(n, x) = 0$ ;

## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

- ▶ if  $|x| > n$ , then  $N(n, x) = 0$ ;
- ▶ if  $x + n$  is odd,

## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

- ▶ if  $|x| > n$ , then  $N(n, x) = 0$ ;
- ▶ if  $x + n$  is odd, then  $N(n, x) = 0$ ;

## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

- ▶ if  $|x| > n$ , then  $N(n, x) = 0$ ;
- ▶ if  $x + n$  is odd, then  $N(n, x) = 0$ ;
- ▶ otherwise,

## Counting trajectories

$N(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n(\omega) = x$ .

- ▶ if  $|x| > n$ , then  $N(n, x) = 0$ ;
- ▶ if  $x + n$  is odd, then  $N(n, x) = 0$ ;
- ▶ otherwise,  $N(n, x) = \binom{n}{\frac{n+x}{2}}$ .

## Counting trajectories

$\tilde{N}(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n = x, S_1, \dots, S_{n-1} \neq 0$ .

## Counting trajectories

$\tilde{N}(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n = x, S_1, \dots, S_{n-1} \neq 0$ .

### Theorem (Reflection principle)

*If  $x > 0$ , then*

$$\tilde{N}(n, x) = N(n-1, x-1) - N(n-1, -x-1).$$

## Counting trajectories

$\tilde{N}(n, x)$  — number of trajectories of  $\{S_k, k \leq n\}$  s.t.  $S_n = x, S_1, \dots, S_{n-1} \neq 0$ .

### Theorem (Reflection principle)

*If  $x > 0$ , then*

$$\tilde{N}(n, x) = N(n-1, x-1) - N(n-1, -x-1).$$

*If  $x < 0$ , then*

$$\tilde{N}(n, x) = N(n-1, x+1) - N(n-1, -x+1).$$



## The proof

Let  $x > 0$ .


## The proof

Let  $x > 0$ .

There exists a bijection between

$\{\text{trajectories s.t. } S_1 = 1, S_n = -x\},$

$\{\text{trajectories s.t. } S_1 = 1, S_n = x, \text{ at least one of } S_2, \dots, S_{n-1} \text{ equals } 0\}.$




## The proof

Let  $x > 0$ .

There exists a bijection between

$\{\text{trajectories s.t. } S_1 = 1, S_n = -x\},$

$\{\text{trajectories s.t. } S_1 = 1, S_n = x, \text{ at least one of } S_2, \dots, S_{n-1} \text{ equals } 0\}.$



## The proof

$$\tilde{N}(n, x) =$$

## The proof

$$\tilde{N}(n, x) =$$

the number of trajectories from  $(1, 1)$  to  $(n, x)$  that never meet  $0 =$

## The proof

$$\tilde{N}(n, x) =$$

the number of trajectories from  $(1, 1)$  to  $(n, x)$  that never meet  $0 =$

$N(n - 1, x - 1)$ —the number of trajectories from  $(1, 1)$  to  $(n, x)$  that meet  $0 =$

## The proof

$$\tilde{N}(n, x) =$$

the number of trajectories from  $(1, 1)$  to  $(n, x)$  that never meet  $0 =$

$N(n - 1, x - 1)$ —the number of trajectories from  $(1, 1)$  to  $(n, x)$  that meet  $0 =$

$$N(n - 1, x - 1) - N(n - 1, -x - 1).$$

## The proof

$$\tilde{N}(n, x) =$$

the number of trajectories from  $(1, 1)$  to  $(n, x)$  that never meet  $0 =$

$N(n - 1, x - 1)$ —the number of trajectories from  $(1, 1)$  to  $(n, x)$  that meet  $0 =$   
 $N(n - 1, x - 1) - N(n - 1, -x - 1).$

The case  $x < 0$  is symmetric.



## Counting probabilities

$$P(S_n = x) =$$

## Counting probabilities

$$P(S_n = x) = \frac{1}{2^n} N(n, x) =$$

## Counting probabilities

$$P(S_n = x) = \frac{1}{2^n} N(n, x) = \frac{1}{2^n} \binom{n}{\frac{n+x}{2}}$$

## Counting probabilities

$$P(S_n = x) = \frac{1}{2^n} N(n, x) = \frac{1}{2^n} \binom{n}{\frac{n+x}{2}}$$

$$P(S_1 > 0, \dots, S_{n-1} > 0, S_n = x) =$$

## Counting probabilities

$$P(S_n = x) = \frac{1}{2^n} N(n, x) = \frac{1}{2^n} \binom{n}{\frac{n+x}{2}}$$

$$P(S_1 > 0, \dots, S_{n-1} > 0, S_n = x) = \frac{\tilde{N}(n, x)}{2^n} =$$

## Counting probabilities

$$P(S_n = x) = \frac{1}{2^n} N(n, x) = \frac{1}{2^n} \binom{n}{\frac{n+x}{2}}$$

$$P(S_1 > 0, \dots, S_{n-1} > 0, S_n = x) = \frac{\tilde{N}(n, x)}{2^n} =$$

$$\frac{N(n-1, x-1) - N(n-1, -x-1)}{2^n} =$$

## Counting probabilities

$$P(S_n = x) = \frac{1}{2^n} N(n, x) = \frac{1}{2^n} \binom{n}{\frac{n+x}{2}}$$

$$P(S_1 > 0, \dots, S_{n-1} > 0, S_n = x) = \frac{\tilde{N}(n, x)}{2^n} =$$

$$\frac{N(n-1, x-1) - N(n-1, -x-1)}{2^n} =$$

$$\frac{1}{2^n} \left( \binom{n-1}{\frac{n+x-2}{2}} - \binom{n-1}{\frac{n-x-2}{2}} \right).$$

## Returning to 0

Find  $P(S_n \text{ never meets } 0)$ .



## Returning to 0

Find  $P(S_n \text{ never meets } 0)$ .

$$P(\forall n \geq 1 \ S_n \neq 0) =$$

## Returning to 0

Find  $P(S_n \text{ never meets } 0)$ .

$$P(\forall n \geq 1 \ S_n \neq 0) =$$

$$2P(\forall n \geq 1 \ S_n > 0) =$$

## Returning to 0

Find  $P(S_n \text{ never meets } 0)$ .

$$P(\forall n \geq 1 \ S_n \neq 0) =$$

$$2P(\forall n \geq 1 \ S_n > 0) =$$

$$\lim_{n \rightarrow \infty} \sum_{x=1}^n P(S_1 > 0, \dots, S_{n-1} > 0, S_n = x)$$

## Returning to 0

Find  $P(S_n \text{ never meets } 0)$ .

$$P(\forall n \geq 1 \ S_n \neq 0) =$$

$$2P(\forall n \geq 1 \ S_n > 0) =$$

$$\lim_{n \rightarrow \infty} \sum_{x=1}^n P(S_1 > 0, \dots, S_{n-1} > 0, S_n = x) = 0.$$

## Law of large numbers

Since  $ES_n = 0$ , we get

$$\frac{S_n}{\sqrt{nf(n)}} \xrightarrow{P} 0$$

for  $f(n) \uparrow \infty$ .

## Law of large numbers

Since  $ES_n = 0$ , we get

$$\frac{S_n}{\sqrt{nf(n)}} \xrightarrow{P} 0$$

for  $f(n) \uparrow \infty$ .

- What about a.s. convergence?

## Law of large numbers

Since  $ES_n = 0$ , we get

$$\frac{S_n}{\sqrt{nf(n)}} \xrightarrow{P} 0$$

for  $f(n) \uparrow \infty$ .

- ▶ What about a.s. convergence?
- ▶ Is there a limit of  $\frac{S_n}{\sqrt{n}}$ ?