

Probability theory

Lecture 3: Probability distribution

Maksim Zhukovskii

MIPT

Distribution in \mathbb{R}

$$\Omega = \mathbb{R},$$

Distribution in \mathbb{R}

$$\Omega = \mathbb{R},$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}),$$

Distribution in \mathbb{R}

$$\Omega = \mathbb{R},$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}),$$

P — probability measure on (Ω, \mathcal{F})

Distribution in \mathbb{R}

$$\Omega = \mathbb{R},$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}),$$

P — probability measure on (Ω, \mathcal{F}) —
distribution

Distribution function

$$F : \mathbb{R} \rightarrow [0, 1],$$

Distribution function

$$F : \mathbb{R} \rightarrow [0, 1], \quad F(x) = P((-\infty, x])$$

Distribution function

$$F : \mathbb{R} \rightarrow [0, 1], \quad F(x) = P((-\infty, x])$$

F — distribution function

Properties

1. F is nondecreasing,

Properties

1. F is nondecreasing,
2. $\lim_{x \rightarrow -\infty} F(x) = 0, \lim_{x \rightarrow \infty} F(x) = 1,$

Properties

1. F is nondecreasing,
2. $\lim_{x \rightarrow -\infty} F(x) = 0, \lim_{x \rightarrow \infty} F(x) = 1,$
3. F is right-continuous.

The proof

From distribution function to probability

Theorem

Let $F : \mathbb{R} \rightarrow [0, 1]$,

From distribution function to probability

Theorem

*Let $F : \mathbb{R} \rightarrow [0, 1]$,
 F have properties 1), 2), 3).*

From distribution function to probability

Theorem

*Let $F : \mathbb{R} \rightarrow [0, 1]$,
 F have properties 1), 2), 3).*

*Then, there exists the only distribution P
such that*

From distribution function to probability

Theorem

*Let $F : \mathbb{R} \rightarrow [0, 1]$,
 F have properties 1), 2), 3).*

*Then, there exists the only distribution P
such that*

- F is the distribution function of P ,*

From distribution function to probability

Theorem

Let $F : \mathbb{R} \rightarrow [0, 1]$,
 F have properties 1), 2), 3).

Then, there exists the only distribution P such that

- F is the distribution function of P ,*
- for all $-\infty \leq a < b < \infty$,*

$$P((a, b]) = F(b) - F(a).$$

Examples

$$F(x) = \begin{cases} 0, & x < 0; \\ x, & 0 \leq x \leq 1; \\ 1, & x > 1. \end{cases}$$

Examples

$$F(x) = \begin{cases} 0, & x < 0; \\ x, & 0 \leq x \leq 1; \\ 1, & x > 1. \end{cases}$$

The respective probability distribution is called **Lebesgue measure** μ on $[0, 1]$.

Examples

$$F(x) = \begin{cases} 0, & x < 0; \\ x, & 0 \leq x \leq 1; \\ 1, & x > 1. \end{cases}$$

The respective probability distribution is called **Lebesgue measure** μ on $[0, 1]$.

For $a < b \in [0, 1]$,

$$\mu((a, b)) = \mu([a, b)) = \mu((a, b]) = \mu([a, b]) = b - a.$$

Examples

$$F(x) = \begin{cases} 0, & x < c; \\ 1, & x \geq c. \end{cases}$$

Examples

$$F(x) = \begin{cases} 0, & x < c; \\ 1, & x \geq c. \end{cases}$$

$$\text{for } A \in \mathcal{B}(\mathbb{R}), \quad P(A) = \begin{cases} 0, & c \notin A, \\ 1, & c \in A, \end{cases}$$

Discrete distribution

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}$ — finite or countably infinite

Discrete distribution

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}$ — finite or countably infinite
- ▶ $P(\{x_k\}) = p_k > 0$ for all k , $P(X) = 1$

Discrete distribution

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}$ — finite or countably infinite
- ▶ $P(\{x_k\}) = p_k > 0$ for all k , $P(X) = 1$

P — discrete distribution

Discrete distribution

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}$ — finite or countably infinite
- ▶ $P(\{x_k\}) = p_k > 0$ for all k , $P(X) = 1$

P — discrete distribution

$$F(x) = \sum_{k: x_k < x} p_k$$

Examples

- ▶ Uniform distrib. on $\{1, \dots, N\}$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$

$$X = \{1, \dots, N\},$$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$

$$X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p
 $X = \{0, 1\},$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$

$$X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$$

- ▶ **Bernoulli** distrib. with parameter p

$$X = \{0, 1\}, \quad P(\{0\}) = 1 - p, \quad P(\{1\}) = p.$$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$

$$X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$$

- ▶ **Bernoulli** distrib. with parameter p

$$X = \{0, 1\}, \quad P(\{0\}) = 1 - p, \quad P(\{1\}) = p.$$

- ▶ **Binomial** distrib. with parameters (n, p) ,

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p
 $X = \{0, 1\}, \quad P(\{0\}) = 1 - p, P(\{1\}) = p.$
- ▶ **Binomial** distrib. with parameters (n, p) ,
 $X = \{0, 1, \dots, n\},$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p
 $X = \{0, 1\}, \quad P(\{0\}) = 1 - p, P(\{1\}) = p.$
- ▶ **Binomial** distrib. with parameters (n, p) ,
 $X = \{0, 1, \dots, n\}, \quad P(\{k\}) = \binom{n}{k} p^k (1 - p)^{n-k}.$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p
 $X = \{0, 1\}, \quad P(\{0\}) = 1 - p, P(\{1\}) = p.$
- ▶ **Binomial** distrib. with parameters (n, p) ,
 $X = \{0, 1, \dots, n\}, \quad P(\{k\}) = \binom{n}{k} p^k (1 - p)^{n-k}.$
- ▶ **Poisson** distrib. with parameter $\lambda > 0$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p
 $X = \{0, 1\}, \quad P(\{0\}) = 1 - p, P(\{1\}) = p.$
- ▶ **Binomial** distrib. with parameters (n, p) ,
 $X = \{0, 1, \dots, n\}, \quad P(\{k\}) = \binom{n}{k} p^k (1 - p)^{n-k}.$
- ▶ **Poisson** distrib. with parameter $\lambda > 0$
 $X = \mathbb{Z}_+,$

Examples

- ▶ **Uniform** distrib. on $\{1, \dots, N\}$
 $X = \{1, \dots, N\}, \quad p_k = \frac{1}{N}.$
- ▶ **Bernoulli** distrib. with parameter p
 $X = \{0, 1\}, \quad P(\{0\}) = 1 - p, P(\{1\}) = p.$
- ▶ **Binomial** distrib. with parameters (n, p) ,
 $X = \{0, 1, \dots, n\}, \quad P(\{k\}) = \binom{n}{k} p^k (1 - p)^{n-k}.$
- ▶ **Poisson** distrib. with parameter $\lambda > 0$
 $X = \mathbb{Z}_+, \quad P(\{k\}) = \frac{e^{-\lambda} \lambda^k}{k!}.$

Continuous distributions

F — distribution function of P

Continuous distributions

F — distribution function of P

Let there exist $p : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

Continuous distributions

F — distribution function of P

Let there exist $p : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}} p(t) dt = 1, \quad F(x) = \int_{-\infty}^x p(t) dt.$$

Continuous distributions

F — distribution function of P

Let there exist $p : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}} p(t) dt = 1, \quad F(x) = \int_{-\infty}^x p(t) dt.$$

Then

Continuous distributions

F — distribution function of P

Let there exist $p : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}} p(t) dt = 1, \quad F(x) = \int_{-\infty}^x p(t) dt.$$

Then

P — absolutely continuous distribution,

Continuous distributions

F — distribution function of P

Let there exist $p : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}} p(t) dt = 1, \quad F(x) = \int_{-\infty}^x p(t) dt.$$

Then

P — absolutely continuous distribution,
 p — density.

From density to distribution

Theorem

Let $p : \mathbb{R} \rightarrow \mathbb{R}_+$, $\int_{\mathbb{R}} p(t) dt = 1$.

From density to distribution

Theorem

Let $p : \mathbb{R} \rightarrow \mathbb{R}_+$, $\int_{\mathbb{R}} p(t) dt = 1$.

Then

$F : \mathbb{R} \rightarrow [0, 1]$,

$$F(x) = \int_{-\infty}^x p(t) dt,$$

is a distribution function.

Proof

Properties of density

- ▶ Let F be a differentiable distribution function.

Properties of density

- ▶ Let F be a differentiable distribution function. Then it has a density $p = F'$.

Properties of density

- ▶ Let F be a differentiable distribution function. Then it has a density $p = F'$.
- ▶ If p — density of P ,
 $B \in \mathcal{B}(\mathbb{R})$,

Properties of density

- ▶ Let F be a differentiable distribution function. Then it has a density $p = F'$.
- ▶ It p — density of P ,
 $B \in \mathcal{B}(\mathbb{R})$,

$$\text{then } P(B) = \int_B p d\mu.$$

Properties of density

► Let F be a differentiable distribution function. Then it has a density $p = F'$.

► If p — density of P ,
 $B \in \mathcal{B}(\mathbb{R})$,

$$\text{then } P(B) = \int_B p d\mu.$$

$$\text{If } \int_B p(x) dx \text{ exists,} \\ \text{then } P(B) = \int_B p(x) dx.$$

Examples

- Uniform distrib. on $[a, b]$:

$$p(t) = \frac{I_{[a,b]}(t)}{b-a}.$$

Examples

- **Uniform** distrib. on $[a, b]$:

$$p(t) = \frac{I_{[a,b]}(t)}{b-a}.$$

- **Normal** distrib. with parameters $(a, \sigma^2 > 0)$

$$p(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-a)^2}{2\sigma^2}}.$$

Examples

- **Uniform** distrib. on $[a, b]$:

$$p(t) = \frac{I_{[a,b]}(t)}{b-a}.$$

- **Normal** distrib. with parameters $(a, \sigma^2 > 0)$

$$p(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-a)^2}{2\sigma^2}}.$$

- **Gamma** distrib. with param. $(\alpha, \beta) \in \mathbb{R}_{>0}^2$

$$p(t) = \frac{\beta^\alpha t^{\alpha-1}}{\Gamma(\alpha)} e^{-\beta t} I_{(0,\infty)}(t).$$

Examples

- **Uniform** distrib. on $[a, b]$:

$$p(t) = \frac{I_{[a,b]}(t)}{b-a}.$$

- **Normal** distrib. with parameters $(a, \sigma^2 > 0)$

$$p(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-a)^2}{2\sigma^2}}.$$

- **Gamma** distrib. with param. $(\alpha, \beta) \in \mathbb{R}_{>0}^2$

$$p(t) = \frac{\beta^\alpha t^{\alpha-1}}{\Gamma(\alpha)} e^{-\beta t} I_{(0,\infty)}(t).$$

- **Exponential** distrib. with parameter $\lambda > 0$:

$$p(t) = \lambda e^{-\lambda t} I_{(0,\infty)}(t).$$

Examples

- **Uniform** distrib. on $[a, b]$:

$$p(t) = \frac{I_{[a,b]}(t)}{b-a}.$$

- **Normal** distrib. with parameters $(a, \sigma^2 > 0)$

$$p(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-a)^2}{2\sigma^2}}.$$

- **Gamma** distrib. with param. $(\alpha, \beta) \in \mathbb{R}_{>0}^2$

$$p(t) = \frac{\beta^\alpha t^{\alpha-1}}{\Gamma(\alpha)} e^{-\beta t} I_{(0,\infty)}(t).$$

- **Exponential** distrib. with parameter $\lambda > 0$:

$$p(t) = \lambda e^{-\lambda t} I_{(0,\infty)}(t).$$

- **Cauchy** distrib. with parameter $\theta > 0$:

$$p(t) = \frac{\theta}{\pi(t^2 + \theta^2)}.$$

n -dimensional distribution

$$\Omega = \mathbb{R}^n,$$

n -dimensional distribution

$$\Omega = \mathbb{R}^n,$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}^n) = \sigma(\{B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R})\})$$

n -dimensional distribution

$$\Omega = \mathbb{R}^n,$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}^n) = \sigma(\{B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R})\})$$

- ▶ P — probability measure on (Ω, \mathcal{F})

n -dimensional distribution

$$\Omega = \mathbb{R}^n,$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}^n) = \sigma(\{B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R})\})$$

- ▶ P — probability measure on (Ω, \mathcal{F}) —
distribution

n -dimensional distribution

$$\Omega = \mathbb{R}^n,$$

$$\mathcal{F} = \mathcal{B}(\mathbb{R}^n) = \sigma(\{B_1 \times \dots \times B_n, B_i \in \mathcal{B}(\mathbb{R})\})$$

- ▶ P — probability measure on (Ω, \mathcal{F}) —
distribution
- ▶ $F : \mathbb{R}^n \rightarrow [0, 1]$, $F(x_1, \dots, x_n) =$
 $P((-\infty, x_1] \times \dots \times (-\infty, x_n])$ —
distribution function

Properties

Notations

► $x \geq y$

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$
- ▶ $(-\infty, \mathbf{x}]$

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$
- ▶ $(-\infty, \mathbf{x}] := (-\infty, x_1] \times \dots \times (-\infty, x_n]$

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$
- ▶ $(-\infty, \mathbf{x}] := (-\infty, x_1] \times \dots \times (-\infty, x_n]$
- ▶ $\mathbf{x}^k \downarrow \mathbf{x}$:

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$
- ▶ $(-\infty, \mathbf{x}] := (-\infty, x_1] \times \dots \times (-\infty, x_n]$
- ▶ $\mathbf{x}^k \downarrow \mathbf{x}$:
 $\mathbf{x}^{k+1} \leq \mathbf{x}^k, k \in \mathbb{N},$
 $\lim_{k \rightarrow \infty} \mathbf{x}^k = \mathbf{x}.$

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$
- ▶ $(-\infty, \mathbf{x}] := (-\infty, x_1] \times \dots \times (-\infty, x_n]$
- ▶ $\mathbf{x}^k \downarrow \mathbf{x}$:
 $\mathbf{x}^{k+1} \leq \mathbf{x}^k, k \in \mathbb{N},$
 $\lim_{k \rightarrow \infty} \mathbf{x}^k = \mathbf{x}.$
- ▶ $\Delta_{a,b}^i F(x_1, \dots, x_n)$

Properties

Notations

- ▶ $\mathbf{x} \geq \mathbf{y}$: $x_i \geq y_i, i \in \{1, \dots, n\}$
- ▶ $(-\infty, \mathbf{x}] := (-\infty, x_1] \times \dots \times (-\infty, x_n]$
- ▶ $\mathbf{x}^k \downarrow \mathbf{x}$:
 $\mathbf{x}^{k+1} \leq \mathbf{x}^k, k \in \mathbb{N},$
 $\lim_{k \rightarrow \infty} \mathbf{x}^k = \mathbf{x}.$
- ▶ $\Delta_{a,b}^i F(x_1, \dots, x_n) :=$
 $F(x_1, \dots, x_{i-1}, b, x_{i+1}, \dots, x_n) -$
 $F(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n).$

Properties

1. For all $a_1 < b_1, \dots, a_n < b_n$,
$$\Delta_{a_1, b_1}^1 \cdots \Delta_{a_n, b_n}^n F(x_1, \dots, x_n) \geq 0.$$

Properties

1. For all $a_1 < b_1, \dots, a_n < b_n$,

$$\Delta_{a_1, b_1}^1 \cdots \Delta_{a_n, b_n}^n F(x_1, \dots, x_n) \geq 0.$$

2.

$$\lim_{x_1 \rightarrow \infty, \dots, x_n \rightarrow \infty} F(x_1, \dots, x_n) = 1,$$

$$\forall i \in \{1, \dots, n\} \quad \lim_{x_i \rightarrow -\infty} F(x_1, \dots, x_n) = 0.$$

Properties

1. For all $a_1 < b_1, \dots, a_n < b_n$,
 $\Delta_{a_1, b_1}^1 \cdots \Delta_{a_n, b_n}^n F(x_1, \dots, x_n) \geq 0$.

2.

$$\lim_{x_1 \rightarrow \infty, \dots, x_n \rightarrow \infty} F(x_1, \dots, x_n) = 1,$$

$$\forall i \in \{1, \dots, n\} \quad \lim_{x_i \rightarrow -\infty} F(x_1, \dots, x_n) = 0.$$

3. If $\mathbf{x}^k \downarrow \mathbf{x}$, then
 $F(\mathbf{x}^k) \rightarrow F(\mathbf{x})$.

From distribution function to probability

Theorem

Let $F : \mathbb{R}^n \rightarrow [0, 1]$,

From distribution function to probability

Theorem

*Let $F : \mathbb{R}^n \rightarrow [0, 1]$,
 F have properties 1), 2), 3).*

From distribution function to probability

Theorem

Let $F : \mathbb{R}^n \rightarrow [0, 1]$,
 F have properties 1), 2), 3).

*Then, there exists the only distribution P
such that*

From distribution function to probability

Theorem

Let $F : \mathbb{R}^n \rightarrow [0, 1]$,
 F have properties 1), 2), 3).

*Then, there exists the only distribution P
such that*

- *F is the distribution function of P ,*

From distribution function to probability

Theorem

Let $F : \mathbb{R}^n \rightarrow [0, 1]$,
 F have properties 1), 2), 3).

Then, there exists the only distribution P such that

- F is the distribution function of P ,*
- for all $a_1 < b_1, \dots, a_n < b_n$,*

$$\Delta_{a_1, b_1}^1 \cdots \Delta_{a_n, b_n}^n F(x_1, \dots, x_n) = \\ P((a_1, b_1] \times \dots \times (a_n, b_n]).$$

Examples

- ▶ If F^1, \dots, F^n — are 1-dimensional distribution functions,

Examples

- ▶ If F^1, \dots, F^n — are 1-dimensional distribution functions,
then $F^1(x_1) \dots F^n(x_n)$ — n -dimensional distribution function.

Examples

- ▶ If F^1, \dots, F^n — are 1-dimensional distribution functions,
then $F^1(x_1) \dots F^n(x_n)$ — n -dimensional distribution function.

▶

$$\left(\prod_{i \in \{1, \dots, n\}: x_i < 1} x_i \right) I(x_i > 0) + I(x_i > 1)$$

Examples

- ▶ If F^1, \dots, F^n — are 1-dimensional distribution functions,
then $F^1(x_1) \dots F^n(x_n)$ — n -dimensional distribution function.



$$\left(\prod_{i \in \{1, \dots, n\}: x_i < 1} x_i \right) I(x_i > 0) + I(x_i > 1)$$

is distribution function of **Lebesgue** measure on $[0, 1]^n$.

Discrete n -dimensional distributions

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}^n$ — finite or countably infinite

Discrete n -dimensional distributions

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}^n$ — finite or countably infinite
- ▶ $P(\{x_k\}) = p_k > 0$ for all k , $P(X) = 1$

Discrete n -dimensional distributions

- ▶ $X = \{x_1, x_2, \dots\} \subset \mathbb{R}^n$ — finite or countably infinite
- ▶ $P(\{x_k\}) = p_k > 0$ for all k , $P(X) = 1$

P — discrete distribution

Continuous n -dimensional distributions

F — distribution function of P

Continuous n -dimensional distributions

F — distribution function of P

Let there exist $p : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

Continuous n -dimensional distributions

F — distribution function of P

Let there exist $p : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}^n} p(t_1, \dots, t_n) dt_1 \dots dt_n = 1,$$

$$F(x) = \int_{-\infty}^{x_1} \dots \int_{-\infty}^{x_n} p(t_1, \dots, t_n) dt_1 \dots dt_n.$$

Continuous n -dimensional distributions

F — distribution function of P

Let there exist $p : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}^n} p(t_1, \dots, t_n) dt_1 \dots dt_n = 1,$$

$$F(x) = \int_{-\infty}^{x_1} \dots \int_{-\infty}^{x_n} p(t_1, \dots, t_n) dt_1 \dots dt_n.$$

Then

Continuous n -dimensional distributions

F — distribution function of P

Let there exist $p : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}^n} p(t_1, \dots, t_n) dt_1 \dots dt_n = 1,$$

$$F(x) = \int_{-\infty}^{x_1} \dots \int_{-\infty}^{x_n} p(t_1, \dots, t_n) dt_1 \dots dt_n.$$

Then

P — absolutely continuous distribution,

Continuous n -dimensional distributions

F — distribution function of P

Let there exist $p : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

$$\int_{\mathbb{R}^n} p(t_1, \dots, t_n) dt_1 \dots dt_n = 1,$$

$$F(x) = \int_{-\infty}^{x_1} \dots \int_{-\infty}^{x_n} p(t_1, \dots, t_n) dt_1 \dots dt_n.$$

Then

P — absolutely continuous distribution,

p — density.

Continuous n -dimensional distributions

- ▶ If F is differentiable distribution function, then $p = \frac{\partial^n}{\partial x_1 \dots \partial x_n} F$.

Continuous n -dimensional distributions

- ▶ If F is differentiable distribution function, then $p = \frac{\partial^n}{\partial x_1 \dots \partial x_n} F$.
- ▶ If p — density, then $P(B) = \int_B p d\mu$.

Continuous n -dimensional distributions

► If F is differentiable distribution function, then $p = \frac{\partial^n}{\partial x_1 \dots \partial x_n} F$.

► If p — density, then $P(B) = \int_B p d\mu$.

$P(B) = \int_B p(x_1, \dots, x_n) dx_1 \dots dx_n$, if the integral exists.