

NNT : ***

n°LAL : ***

Thèse de doctorat

Search of the $0\nu\beta\beta$ decay with the SuperNEMO demonstrator

Thèse de doctorat de l'Université Paris-Saclay
préparée à l'Université Paris Saclay au sein du Laboratoire Irène-Joliot Curie
(anciennement Laboratoire de l'Accélérateur Linéaire)

École doctorale n°576 Particles, Hadrons, Energy, Nuclei, Instrumentation,
Imaging, Cosmos et Simulation (PHENIICS)
Spécialité de doctorat : Physique des particules

Thèse présentée et soutenue à Orsay, le ***, par

CLOÉ GIRARD-CARILLO

Composition du Jury :

Président

Rapporteur

Rapporteur

Christine Marquet
CENBG - Bordeaux-Gradignan

Examineur

Examineur

Examineur

Laurent Simard
LAL - Orsay

Directeur de thèse

Mathieu Bongrand
LAL - Orsay

Co-directeur de thèse

Contents

Contents	3
Introduction	7
1 Phenomenology of particle physics	9
1.1 The Standard Model of particle physics	9
1.1.1 Bosons	9
1.1.2 Fermions	9
1.1.3 $2\nu\beta\beta$ decay	9
1.1.4 Where the Standard Model ends	9
1.2 Going beyond the Standard Model with neutrinos	9
1.2.1 Neutrino flavors and oscillations	9
1.2.2 Neutrino masses and nature	9
1.2.3 Other searches beyond the Standard Model with neutrinos	9
2 $0\nu\beta\beta$ experiment status	11
2.1 Experimental design criteria	11
2.1.1 Aspects of the nuclear matrix elements	12
2.1.2 Quenching	12
2.2 $0\nu\beta\beta$ direct search experiments	12
2.2.1 Semiconductors	12
2.2.2 Bolometers	14
2.2.3 Time projection chambers	14
2.2.4 Scintillators	16
2.2.5 Tracking calorimeters	16
3 The SuperNemo demonstrator	18
3.1 The SuperNemo demonstrator	18
3.1.1 Comparison with Nemo3 experiment	18
3.1.2 Experimental design	18
3.1.3 Sources	18
3.1.4 Tracker	18
3.1.5 Calorimeter	18
3.1.5.1 Scintillator	18
3.1.5.2 Photomultiplier	18
3.1.6 Calibration systems	18
3.1.7 Control Monitoring system	18

3.1.8	Electronics	18
3.2	The background of SuperNEMO	18
3.2.1	Internal background	18
3.2.2	External background	18
3.2.3	Background specifications	18
3.2.4	Measured demonstrator background levels	18
3.3	The SuperNemo software	18
3.3.1	Simulation	18
3.3.2	Reconstruction	19
4	Analysis tools	21
4.0.1	Internal probability	21
4.1	Simulations	22
4.1.1	Modifications of simulation software	22
4.1.2	Internal background simulations	22
4.1.3	$0\nu\beta\beta$ simulations	22
5	Time difference	23
5.1	Principle and goal	23
5.1.1	Internal conversion	23
5.2	Analysis	24
5.2.1	Topological cuts	24
5.2.2	Exponentially modified Gaussian	24
5.2.3	Results	24
5.3	Conclusion	24
6	Sensitivity of the SuperNEMO demonstrator to the $0\nu\beta\beta$	27
6.1	Signal and backgrounds considered	27
6.2	Optimisation of event selection	29
6.3	Expected number of background events	29
6.4	Demonstrator sensitivity	29
6.4.1	avec B	30
6.4.2	sans B	30
6.4.3	Champ mappé	30
6.5	HyperNEMO	30
6.6	Other isotopes	30
6.7	Conclusion	30
7	Detector commissioning	31
7.1	Reflectometry analysis	31
7.1.1	Goal of the reflectometry analysis	31
7.1.2	Pulse timing: controlling cable lengths	32
7.1.3	Signal attenuation	37
7.1.4	Pulse shape analysis	39
7.1.5	Comparison with ^{60}Co	39
7.1.6	Conclusion	39
7.2	Calibrating the electronic boards	39

7.2.1	Principle	39
7.2.2	Measuring the time offset of front end boards	39
7.2.3	Results	39
7.3	Energy calibration of optical modules	39
7.4	Baseline studies	39
7.5	Light Injection System	39
8	Characterisation of the calorimeter time resolution	41
8.1	Interaction of particles in the SuperNEMO scintillators	42
8.1.1	Interaction of electrons	42
8.1.2	Interaction of photons	42
8.2	Measurement of the time resolution with a ^{60}Co source	43
8.2.1	Description of Cobalt 60 nucleus	44
8.2.2	Time response of optical modules	44
8.2.3	Final experimental design	47
8.2.4	Signal events selection	49
8.2.5	Background estimation	51
8.2.6	Detector efficiency	55
8.2.7	Determination of the individual timing resolution of each optical module	56
8.2.8	Conclusion	60
8.3	The Light Injection System	60
8.3.1	Light injection system commissioning	61
8.3.2	Time resolution of optical modules	61
	Conclusion	63
	Bibliography	65

Chapter 6

Sensitivity of the SuperNEMO demonstrator to the $0\nu\beta\beta$

In this chapter, we present the SuperNEMO sensitivity to the $0\nu\beta\beta$ decay half-life, and the corresponding effective neutrino masses, for several isotopes. The SuperNEMO final detector is expected to exclude $0\nu\beta\beta$ half-lives up to 1.2×10^{26} y (90% CL) if $0\nu\beta\beta$ decays through the mass mechanism, with a detector exposure of 500 kg.y [7]. The sensitivity is given as a limit, in case we do not observe the expected signal. In 2010 began the demonstrator installation at the Laboratoire Souterrain de Modane. With an exposure of 17.5 y, the demonstrator could set a limit on the $0\nu\beta\beta$ process of 5.35×10^{24} y (90% CL) [8].

This study aims to explore the impact on the sensitivity of the presence of a magnetic field, and will participate in the final decision on the installation of the coil. In a context of investigating the demonstrator and final detector capabilities, different internal source contamination levels are explored. The topology of interest is the two electrons topology, and we use the $2e$ energy sum to discriminate the signal from the background events. Thanks to SuperNEMO tracking capabilities, topological informations are exploited to improve the SuperNEMO sensitivity.

6.1 Signal and backgrounds considered

A full simulation for the SuperNEMO demonstrator was performed, in order to determine the longest $0\nu\beta\beta$ half-life that can be probed with SuperNEMO using the distribution of the sum of electron energies, in the case where the $0\nu\beta\beta$ decay were not observed. In the Tab. 6.1 is summarised the expected number of signal and background events, both for the SuperNEMO demonstrator and final detector, and we present the amount of simulated Monte-Carlo events for each considered decay.

The $0\nu\beta\beta$ signal

In the following, the assumed underlying mechanism for the $0\nu\beta\beta$ decay is the mass mechanism (MM), as it is the most natural and widespread mechanism. The hypothetical $0\nu\beta\beta$ signal would be detected as an excess of events in the region

of interest, with respect to the predicted background contamination level. The 10^7 $0\nu\beta\beta$ Monte-Carlo events are generated using the DECAY0 software [9]. The simulations are normalised assuming a $T_{1/2}^{0\nu} = 6.0 \cdot 10^{24}$ y half-life [citation].

Internal backgrounds

As described in Sec. 3.2.1, source foils contaminations by isotopes such as ^{208}Tl or ^{214}Bi constitute the principal internal backgrounds with the $2\nu\beta\beta$ decay. These backgrounds are processed by the same detector simulation as the $0\nu\beta\beta$ signal, using DECAY0. Since internal backgrounds have very low efficiencies in the $2e$ topology, we simulated an important amount of Monte-Carlo events. The target background activities were defined so that each background has a similar contribution to that of the $2\nu\beta\beta$ in the region of interest [10].

The dominant two neutrino $2\nu\beta\beta$ background and the background due to foil contamination were normalised assuming a detector exposure of 500 kg.y.

Tracking volume background

External background

All external backgrounds from outside the foil, apart from ^{222}Rn in the tracking volume, are expected to be negligible and were not simulated.

Table 6.1: Expected and simulated decays for different processes, both for the demonstrator (17.5 kg.y) and for the final detector (500 kg.y).

	Expected decays		Simulated decays
	Demonstrator	Final detector	
$0\nu\beta\beta$ ($T_{1/2}^{0\nu} = 6.0 \cdot 10^{24}$ y)	$1.5 \cdot 10^1$	$2.7 \cdot 10^7$	$1.0 \cdot 10^7$
$2\nu\beta\beta$	$9.5 \cdot 10^5$	$4.2 \cdot 10^2$	$1.0 \cdot 10^7$
^{208}Tl	$5.5 \cdot 10^3$	$1.6 \cdot 10^5$	$1.0 \cdot 10^7$
^{214}Bi	$1.1 \cdot 10^3$	$3.1 \cdot 10^4$	$1.0 \cdot 10^7$
^{222}Rn	$1.8 \cdot 10^5$	$7.2 \cdot 10^6$	$1.0 \cdot 10^7$

Justifier bdf externe avec article nemo3 (plus diff roi et meilleure eff) Activités bkg considérées à justifier, bkg interne: balek externe à justifier

Demies vies 2ν à justifier

Se, Nd avec et sans champs

présentation du PID de Steven (peut être à bouger dans Tl selon développement du plan ou généralités)

Influence des quantités de contaminations sur la sensibilité




Figure 6.1

6.2 Optimisation of event selection

plot S/\sqrt{B} en fonction $E_l E_{\text{min}}$

Quel est le signal qu'on cherche
 présentation des cuts
 efficacité des cuts/ signal + bkg
 cuts premier et second ordre

6.3 Expected number of background events

plot energy tot plus dans région intérêt

6.4 Demonstrator sensitivity

Résultats $B = 0$, avec activités nominales, puis avec activités caca
 Efficiency spretra
 Energy spectra

6.4.1 avec B

Parler du champ non uniforme/attenuation ROI optimization: avec variation
coupure énergie

6.4.2 sans B

avec variation coupure énergie

6.4.3 Champ mappé

6.5 HyperNEMO

results for 500kg.y exposure

6.6 Other isotopes

bam bam le Nd

distribution $t_{1/2}$ avec différents échantillons de simus (17.5 kg.y)

6.7 Conclusion

Faut arrêter SN

Etude plus générale avec bkg externe+lab (reprendre chiffres NEMO3) + neutrons
(cf NEMO3)

Bibliography

- [1] M. et al. Agostini. Probing majorana neutrinos with double- β decay. *Science* 365, 1445, 2019.
- [2] S.I. et al Alvis. Search for neutrinoless double-beta decay in ^{76}Ge with 26 kg-yr of exposure from the majorana demonstrator. *Phys. Rev. C*, 100, 2019.
- [3] O. et al. Azzolini. First result on the neutrinoless double- β decay of ^{82}Se with cupid-0. *Phys. Rev. Lett.*, 120:232502, Jun 2018.
- [4] C. et al. Alduino. First results from cuore: A search for lepton number violation via $0\nu\beta\beta$ decay of ^{130}Te . *Phys. Rev. Lett.*, 120:132501, Mar 2018.
- [5] J. B. et al. Albert. Search for neutrinoless double-beta decay with the upgraded exo-200 detector. *Phys. Rev. Lett.*, 120:072701, Feb 2018.
- [6] A. et al. Gando. Search for majorana neutrinos near the inverted mass hierarchy region with kamland-zen. *Phys. Rev. Lett.*, 117:082503, Aug 2016.
- [7] R. et al. Arnold. Probing new physics models of neutrinoless double beta decay with supernemo. *Eur. Phys. J. C*, 2010.
- [8] S. Clavez. *Development of reconstruction tools and sensitivity of the SuperNEMO demonstrator*. PhD thesis, Université Paris Sud, 2017.
- [9] Tretyak V.I. Ponkratenko O.A. and Zdesenko Yu.G. The event generator decay4 for simulation of doublebeta processes and decay of radioactive nuclei. *Phys. At. Nucl.*, 63:1282–1287, Jul 2000.
- [10] Gomez-Cadenas et al. Physics case of supernemo with ^{82}Se source. Internal presentation, 2008.
- [11] Nucleid database.