

NNT : ***

n°LAL : ***

Thèse de doctorat

Search of the $0\nu\beta\beta$ decay with the SuperNEMO demonstrator

Thèse de doctorat de l'Université Paris-Saclay
préparée à l'Université Paris Saclay au sein du Laboratoire Irène-Joliot Curie
(anciennement Laboratoire de l'Accélérateur Linéaire)

École doctorale n°576 Particles, Hadrons, Energy, Nuclei, Instrumentation,
Imaging, Cosmos et Simulation (PHENIICS)
Spécialité de doctorat : Physique des particules

Thèse présentée et soutenue à Orsay, le ***, par

CLOÉ GIRARD-CARILLO

Composition du Jury :

Président

Rapporteur

Rapporteur

Christine Marquet
CENBG - Bordeaux-Gradignan

Examineur

Examineur

Examineur

Laurent Simard
LAL - Orsay

Directeur de thèse

Mathieu Bongrand
LAL - Orsay

Co-directeur de thèse

Contents

Contents	3
Introduction	7
1 Phenomenology of particle physics	9
1.1 The Standard Model of particle physics	9
1.1.1 Bosons	9
1.1.2 Fermions	9
1.1.3 $2\nu\beta\beta$ decay	9
1.1.4 Where the Standard Model ends	9
1.2 Going beyond the Standard Model with neutrinos	9
1.2.1 Neutrino flavors and oscillations	9
1.2.2 Neutrino masses and nature	9
1.2.3 Other searches beyond the Standard Model with neutrinos	9
1.3 $0\nu\beta\beta$ experiment status	9
1.3.1 Experimental design criteria	9
1.3.2 $0\nu\beta\beta$ direct search experiments	10
1.3.3 Bolometers	12
1.3.4 Time projection chambers	12
1.3.5 Scintillators	14
1.3.6 Tracking calorimeters	14
2 The SuperNemo demonstrator	15
2.1 The SuperNemo demonstrator	15
2.1.1 Comparison with Nemo3 experiment	15
2.1.2 Experimental design	15
2.1.3 Sources	15
2.1.4 Tracker	15
2.1.5 Calorimeter	15
2.1.6 Calibration systems	15
2.1.7 Control Monitoring system	15
2.1.8 Electronics	15

2.2	The background of SuperNEMO	15
2.2.1	Internal background	15
2.2.2	External background	16
2.2.3	Background specifications	17
2.2.4	Measured demonstrator background levels	17
2.3	Magnetic field	17
2.4	The SuperNemo software	17
2.4.1	Simulation	17
2.4.2	Reconstruction	17
2.4.3	Modifications of simulation software	17
2.5	Analysis tools	17
2.5.1	Internal probability	17
2.5.2	External probability	19
3	Sensitivity of the SuperNEMO demonstrator to the $0\nu\beta\beta$	21
3.1	The $0\nu\beta\beta$ signal and background model	21
3.1.1	The $0\nu\beta\beta$ signal	23
3.1.2	Inside detector backgrounds	23
3.1.3	External backgrounds	24
3.2	Event selection	24
3.3	Demonstrator sensitivity to the $0\nu\beta\beta$ decay of ^{82}Se	27
3.3.1	Sensitivity to the $0\nu\beta\beta$ half-life	27
3.3.2	Limit on the effective neutrino mass	29
3.4	Impact of sources contamination levels on the sensitivity	30
3.4.1	Influence of the contamination levels	30
3.4.2	Optimisation of the event selection	33
3.5	Impact of the magnetic field on the sensitivity	39
3.5.1	Simulations of the magnetic field inside the demonstrator and reconstructed track fit	39
3.5.2	Impact of the magnetic field on signal and background selections	40
3.5.3	Influence of the magnetic field on optical modules and reconstruction efficiency	42
3.5.4	Simulations with a non-uniform magnetic field	43
3.6	Searching for the Neodymium-150 $0\nu\beta\beta$ decay	43
3.6.1	Searching for the $0\nu\beta\beta$ of other isotopes	44
3.6.2	Sensitivity to the $0\nu\beta\beta$ of ^{150}Nd	44
3.7	The final detector sensitivity	46
3.8	Conclusion	46
4	Improvement of the rejection of the internal Thallium-208 background	49
4.1	Motivations	49
4.2	The internal ^{208}Tl background	50
4.2.1	The internal conversion process	51
4.2.2	Selection of ^{208}Tl disintegrations in the 2e channel	51
4.3	Rejection of ^{208}Tl with a time-of-flight criterion	53

4.3.1	The internal probability	53
4.3.2	The exponential probability for ^{208}Tl events	55
4.4	Event selection	57
4.4.1	Selection on particle detection times	57
4.5	Impact of ^{208}Tl rejection on the experiment's sensitivity	58
4.5.1	Influence of the calorimeter time resolution	58
4.6	Conclusions	61
5	Characterisation of the calorimeter time resolution	63
5.1	Interaction of particles in the SuperNEMO scintillators	64
5.1.1	Interaction of electrons	64
5.1.2	Interaction of photons	64
5.2	Measurement of the time resolution with a ^{60}Co source	65
5.2.1	Description of Cobalt 60 nucleus	66
5.2.2	Time response of optical modules	66
5.2.3	Final experimental design	69
5.2.4	Signal events selection	71
5.2.5	Background estimation	73
5.2.6	Detector efficiency	77
5.2.7	Determination of the individual timing resolution of each optical module	77
5.2.8	Conclusion	82
5.3	The Light Injection System	82
5.3.1	Light injection system commissioning	83
5.3.2	Time resolution of optical modules	83
6	Detector commissioning	85
6.1	Reflectometry analysis	85
6.1.1	Goal of the reflectometry analysis	85
6.1.2	Pulse timing: controlling cable lengths	86
6.1.3	Signal attenuation	91
6.1.4	Pulse shape analysis	93
6.1.5	Comparison with ^{60}Co	93
6.1.6	Conclusion	93
6.2	Calibrating the electronic boards	93
6.2.1	Principle	93
6.2.2	Measuring the time offset of front end boards	93
6.2.3	Results	93
6.3	Energy calibration of optical modules	93
6.4	Baseline studies	93
6.5	Light Injection System	93
	Conclusion	95
	Bibliography	97

Sensitivity of the SuperNEMO demonstrator to the $0\nu\beta\beta$

In this chapter, we present a study aiming to evaluate the SuperNEMO's sensitivity to the $0\nu\beta\beta$ decay, and the corresponding effective neutrino mass. Studies of this kind have already been conducted, and the final detector, based on the NEMO-3 technology, is expected to exclude half-lives up to 1.2×10^{26} y (90% CL), with an exposure of 500 kg.y for the ^{82}Se ¹ [8]. In order to assess the feasibility of such a large-scale detector, the SuperNEMO demonstrator started to be installed in early 2015, at the Laboratoire Souterrain de Modane. With an exposure of 17.5 kg.y, this demonstrator could reach a sensitivity on the $0\nu\beta\beta$ process of ^{82}Se 5.3×10^{24} y (90% CL) [9].

A copper coil was designed to deliver a uniform 25 Gauss magnetic field inside the wire chamber, to bend the charged particles trajectories, hence making it possible to discriminate between electrons and positrons. However, studies lead by the collaboration determined that this field could be impacted by the detector material (especially by the calorimeter magnetic shields), producing notable variations in intensity and a loss of energy resolution [9][10]. We aim to explore the impact, on both the demonstrator and final detector sensitivity, of the presence of this magnetic field. The findings of this study will participate in the ultimate decision on the installation of the coil. In a context of investigating the demonstrator and final detector's capabilities, different internal source contamination levels are considered. The topology of interest is the two electrons topologies, and we use the total energy sum to discriminate the signal from the background events. Thanks to SuperNEMO tracking capabilities, topological informations are exploited to improve the sensitivity. To go further, we also explore the possibility of studying the $0\nu\beta\beta$ decay of other $\beta\beta$ isotopes.

3.1 The $0\nu\beta\beta$ signal and background model

A full simulation for the demonstrator was performed, to determine the upper limit on $0\nu\beta\beta$ half-life that can be probed with SuperNEMO. This sensitivity depends

¹Supposing the $0\nu\beta\beta$ decay of ^{82}Se occurs through the exchange of a light Majorana neutrino.

on the number of selected background events.

Due to the time it would take to simulate every expected background contribution, the choice of a simplified model, containing only the most harmful backgrounds to the $0\nu\beta\beta$ decay search, was adopted. These backgrounds are the $2\nu\beta\beta$ decay, contaminations of the source in ^{208}Tl and ^{214}Bi , and contamination of the tracker gas with Radon. The number of natural isotope decay events expected depends on their activities inside the source foils (for ^{208}Tl and ^{214}Bi), or on the tracker's wires (for ^{222}Rn decaying in ^{214}Bi). It was necessary, during the design of the detector, to constrain the maximal tolerable activities, to guarantee a high sensitivity to the $0\nu\beta\beta$ disintegration [11]. The collaboration then established recommendations for maximum levels of the internal backgrounds, expressed in number of disintegrations per second, for a unit mass of $\beta\beta$ isotope, or for a unit volume of gas. These *specified activities* have been calculated in order to achieve the expected sensitivity of the final detector. The amount of expected double β decays is driven by its half-life value: the higher the half-life, the lower its contribution in the total number of expected background. For ^{82}Se sources, we take the $2\nu\beta\beta$ half-life measured by NEMO-3, $T_{1/2}^{2\nu} = 9.39 \pm 0.17$ (stat) ± 0.58 (syst) $\times 10^{19}$ years [12]. Moreover, we choose to consider the best limit set on the $0\nu\beta\beta$ process, of $T_{1/2}^{0\nu} > 2.5 \times 10^{23}$ y [12]. This value is given for illustration purposes only and is not used to estimate the sensitivity of the detector. In the Tab. 3.1 is summarised the expected number of signal and background events taking into account the specified activities, for the demonstrator and for the final detector. These figures are presented both for the

Process	Expected decays		Simulated decays
	Demonstrator	Final detector	
$0\nu\beta\beta$ ($T_{1/2}^{0\nu} = 2.5 \times 10^{23}$ y)	3.6×10^2	1.0×10^4	1.0×10^7
$2\nu\beta\beta$ ($T_{1/2}^{2\nu} = 9.39 \times 10^{19}$ y)	9.5×10^5	2.7×10^7	1.0×10^7
^{208}Tl ($\mathcal{A}^{\text{Tl}} = 2 \mu\text{Bq/kg}$)	1.1×10^3	3.1×10^4	1.0×10^7
^{214}Bi ($\mathcal{A}^{\text{Bi}} = 10 \mu\text{Bq/kg}$)	5.5×10^3	1.6×10^5	1.0×10^7
^{222}Rn ($\mathcal{A}^{\text{Rn}} = 0.15 \text{ mBq/m}^3$)	1.8×10^5	7.2×10^6	1.0×10^8

Table 3.1: Expected and simulated decays for different processes, both for the demonstrator (17.5 kg.y) and for the final detector exposures (500 kg.y), assuming target background activities are reached: $\mathcal{A}^{\text{Tl}} = 10 \mu\text{Bq/kg}$, $\mathcal{A}^{\text{Bi}} = 2 \mu\text{Bq/kg}$, $\mathcal{A}^{\text{Rn}} = 0.15 \text{ mBq/m}^3$. The measured half-life $T_{1/2}^{2\nu} = 9.39 \times 10^{19}$ y for ^{82}Se 's $2\nu\beta\beta$ is taken, and we assume $T_{1/2}^{0\nu} = 2.5 \times 10^{23}$ y [12].

SuperNEMO demonstrator and final detector. They represent the total number of disintegrations, without taking into account any technique to reject background, and for the full energy range. Nevertheless, the number of decays presented therein table are expected to be extremely reduced, notably by the application of event selections aimed at maximising the sensitivity to the excluded $0\nu\beta\beta$ half-life (Sec. 3.2). This effect is augmented by the fact that, for the current sensitivity analysis, we focus on a narrow energy window, called *region of interest*, whose usefulness is described in detail in Sec. 3.2. Therefore, to properly conduct this

sensitivity study, it was necessary to simulate a considerable number of events, so that the signal and backgrounds are correctly represented in the region of interest. Within the table is presented the total number of simulated events for each process.

3.1.1 The $0\nu\beta\beta$ signal

The SuperNEMO detector was designed to search for the never-observed $0\nu\beta\beta$ decay. In the following, we assume the underlying mechanism for this decay is the exchange of a light Majorana neutrino, the so-called mass mechanism (MM), as it is the most widespread. The hypothetical $0\nu\beta\beta$ signal would be detected as an excess of events in the region of interest, with respect to the predicted background contamination levels. Some 10^7 $0\nu\beta\beta$ events were simulated inside the source foils, using the DECAY0 software [13].

3.1.2 Inside detector backgrounds

In addition with the $0\nu\beta\beta$ decay, we simulated numerous types of backgrounds, that could mimic the searched signal.

3.1.2.1 Internal backgrounds

The so-called *internal backgrounds* stand for decays occurring inside the source foils, presenting the same signature as the $0\nu\beta\beta$ signal. These backgrounds, already introduced earlier, are the $2\nu\beta\beta$ of the source isotope, as well as disintegrations of ^{208}Tl and ^{214}Bi inside the source foils.

The $2\nu\beta\beta$ process

In the full energy range, the allowed $2\nu\beta\beta$ decay stands as the dominant internal background type. The corresponding two-electrons energy sum spectrum is a continuum, whose ending point should stand at $Q_{\beta\beta} = 2.99$ MeV, but is subtly shifted by the detector's energy resolution. We simulated 10^7 events of this decay inside the source foils, in the full energy window. However, above a certain energy value, the number of $2\nu\beta\beta$ events decreases very quickly. To offset this effect, we simulated additional 10^7 of this decay on a slightly reduced energy range, that is to say above 2 MeV. The second set of simulations is normalised with the first one. In this way, the lack of $2\nu\beta\beta$ simulated events in the high-energy tail is avoided, without requiring too high computational resources.

Source foils contamination by natural isotopes

As described in Sec. 2.2.1, after sources purification, remaining natural isotopes such as ^{208}Tl or ^{214}Bi can still be present inside the foils, constituting the principal internal source of background, with the $2\nu\beta\beta$ decay. We simulated 10^7 decays for each of the two isotopes, inside the source foils.

3.1.2.2 Tracker contamination by natural isotopes

The presence of gaseous ^{222}Rn inside the tracker, mainly deposited on the tracker wires, can produce events similar to internal ones. In fact, one of the progeny of ^{222}Rn , the ^{214}Bi , can decay on (or near) a foil, and appear with a two-electron ($2e$) topology, becoming hard to distinguish from a double beta decay candidate. As this isotope is distributed throughout the whole tracking detection volume, to study the experiment's sensitivity, we simulated a large quantity of this decay, that is to say 10^8 decays on the tracker wires.

3.1.3 External backgrounds

This background category is populated by the external γ -ray flux produced by radioactive isotope decays (mostly ^{40}K , ^{214}Bi and ^{208}Tl) in detector components or surrounding laboratory rocks, as well as neutron interactions in the external iron shield. As simulating external backgrounds would be very consuming in terms of computing resources due to their very low probability to produce a $2e$ -topology, it is important to ensure that they are necessary. The NEMO-3 experiment set a limit on the external background number of counts, of < 0.2 events in the $2e$ topology, for the energy range $[2.8; 3.2]$ MeV (two electrons energy sum), for an exposure of $34.3 \text{ kg}\cdot\text{y}$, with ^{100}Mo sources [14]. Radiopurity measurements of the SuperNEMO PMTs allow to conclude that the total ^{214}Bi activity is 35% lower than for those of NEMO-3 [15], which is encouraging. Unfortunately, these measurements also revealed that the total budget in ^{208}Tl isotope is 150% higher than NEMO-3. This could lead us to think that the external background contribution for SuperNEMO could be higher than that of NEMO-3. However, on that level, the most notorious difference between the two detectors is the fact that the SuperNEMO scintillator blocks are thicker than those of NEMO-3. Therefore, a gamma is more likely to be detected before crossing the source foils and, in that case, the event would not contribute to the background in the $2e$ channel. Even if the regions of interest are slightly different between these two experiments, it produces a negligible increase on the external background contribution². After all, given the fact that SuperNEMO is expected to be better than NEMO-3 at rejecting external background events, we consider that all external backgrounds from outside the foil, apart from ^{222}Rn in the tracking volume, are expected to be negligible, and were not simulated.

3.2 Event selection

For SuperNEMO, the $0\nu\beta\beta$ signature is two-electrons events, emitted simultaneously from the same vertex on the source foils, with an energy sum compatible with $Q_{\beta\beta} = 2.99 \text{ MeV}$ for the ^{82}Se . Therefore, we conducted this

²A study conducted by the SuperNEMO collaboration shown that at most 0.73 additional external background events would have been expected for the NEMO-3 detector, if instead of taking the $[2.8; 3.2]$ MeV energy range, we would have considered the $[2.7; 3.15]$ MeV region of interest.


analysis selecting only events matching the $2e$ topology, where a reconstructed particle is tagged as an electron if it has

- a vertex on the source foil,
- a reconstructed track inside the wire chamber,
- an associated calorimeter hit,
- and a final criterion depending on the case under consideration. In fact, as announced, this study also aims at considering two separate magnetic field cases. One where the magnetic field is aligned with the Z (vertical) axis of the detector with a uniform value of 25 Gauss, and one where the field is turned off. In the first case, particles such as electrons and positrons of a few MeV have a curved trajectory in the tracker. In the second case, the tracks of the particles may be similar to straight lines (not to mention the possible multiple scattering on the wires of the tracker). It is then necessary to adapt the selection of events to each case. We then consider an additional criterion: a particle is identified as an electron if its track has a negative curvature³. In the following, we present results where the magnetic field is turned on. The off-field study is addressed in Sec. 3.5.


All these selections represent the so-called *first-order* cut-offs. We present the total energy spectra for each simulated process, after application of these cut-offs, in Fig. 3.1. We present results for the specific case of the demonstrator (^{82}Se sources, 17.5 kg.y exposure), with the specified internal contamination levels ($\mathcal{A}^{\text{Tl}} = 2 \mu\text{Bq/kg}$ and $\mathcal{A}^{\text{Bi}} = 10 \mu\text{Bq/kg}$).

If the $0\nu\beta\beta$ decay is detected, its energy distribution would be a peak, located at the end-point of the $2\nu\beta\beta$ energy distribution, that is to say at the total available energy, $Q_{\beta\beta} = 2.99 \text{ MeV}$. In principle, this peak is infinitely thin, as the two electrons of this decay would share the total available energy. However, a widening of this peak is expected, due to the fact that there is not always the same energy lost by the electrons for the same path considered, and on the other hand that the path in the source is more or less long depending on the place of disintegration and the angle of emission of the electrons. Moreover, the two emitted electrons may lose some of their energy inside the detector, in the wire chamber for instance. This peak is also expected to be shifted towards small energies, by the calorimeter energy resolution and the straggling of energy losses in the detector. All this is at the origin of the asymmetrical energy distribution shown in the figure. Concerning the $2\nu\beta\beta$ energy spectrum, we present the events simulated in the total energy range, in addition with the $2\nu\beta\beta$ simulations with an energy $> 2 \text{ MeV}$. At the end, the two sets of simulation are normalised. Whatever their origin, either ^{222}Rn contaminations inside the tracker gas, or internal contaminations of the source foils, the two ^{214}Bi energy distributions have nearly the same shapes. The ^{208}Tl spectrum reveals the internal conversion of the 2.614 MeV gamma, emitted

³A trajectory is said to be negative if it has the same curvature as that of an electron moving from the source to the calorimeter, in a magnetic field oriented according to $+Z$



(a) Full energy range.



(b) Zoom on ROI.

Figure 3.1: Total energy spectra for the $0\nu\beta\beta$ signal and main backgrounds, for (a) the full energy range, and (b) for the $[2.7;3.15]$ MeV energy range, whose optimisation is discussed in Sec. 3.3. The $2\nu\beta\beta$ spectrum is normalised to $T_{1/2}^{2\nu} = 9.39 \times 10^{19}$ y, and ^{208}Tl , ^{214}Bi and ^{222}Rn backgrounds are normalised to the nominal activities. The amplitude of the $0\nu\beta\beta$ is arbitrarily set at the limit obtained with NEMO-3.

after ^{208}Tl β^- disintegrations. All spectra are normalised to the natural isotopes specified activities, and the half-lives values.

A widespread technique consists in constraining the $0\nu\beta\beta$ decay searches to a narrow energy range, the so-called *region of interest* (ROI), materialised by the two vertical dashed lines in the figure. In the following, we expose general principles leading to the determination of the best limit on $T_{1/2}^{0\nu}$, in the appropriate region

of interest. We illustrate the reasoning with the demonstrator, specified activities, on-magnetic field case. However, the technique presented remain valid for all exposures, internal contamination levels and field conditions. The impact of all these parameters is detailed in the rest of this chapter.

3.3 Demonstrator sensitivity to the $0\nu\beta\beta$ decay of ^{82}Se

The SuperNEMO demonstrator is designed to measure $\beta\beta$ decays of radioactive emitters. In case a the non-observation of this process, the demonstrator would set an upper-limit on the half-life $T_{1/2}^{0\nu}$, and on the effective neutrino mass $m_{\beta\beta}$.

3.3.1 Sensitivity to the $0\nu\beta\beta$ half-life

In case of the non-observation of a $0\nu\beta\beta$ signal, the expected upper limit on the half-life is provided for a given energy range $[E_{\min}; E_{\max}]$ for the two electrons energy sum, and depends on the characteristics of the detector. First, it depends on the signal detection efficiency, $\epsilon_{0\nu}$, in this energy window, secondly on the source isotope nature, as well as the detector's exposure $m \times t$. It follows

$$T_{1/2}^{0\nu} > \frac{N_A \ln 2}{M} \times \frac{\epsilon_{0\nu} \times m \times t}{N_{0\nu}^{\text{excl.}}}, \quad (3.1)$$

with N_A the Avogadro number, m the mass of isotope in the source foils, M its molar mass, and t the total time of data acquisition. $N_{0\nu}^{\text{excl.}}$ is the number of signal events excluded, calculated with the Feldman-Cousins statistics from the total expected number of background events. The Feldman-Cousins statistics [16] is a wide-used method in rare events search experiments, providing confidence intervals for upper limits in the case of background events following a Poissonian probability law. We use this method in the framework of this analysis to provide a limit, at 90% CL, on the number of excluded signal events $N_{0\nu}^{\text{excl.}}$, on the basis of the expected number of background events, given below.

- The $2\nu\beta\beta$ background

In Eq. (3.1), we defined the upper limit on $T_{1/2}^{0\nu}$ from the number of excluded signal events, and the signal selection efficiency $\epsilon_{0\nu}$. In a similar manner, we can define the number of expected $2\nu\beta\beta$ events, $N_{2\nu}$, from the half-life $T_{1/2}^{2\nu}$ and the $2\nu\beta\beta$ selection efficiency, $\epsilon_{2\nu}$, as

$$N_{2\nu} = \frac{N_A \ln 2}{M} \times \frac{\epsilon_{2\nu} \times m \times t}{T_{1/2}^{2\nu}}. \quad (3.2)$$

- Natural radioactive backgrounds

We consider the background massic activities $A_{\text{rad.}}$, and $\epsilon_{\text{rad.}}$ their selection efficiencies in a given energy window. The number of background events is therefore given, for the ^{208}Tl and ^{214}Bi internal contaminations, as

$$N_{\text{rad.}}^m = A_{\text{rad.}}^m \epsilon_{\text{rad.}}^m \times m \times t, \quad (3.3)$$

where $A_{\text{rad.}}$ is given in Bq/kg. Similarly, for the ^{222}Rn background,

$$N_{\text{rad.}}^V = A_{\text{rad.}}^V \epsilon_{\text{rad.}}^V \times V \times t, \quad (3.4)$$

with $V = 15.3 \text{ m}^3$ the tracker volume, and where $A_{\text{rad.}}$ represents here a volumic activity, given in Bq/ m^3 .

All these equations, similarly as Eq. (3.1), are valid for a given energy range $[E_{\text{min}}; E_{\text{max}}]$. To find the optimal energy interval for the search of the $0\nu\beta\beta$ decay, that is to say the one maximising the limit on $T_{1/2}^{0\nu}$, we must vary the E_{min} and E_{max} bounds.

As can be seen in Fig. 3.1, beyond 3 MeV, the number of background events becomes negligible compared with the $0\nu\beta\beta$ decay. Indeed, the ^{208}Tl background dominates at these energies, with 0.03 count for $E > 3.2 \text{ MeV}$. This is why the upper limit E_{max} of the energy interval has only a limited impact on the search for the best ROI. It is then natural to study mainly the influence of the upper limit E_{min} . Therefore, the selection efficiencies, entering in the calculation of the $T_{1/2}^{0\nu}$ upper limit, are presented in Fig. 3.2, as a function of the lower bound E_{min} . We remind




Figure 3.2: Efficiency spectra as a function of $E > E_{\text{min}}$, for the $0\nu\beta\beta$ signal (dashed black line) and for the main backgrounds (plain lines). The two vertical grey lines represent the final ROI optimised for the case of the demonstrator, taken the specified isotope activities.

a selection efficiency ϵ is the ratio of the number of selected events, to the number of simulated events. As a matter of fact, we look for an energy region where $\epsilon_{0\nu}$ is high, and where selection efficiencies for the background are low. This choice directly determines the best value for $T_{1/2}^{0\nu}$ (90% CL), whose variation as a function of E_{min} and E_{max} are presented in Fig. 3.3. As we said, the energy range chosen as the region of interest for the search of the $0\nu\beta\beta$ decays is the one maximising the $T_{1/2}^{0\nu}$. We found that, for the demonstrator exposure, with ^{82}Se sources, with a 25 Gauss magnetic field, and for the specified background activities, the best




Figure 3.3: Two-dimensional histogram showing the evolution of the $T_{1/2}^{0\nu}$ value as a function of the lower and upper energy bounds. The maximal upper limit of $T_{1/2}^{0\nu} > 5.7 \times 10^{24}$ y (90% CL) is retained, in the [2.7;3.15] MeV region of interest.

ROI is [2.7;3.15] MeV. The choice of the upper bound value does not impact the results for $E_{\text{max}} > 3.15$ MeV, so this value is kept in order to cut off possible external background. In this optimised energy range, the sensitivity expected for the SuperNEMO demonstrator stands at

$$T_{1/2}^{0\nu} > 5.7 \times 10^{24} \text{ y} \quad (90\% \text{CL}). \quad (3.5)$$

This result is compatible with previous SuperNEMO analysis [9].

3.3.2 Limit on the effective neutrino mass

The decay rate for the light Majorana exchange mechanism is given by:

$$(T_{1/2}^{0\nu})^{-1} = g_A^4 G^{0\nu} |M^{0\nu}|^2 \left| \frac{m_{\beta\beta}}{m_e} \right|^2. \quad (3.6)$$

where $G^{0\nu}$ is the two particles phase space factor, depending on $Q_{\beta\beta}$ and Z , $M^{0\nu}$ is the nuclear matrix elements for the $0\nu\beta\beta$ process, and $m_{\beta\beta}$ is the effective Majorana neutrino mass, defined as

$$\langle m_{\beta\beta} \rangle = \left| \sum_i m_i U_{ei}^2 \right|, \quad (3.7)$$

where m_i are the neutrino masses, and U_{ei}^2 is the mixing matrix. Therefore, the effective mass takes into account the neutrino mixing. Consequently, observing the $0\nu\beta\beta$ decay would not only prove the Majorana nature of neutrinos but, assuming the mass mechanism, could also help constraining the absolute neutrino masses. Given g_A , $G^{0\nu}$ and $M^{0\nu}$ [17][18][19][20][21][22][23][24][25][26], we find the SuperNEMO demonstrator could reach a limit on the effective neutrino mass of

$$\langle m_{\beta\beta} \rangle = [0.24 - 0.47] \text{ eV} . \quad (3.8)$$

In this section, we presented the general procedure leading to an optimised result on the $T_{1/2}^{0\nu}$ limit, and we gave the best limit on $T_{1/2}^{0\nu}$, driven by Eq. (3.1), for the demonstrator case. Thereafter, we discuss the results obtained for different detector exposures (demonstrator and final detector), and different internal background activities. Also, and this is the main purpose of this study, we discuss the influence of the presence of the magnetic field on the final detector's sensitivity.

3.4 Impact of sources contamination levels on the sensitivity

We study the impact of the isotope contamination levels (inside the source foils, as well as on the tracker's wires) on the $0\nu\beta\beta$ sensitivity. We also optimise additional event selections aimed at improving the final sensitivity result.

3.4.1 Influence of the contamination levels

Specified contamination levels have been established in order to achieve the $0\nu\beta\beta$ half-life target of $\sim 1 \times 10^{26}$ years for the final detector. The ^{82}Se demonstrator source is segmented in 34 foils, whose production was the responsibility of different laboratories (Dubna, LAPP and Tomsk). The sources have undergone different purification treatments, in order to investigate new techniques, and to compare them with those of NEMO-3. After the sources production and purification, preliminary measurements have been performed with the BiPo-3 detector to determine the actual ^{208}Tl and ^{214}Bi contamination levels inside the foils [27]. The level of radon emissions inside the tracker was also measured by the collaboration, for each of the four sections of the chamber, using a concentration line. We summarise all these contamination levels in Tab. 3.2, and give a comparison with the detector initial specifications. The targeted ^{208}Tl level is not reached, being almost 27 times higher than expected, and 3.0×10^4 internal Thallium events are expected in 2.5 years. Nevertheless, on average, the activity of the sources was improved by a factor of 2 compared to the ^{100}Mo sources of NEMO-3. In addition, valuable information has been accumulated on the different production techniques, which are of great importance for the final detector construction. In particular, the two best ^{208}Tl sources activities were reached by inverse chromatography, reaching a $20 \pm 10 \mu\text{Bq/kg}$ level, an improvement by a factor 5 compared to NEMO-3. This encourages for further investigations in this direction. The sensitivity of

	Specified activities	Measured activities
^{208}Tl	$2 \mu\text{Bq.kg}^{-1}$	$54 \mu\text{Bq.kg}^{-1}$ [26 - 102]
^{214}Bi	$10 \mu\text{Bq.kg}^{-1}$	$< 290 \mu\text{Bq.kg}^{-1}$
^{222}Rn	0.15 mBq.m^{-3}	$0.15 \pm 0.02 \text{ mBq.m}^{-3}$

Table 3.2: Real and targeted specified activities for the SuperNEMO detector. The ^{222}Rn tracker contamination is measured with a concentration line [28], extrapolated with a $2 \text{ m}^3/\text{h}$ flow rate. The limit on ^{214}Bi contamination is provided by BiPo measurements for a 90% CL [27].

BiPo detector only allowed to give an upper limit on the level of internal ^{214}Bi (an activity of $290 \mu\text{Bq/kg}$ would correspond to 1.6×10^5 internal Bismuth events in 2.5 years). Precise measurements are expected from the demonstrator calibration. Radon emissions from the tracker were also measured, and extrapolated with an air flow rate of $2 \text{ m}^3/\text{h}$ inside the chamber, showing the targeted level of 0.15 mBq.m^{-3} was reached.

In Sec. 3.3, we developed the general procedure allowing to set a 90% confidence interval limit on $T_{1/2}^{0\nu}$. For the demonstrator, supposing the specified activities are reached, the demonstrator would achieve a sensitivity of 5.7×10^{24} years on the searched decay, in 2.5 years of data acquisition, with 7 kg of ^{82}Se . This sensitivity could be affected by the higher-than-specified level of contaminations, measured by BiPo, inside the source foils.

In this sub-section, four distinct levels of internal contaminations are considered:

- the *zero activities* case, a hypothetical case where the source foils and the tracker are non contaminated at all,
- the *specified activities* case, where the targeted level of contaminations would have been reached,
- and two *measured* cases, that takes into account the measured levels or limits on the contaminations at 90% CL. As the ^{214}Bi activity is provided by BiPo measurements as an upper limit, it is possible for this level to be lower than $290 \mu\text{Bq/kg}$. We therefore present the results, either for sources that would not be contaminated by this isotope (the *without ^{214}Bi* case), or considering that the activity reached is $290 \mu\text{Bq/kg}$ (*with ^{214}Bi*).

The fact that we are showing results for a hypothetical zero isotope contamination is to illustrate an important statistical phenomenon. Indeed, we notice that there is no difference in sensitivity, or ROI, between the first two cases of contamination, which may seem surprising at first glance. This is explained by the Feldman-Cousins statistics employed to determine the number of excluded signal events, $N_{0\nu}^{\text{excl.}}$, given the number of observed background events (defined from Eq. (3.2) to Eq. (3.4)).

Clarifications on Feldman-Cousins statistics When the expected number of background events is negligible (which is the case for the two first levels presented), the probability p to observe n_s signal events, expecting s events, is given by the Poisson distribution

$$p = \frac{e^{-s} s^{n_s}}{n_s!}. \quad (3.9)$$

Let's now put ourselves in the situation where no signal event is observed - that is what we assume to put an upper limit on the $0\nu\beta\beta$ half-life. Then $n_s \rightarrow 0$, and $p \rightarrow e^{-s}$. If zero signal event is *observed*, it is incorrect to assume that zero signal events were *produced* during the experiment. We only can say that no signal event has been observed *a priori*. To account for this particular case, the quantity s should no longer be viewed as the number of expected signal events, but as the number of excluded signal events, $N_{0\nu}^{\text{excl.}}$. In the end, for a negligible expected number of background events, and no signal event observed, we can set an upper limit on the number of excluded signal events, excluding values for which $p < \alpha$. Taking a 90% confidence interval, that is to say $\alpha = 10\%$, we obtain $s \leq 2.303$.

Tab. 3.3 summarises the expected number of background for each non-zero contamination case presented in Fig. 3.4. For each case, the ROI is optimised.

Activity ROI	Specified [2.7;3.15] MeV	Measured (w/o ^{214}Bi) [2.75;2.95] MeV	Measured (w/ ^{214}Bi) [2.7;2.9] MeV
$\epsilon_{0\nu}$ (%)	14.7	11.3	14.3
$2\nu\beta\beta$	0.418	0.122	0.418
^{208}Tl	0.0475	0.688	0.699
^{214}Bi	0.0546	0	1.55
^{222}Rn	0.292	0.173	0.287

Table 3.3: Expected number of background events in the $2e$ topology, in the [2.7;3.15] MeV ROI, for the SuperNEMO demonstrator (17.5 kg.y), for specified activities. The selection efficiency of $0\nu\beta\beta$ events, $\epsilon_{0\nu}$, is also given.

A direct consequence of the statement on Feldman-Cousins statistics is that, considering the background levels for the two first cases are negligible (or null) in the region of interest, they both reach this limit of 2.303 on the excluded number of signal events. For the two measured activities cases considered, the expected number of background events increases, although the narrow ROI for the case without ^{214}Bi limits it. Indeed, both region of interests are highly reduced, especially for the case without ^{214}Bi , where the lower bound is increased from 2.7 to 2.75 MeV. As this 50 keV wide energy region is populated with a non-negligible number of background events, this change in E_{min} usefully reduces the $2\nu\beta\beta$ background contribution, thereby limiting the increase of total expected number of background.

We show in Fig. 3.4 the 90 % CL $T_{1/2}^{0\nu}$ limit as a function of the contamination levels, as well as the corresponding chosen region of interest. As expected from previous conclusions on the Feldman-Cousins statistics, no difference is observed




Figure 3.4: The 90% CL limit on the $0\nu\beta\beta$ half-life (top pad), and the corresponding ROI (bottom pad), as a function of the contamination level considered. For the *zero activities* case, we consider hypothetical contamination levels where $\mathcal{A}^{\text{Bi}} = \mathcal{A}^{\text{Tl}} = 0$ Bq/kg. The *specified activities* are presented in Tab. 3.2. The *measured activities*, provided by the BiPo detector [27], are presented in the same table. We consider successively a null ^{214}Bi contamination (*measured act. w/o ^{214}Bi*), or equals to the $290\mu\text{Bq/kg}$ upper limit (*measured act. w/ ^{214}Bi*).

in terms of half-life limits between the zero and specified activities cases. However, when considering the measured internal activities, a decrease in sensitivity is observed. Indeed, the number of background events in the ROI is no more negligible, and influence significantly the value of $T_{1/2}^{0\nu}$, decreasing the experiment's sensitivity by 23% (without ^{214}Bi) and 37% (with ^{214}Bi).

The degradation of the limit on the $0\nu\beta\beta$ half-life with the level of contamination remains acceptable. However, we can try improving the situation by exploring new background rejection techniques. This would be especially useful for the final detector case, where a slight increase in internal contaminations could be highly harmful, all the more so as the upper limit given for ^{214}Bi turns out to be the true contamination level.

3.4.2 Optimisation of the event selection

The measured level of ^{208}Tl isotope inside the source foils is greater than the specifications. Moreover, an upper limit, higher than the specified level for SuperNEMO, has been set by the BiPo detector, regarding the internal ^{214}Bi level. Consequently, we can imagine implementing a tighter selection to reject more background. Most of the double beta experiments are only sensitive to the total electron energy sum. The unique SuperNEMO tracko-calorimetry technology confers the experiment the ability to characterise single particles (individual energies, emission angles...). Based on previous studies [9] [29], *topological cuts*, relying on these

additional observables, have been set up. They are especially designed to reject events where the two electrons are not emitted simultaneously, or from the same location on the source foils.

The internal probability Based on time-of-flight (TOF) computation, the internal probability (P_{int}) is derived from the internal χ^2 (see details in Sec. 2.5.1). In Fig. 3.5 are presented the internal probability spectra for the $0\nu\beta\beta$ signal and all background processes, after the first-order selections. These distributions are






Figure 3.5: Internal probabilities for all processes. First-order cuts have been applied. $\beta\beta$ distributions are normalised to the half-lives, and background processes are normalised to the specified activities.



normalised to the double beta half-lives, and the nominal activities. Equivalent distributions, but with different ^{214}Bi and ^{208}Tl contamination levels, can be derived for the case of measured activities. The internal probability distributions for the $0\nu\beta\beta$ and $2\nu\beta\beta$ processes follow the expected flat distribution for electrons emitted simultaneously from the source. As internal Bismuth disintegration actually takes place inside the sources, the ^{214}Bi distribution is also flat. The same could have been assumed for Thallium, however, the distribution is distorted at low internal probabilities. This might be explained by the existence of a metastable excited state ($\tau_{1/2} = 294\text{ps}$) of the daughter nuclei, which would slightly delay the second electron emitted via internal conversion. This feature is addressed in detail in Chap. 4. The Radon, being a non-internal background, presents a peak at low internal probabilities.

We want to evaluate the influence of a cut-off on the simulations using internal probability as a rejection criterion: simulated events are selected for P_{int} values upper than a given limit. The standard value applied in NEMO-3 analyses was $P_{int} > 4\%$. We wish to establish the most adequate P_{int} selection level for the SuperNEMO demonstrator. For each internal probability cut-off applied to the simulations, we evaluate the $T_{1/2}^{0\nu}$ at a 90 % confidence interval, as well as the

best ROI. The internal probability value that will be chosen will be the one that maximises the sensitivity, and depends on the internal level of contamination. To better understand this optimisation, let us present the $\epsilon_{0\nu}$ values for each P_{int} selection case, in Fig. 3.6a, both for the specified and measured contamination levels. For the latter, we assume the Bismuth upper limit of 290 $\mu\text{Bq/kg}$ activity is


 (a) $0\nu\beta\beta$ selection efficiency.


(b) ROI optimisation.


 (c) Limit on $T_{1/2}^{0\nu}$.


(d) Total expected number of background.




Figure 3.6: The $0\nu\beta\beta$ selection efficiency, $\epsilon_{0\nu}$ (a), evolution of the region of interests (b), best limits set on $T_{1/2}^{0\nu}$ at 90% CL (c), and total number of expected background (d), as a function of the cut-off applied on internal probability, P_{int} . The ROI is optimised for each P_{int} value. Results are displayed for two contamination levels: the specified (blue) and the measured (orange) activities (taking into account the upper limit provided for ^{214}Bi). An exposure of 17.5 kg.y is considered. Two vertical dashed lines in (a) and (c) display the best P_{int} selections to be applied in order to improve the $T_{1/2}^{0\nu}$ sensitivity of the experiment.

reached. As expected, in both cases, the $\epsilon_{0\nu}$ selection efficiency globally decreases with the P_{int} cut-off applied. However, for the measured contamination case, we observe a slight increase of the $0\nu\beta\beta$ selection efficiency, for $P_{int} > 1\%$. Fig. 3.6b, showing the variations of the regions of interest as a function of the P_{int} rejection criterion, allow to understand this observation. For the measured activities case, when the minimal acceptable P_{int} is changed from 0 to 1 %, then the ROI upper bound increases from 2.9 to 3.05 MeV. Usually, the variation of this bound does not have such a great impact on the event selection. Nevertheless, in the measured activities case, for a $P_{int} > 0\%$ level, the ROI is optimised at

the narrow [2.7;2.9] MeV interval, where the upper bound is located in an energy region still populated by signal and background (see Fig. 3.4). Therefore, even small variations in this ROI has a great impact on the results, explaining the local increase in efficiency. For P_{int} selections greater than 1 %, we come back in cases where the upper limit of the ROI no longer has an impact on $\epsilon_{0\nu}$. For the specified activities case, the ROI bounds are stable, not impacting the selection efficiency. In Fig. 3.6d, the total expected number of background is displayed. For specified activities, this number decreases with the P_{int} cut-off, as the ROI is stable. However, this number tends to reach a plateau after increasing, again explained by the ROI variations.

As the limit set on $T_{1/2}^{0\nu}$ depends directly on $\epsilon_{0\nu}$, the variations presented in Fig. 3.6a fully explain the results displayed in Fig. 3.6c, presenting the evolution of $T_{1/2}^{0\nu}$ with the internal probability selection level. The sensitivities displayed for a 0% cut-off on P_{int} of course correspond to the results given in Fig. 3.4. The main conclusion is that this rejection criterion has only a limited impact on the improvement of $T_{1/2}^{0\nu}$ sensitivity, because of the very low contamination levels considered. Indeed, paradoxically, the selection on internal probability worth it only if there is enough background events to be rejected. The former 4 % value recommended by the NEMO-3 analysis is then too high for the SuperNEMO source contamination.

NEMO-3 analyses also used the distance between the reconstructed vertices on the source foils as a background rejection criterion. As we shown that the additional P_{int} cut-off is poorly adapted for the low activities of SuperNEMO sources, it is interesting to know if we can improve the results by using this second selection.

Vertices distance As discussed in Sec. 3.2, an electron is defined as a reconstructed track inside the wire chamber, with one extremity on a calorimeter block, and another on a source foil. Therefore, the $2e$ topology involves two distinct calorimeter hits, two reconstructed tracks, and two foil vertices. Thanks to the trajectory fitting algorithm, we have access to the (Y, Z) coordinates of the latter, and by extension, to the distance between them. In the previous studies, the choice was made to look at the effect of this selection, separately on the Y (perpendicular to the wires) and Z (parallel to the wires) directions. We choose to follow the same approach, and we give the results for a cut along the Z axis, but the conclusions would remain valid for the Y direction. Fig. 3.7 shows the distributions of the absolute value of the distance between foil vertices for each process studied. We would use this information in order to maximise the double β decays to be selected, while rejecting natural isotope disintegrations.

Fig. 3.8 displays the limit set on $T_{1/2}^{0\nu}$ as a function of this event selection, allowing to study the impact of the vertices distance cut-off on the final sensitivity. In the same way as previous paragraph, we study the influence of the level of contaminations by showing results for the two specified and measured activities. The two distributions reach a plateau, corresponding to the sensitivities achieved with the first-order cuts (Fig. 3.4). Regardless of the level of contamination, the number of background events is too low in the ROI for the $|\Delta Z|$ cut-off to




Figure 3.7: Distance along the Z direction between the vertices of the 2 reconstructed electrons, for each process considered. The $2\nu\beta\beta$ spectrum is normalised to $T_{1/2}^{2\nu} = 9.39 \times 10^{19}$ y, and ^{208}Tl , ^{214}Bi and ^{222}Rn backgrounds are normalised to the nominal activities. The amplitude of the $0\nu\beta\beta$ is arbitrarily set at the limit obtained with NEMO-3. No energy cut is applied.




Figure 3.8: Limit reached for $T_{1/2}^{0\nu}$ (90% CL), as a function of the cut-off applied on the Z distance between vertices, $|\Delta Z|$. Results are displayed for specified (blue), and measured contaminations taking into account the upper limit for ^{214}Bi activity (orange). The ROI is optimised for each P_{int} value. No improvement in sensitivity is obtained with this event selection.

be worthwhile. Therefore, the cut-off only degrades the sensitivity, as does the

3. SENSITIVITY OF THE SUPERNEMO DEMONSTRATOR TO THE $0\nu\beta\beta$

selection on internal probability. The same conclusions apply to the $|\Delta Y|$ cut-off.

However, such a cut-off could be useful for rejecting unexpected background (coincidence between independent events, for instance). For example, we note that a cut-off at $|\Delta Z| < 80$ mm does not significantly degrade sensitivity. In practice, a selection on vertex distance will always be applied, even if it is very loose.

The idea of having implemented these two selections (on the internal probability and on the distance between vertices) comes from a previous NEMO-3 analysis on the background rejection. For the SuperNEMO demonstrator case, the levels of contaminations we are dealing with is remarkably low for most of the topological cut-offs to be worth applying. However, in practice, applying loose topological selections on the data remains necessary, especially to reject external background events. The minimal cut-off level to be applied is $P_{int} > 4\%$ and $|\Delta Z| < 80$ mm (samely for $|\Delta Y|$).

For future studies, it is useful to give the efficiencies of these loose selections, for the signal and for each background considered (Tab. 3.4a), as well as the expected number of background in the ROI (Tab. 3.4b).

Cut-off	First-order cuts (%)	Internal probability (%) $P_{int} > 4\%$	Vertex distance (%) $ \Delta Z < 80$ mm
$0\nu\beta\beta$	26.9	25.3	24.7
$2\nu\beta\beta$	9.15	8.56	8.21
^{208}Tl	0.106	0.0889	0.0846
^{214}Bi	0.168	0.151	0.144
^{222}Rn	0.0177	7.91×10^{-3}	5.34×10^{-3}

(a) Selection efficiencies (number of selected $2e$ topologies compared with the total number of simulated decays), for the three levels of selection (first-order, P_{int} and vertex distance), in the full energy range.

Activity	Specified		Measured (w/ ^{214}Bi)	
Cut-off	$P_{int} > 4\%$	$ \Delta Z < 80$ mm	$P_{int} > 4\%$	$ \Delta Z < 80$ mm
ROI (MeV)	[2.7;3.15]	[2.7;3.15]	[2.7;3.25]	[2.7;3.3]
$\epsilon_{0\nu} (\%)$	14.1	13.9	14.1	13.9
$2\nu\beta\beta$	0.392	0.383	0.392	0.383
^{208}Tl	0.0338	0.0323	1.08	1.09
^{214}Bi	0.0491	0.0491	1.42	1.42
^{222}Rn	0.115	0.0782	0.115	0.0782

(b) Expected number of background events in the optimised ROI, for successive application of topological selections. The selection efficiency of $0\nu\beta\beta$ events in the ROI, $\epsilon_{0\nu}$, is also given. Specified and measured activities (taking into account the upper limit for ^{214}Bi contamination) are considered. The SuperNEMO demonstrator exposure of 17.5 kg.y is taken.

Table 3.4: The selection efficiencies and expected number of background events for the topological selections.

Regions of interest are optimised for each selection. The topological cuts have a huge impact on Radon, as they are especially designed to reject non-internal

events. They are also efficient in rejecting Thallium internal events, because of the existence of a metastable excited state, described earlier. A special technique to reject efficiently ^{208}Tl background is also addressed in Chapter 4.

After the topological cut-off optimisation, the SuperNEMO demonstrator would reach a sensitivity of $T_{1/2}^{0\nu} > 5.6 \times 10^{24}$ y, and a corresponding effective neutrino mass of $\langle m_{\beta\beta} \rangle < [0.25 - 0.48]$ MeV. After looking at the effect of contaminations on the sensitivity, we review the influence of the magnetic field inside the detector.

3.5 Impact of the magnetic field on the sensitivity

The SuperNEMO demonstrator was originally designed with a copper coil, similarly to NEMO-3, delivering a magnetic field inside the tracker volume, aiming to provide an electron/positron discrimination. This 25 Gauss magnetic field is high enough to bend the trajectory of the few MeV electrons and positrons of interest for SuperNEMO, without too strongly preventing them from reaching the calorimeter. In practice, this magnetic field is mainly used to identify and reject the electron-positron pairs created by high energy γ 's, themselves emitted after a neutron capture. However, as explained in sub-section 3.1.3, we choose to not consider the contribution of this external background. We therefore focus on evaluating the influence of the presence of the magnetic field on the rejection of internal and wire chamber backgrounds.

3.5.1 Simulations of the magnetic field inside the demonstrator and reconstructed track fit

In order to study the influence of the magnetic field on the ^{82}Se $0\nu\beta\beta$ sensitivity, the simulations and reconstructions of decays described in Sec. 3.1 have been performed in two different conditions.

- Simulations with a uniform 25 Gauss magnetic field (following recommendations [9]). Results about the final sensitivity achieved in this condition have already been presented earlier in this chapter. The possible variations of the field intensity, mainly due to the calorimeter magnetic shields, are not taken into account for these simulations. This will be discussed in sub-section 3.5.4.
- Simulations where the magnetic field is turned off.

Each magnetic field condition has the same number of simulated events, as summed up in Tab. 3.1. Depending on the case considered, the electrons do not have the same trajectory curvature. In the first uniform on-field case, the best track fit is performed by an helix. In the second off-field case, the trajectory of an electron is modelised by a straight line. The fitting algorithm is thus modified to match line trajectories.

3.5.2 Impact of the magnetic field on signal and background selections

Among the various event selection criteria considered in Sec. 3.2, the one on the trajectory curvature is of primary importance with regard to the influence of the magnetic field on the final sensitivity of the detector. Indeed, when the magnetic field is switched on, the charged particles of few MeV (as electrons and positrons) have curved trajectories. A particle is then identified as an electron when the trajectory fitting results in a negative curvature. When the magnetic field is switched off, the trajectory of the charged particles takes place in a straight line⁴. This last selection criterion on the track curvature is then no longer applied. Consequently, the number of identified $2e$ topologies, selected by the first-order cuts, are increased, for the signal and background simulated events. To illustrate this effect, we give in Fig. 3.9 the selection efficiencies of signal and background as a function of the $2e$ total energy, for the two cases of magnetic field presented above. The two coloured stripes represent the corresponding region of interests,




Figure 3.9: The $0\nu\beta\beta$ selection efficiency (plain line), and expected number of background events (dashed line), as a function of the $2e$ energy, for the on-field (blue) and the off-field (orange) cases. Results are presented for a 17.5 kg.y exposure, for the specified isotope activities. The two corresponding region of interests are also displayed by coloured stripes. Here no optimised topological cut-offs have been applied.

of [2.7;3.15] MeV for the *on-field* case, and [2.75;3.2] MeV for the *off-field* case. These results are presented for the specified contamination levels, but remain valid in all cases. We clearly notice that, for the total energy interval [0;4] MeV, the number of signal and background selected events is higher in the off-field case, which confirms what we were saying above.

⁴In saying this, we do not take into account possible deviations in the trajectory of the particles, due in particular to multiple scattering in the tracker.

Following Eq. (3.1), the ratio $\epsilon_{0\nu}/N_{0\nu}^{\text{excl.}}$ in the ROI impacts the final sensitivity. Nevertheless, as we explained in the previous section, the low contamination level considered causes $N_{0\nu}^{\text{excl.}}$ to reach the 2.303 limit. Thus, only variations of $\epsilon_{0\nu}$ in the ROI, between on-field and off-field cases, are the source of modifications on the final sensitivity. Integrating $\epsilon_{0\nu}$ over the ROI, we obtain $\epsilon_{0\nu}^{\text{on-field}} = 0.15\% > \epsilon_{0\nu}^{\text{off-field}} = 0.12\%$. The selection efficiency for the on-field case is favoured by the lower bound of the ROI. In fact, the further this lower bound is shifted towards lower energies, the greater the selection efficiency. Besides, slight variations of the upper bound have almost no impact. As expected, this results in a decrease in sensitivity when the field is switched off, giving

$$T_{1/2}^{0\nu} > 4.80 \times 10^{24} \text{ y} \quad (90\% \text{CL}) \text{ (off-field)}. \quad (3.10)$$

We present in Tab. 3.5 the selection efficiencies of the signal and backgrounds, in the ROI, for the two field cases. We principally notice that the ^{208}Tl and ^{222}Rn

Process	On-field [2.7;3.15] MeV	Off-field [2.75;3.2] MeV
$\epsilon_{0\nu}$ (%)	14.7	12.4
$2\nu\beta\beta$	0.40	0.063
^{208}Tl	0.047	0.060
^{214}Bi	0.054	0.045
^{222}Rn	0.29	0.55

Table 3.5: Expected number of background events in the $2e$ topology, in the ROI, for the SuperNEMO demonstrator (17.5 kg.y), for specified activities. The selection efficiency of $0\nu\beta\beta$ events, $\epsilon_{0\nu}$, is also given. The two on- and off-field cases are compared.

contributions are higher in the second case. As concluded in Sec. 3.4, topological selections are especially efficient in rejecting these two backgrounds. Therefore, the application of these additional cut-offs, for the off-field case, could be interesting, in order to increase the sensitivity. Following the work presented in the previous section, we optimise these selections for the particular off-field case, both for the specified and measured contamination levels⁵. The results in the sensitivity are summarised in Fig. 3.10. The left part of the panel gives information on the evolution of sensitivity, when only the first-order cut-offs are applied. We come back to the conclusions given above: when the magnetic field is switched-off, we lose sensitivity, regardless of the level of contamination considered. On the right side of the figure, we present the results when the topological cuts are applied. For the on-field case, the addition of these selections have almost no effect on the sensitivity, as concluded in sub-section 3.4.2. However, as predicted, we are beginning to see the usefulness of these selections in the off-field case, as a higher number of ^{208}Tl and ^{222}Rn events passed the first-order selections. For instance, for the specification case, $T_{1/2}^{0\nu}$ goes from $4.80 \times 10^{24} \text{ y}$ to $6.1 \times 10^{24} \text{ y}$, an improvement of $\sim 30\%$.

⁵As done in sub-section 3.4.2, for the Bismuth measured contamination, we consider here the upper limit where $\mathcal{A}^{\text{Bi}} = 290 \mu\text{Bq/kg}$.




Figure 3.10: $T_{1/2}^{0\nu}$ (90% CL) considering various conditions: on- and off-field (white and gray stripes), first-order and addition of topological cut-offs (left/right parts of the panel), specified and measured activities (blue and orange triangle markers). The measured activities are $\mathcal{A}^{\text{Tl}} = 54 \mu\text{Bq/kg}$, $\mathcal{A}^{\text{Bi}} = 290 \mu\text{Bq/kg}$ and $\mathcal{A}^{\text{Rn}} = 0.15 \text{ mBq/m}^3$.

Finally, even if the absence of the magnetic field has the effect of reducing the sensitivity to the $0\nu\beta\beta$ decay, topological cuts allow this effect to be compensated for, making it possible to reach higher values of $T_{1/2}^{0\nu}$.

3.5.3 Influence of the magnetic field on optical modules and reconstruction efficiency

In the previous sub-section, a comparative study has been lead to evaluate the influence of the presence of a magnetic field on the event selection, and thus on the final sensitivity. However, as things stand now, some features of the demonstrator are not yet implemented in the simulations, and could have a great impact on the results presented above. In particular, studies have been lead by the collaboration to evaluate the influence a 25 Gauss magnetic field on the optical modules, as well as on the event reconstruction [9][10].

SuperNEMO PMTs are protected from the external magnetic field by individual, cylindrical, iron shields. Unfortunately, the latter do not perfectly protect the PMTs, and a residual magnetic field is measured inside the shieldings, leading to losses in charge collected by PMTs close to 8%. This study also revealed the energy resolution would be worsened with a relative decrease of 3% of the initial value of 8% at 1 MeV. Moreover, the PMTs shieldings could themselves severely impact the shape of the field lines, as well as its strength. In fact, with a 25 Gauss magnetic field generated by the copper coil, the magnetic shields are responsible

for the field strength decreasing, and barely 10 G is expected near the source foils. Worse, the magnetic field strength decreases very quickly as we get closer to the calorimeter walls where nearly 0G could be expected. The reconstruction efficiency could therefore be greatly impacted: the magnetic field intensity varying from the source foils to the calorimeter wall, electrons trajectory curvatures are not constant, and the track-fitting algorithm is less performing. A non correct description of the distribution of the magnetic field would more strongly impact low-energy electrons.

In the light of these conclusions, it could be interesting to study the evolution of the sensitivity, considering field simulations with more realistic variations inside the detector.

3.5.4 Simulations with a non-uniform magnetic field

Simulations with a 25 Gauss *mapped* magnetic field have been performed, taking into account more realistic variations of the field inside the detector [30]. Unfortunately, not all isotopes could be simulated with this magnetic field configuration. In particular, radon, as it is present in the entire wire chamber, would have required too many additional storing resources. Thus, final conclusions on the sensitivity can't be given. However, it is possible to assess the selection efficiencies of the different processes, and thus get an idea of the influence of realistic variations of the field on the final results.

Tab. 3.6 compares the selection efficiencies, for the three field cases (uniform field, mapped field and off-field), in the total energy range [0;4] MeV. The mapped

Process	Selection efficiencies (%)		
	On-field	Off-field	Mapped-field
$0\nu\beta\beta$	25.8	29.7	20.0
$2\nu\beta\beta$	8.64	9.93	6.51
^{208}Tl	0.0986	0.152	0.0910
^{214}Bi	0.153	0.220	0.134

Table 3.6: Signal and background selection efficiencies of on-field, off-field and mapped-field cases, in the energy range [0;4] MeV.

field case has lower selection efficiencies in the total energy range, compared with uniform field simulations. As announced in the previous sub-section, the magnetic shields distort the field intensity accross the detector. Therefore, the fitting algorithm is less performant in identifying particles with a negative curvature inside the tracker.

3.6 Searching for the Neodymium-150 $0\nu\beta\beta$ decay

This study was conducted jointly with the PhD student Axel Pin, from CENBG [31]. Although we both worked on the whole of the analysis, I presented

in detail, in the previous sections, the results regarding the influence of the magnetic field. Meanwhile, Axel Pin present in detail the possibility of changing the Selenium material by other $\beta\beta$ isotopes. Indeed, on the model of the NEMO-3 detector, which housed 6.914 kg of ^{100}Mo and 0.932 kg of ^{82}Se , the SuperNEMO detector has the technical possibility of exchanging the source material and study several $\beta\beta$ isotopes. In the case SuperNEMO demonstrates the feasibility of a large-scale tracko-calorimeter experiment, it would be natural to evaluate the sensitivity of SuperNEMO to the $0\nu\beta\beta$ decay of other isotopes than ^{82}Se .

3.6.1 Searching for the $0\nu\beta\beta$ of other isotopes

As explained in Chapter 2, the 2.615 MeV γ -ray produced in the decay of ^{208}Tl is a troublesome source of background. It is then important to select $\beta\beta$ candidates with a $Q_{\beta\beta}$ value above this transition. The natural isotopic abundance is another useful criterion, and typically considering only isotopic abundances greater than 2% is a good basis for making a choice. Five nuclei satisfy these two criteria: ^{116}Cd , ^{82}Se , ^{100}Mo , ^{96}Zr and ^{150}Nd (with respective $Q_{\beta\beta}$ values of 2.80, 2.99, 3.03, 3.35 and 3.36 MeV and respective isotope abundance values of 7.5, 9.2, 9.6, 2.8 and 5.6 % [32]). Given the availability of ^{100}Mo , much effort has been focused by the NEMO-3 collaboration on this isotope (it had already been studied by the NEMO-2 prototype [33]). As the ^{150}Nd isotope has the highest $Q_{\beta\beta}$ value, the current section focuses on evaluating the SuperNEMO sensitivity to the $0\nu\beta\beta$ decay of this isotope, supposing we have several kg at our disposal.

3.6.2 Sensitivity to the $0\nu\beta\beta$ of ^{150}Nd

Until recently, Neodymium was not enrichable in large quantities. Recent developments have resulted in the production of several grams of enriched Neodymium, making this $\beta\beta$ isotope interesting for the search of $0\nu\beta\beta$. The best limit for the search for neutrinoless double β decay of ^{150}Nd was reached by the NEMO-3 detector, using 36.6 g of enriched material for 5.25 years of data acquisition. The collaboration achieved $T_{1/2}^{0\nu} < 2.0 \times 10^{23}$ y (90 % CL), corresponding to an upper limit on the effective neutrino mass of $\langle m_{\beta\beta} \rangle < [1.6 - 5.3]$ eV. The collaboration also measured the $2\nu\beta\beta$ half-life, with $T_{1/2}^{2\nu} = [9.34 \pm 0.22 \text{ (stat.)} \pm 0.60^{+0.62}_{-0.60} \text{ (syst.)}] \times 10^{18}$ y [34].

We wish to determine the limit on the $0\nu\beta\beta$ of the ^{150}Nd that could be reached by the SuperNEMO demonstrator. We lead this study considering the specified activities given for the ^{82}Se sources are reached. We use simulations with the 25 Gauss uniform magnetic field. Fig. 3.11 presents the normalised energy distributions for the $2e$ topologies selected after application of first-order and topological selections. Signal and background selection efficiencies for ^{150}Nd sources, in the total energy range, are given in Tab. 3.7. In this energy range, the background is dominated by the $2\nu\beta\beta$ decay. The selection efficiencies of backgrounds are lower for ^{150}Nd sources than for ^{82}Se sources. This is caused by the higher number of protons in the Neodymium nucleus which induces a stronger Coulombian effect. Indeed, internal electrons are more likely to interact with




Figure 3.11: Total energy spectra for the $0\nu\beta\beta$ signal and main backgrounds, for ^{150}Nd sources. The $2\nu\beta\beta$ spectrum is normalised to $T_{1/2}^{2\nu} = 9.39 \times 10^{19}$ y, and ^{208}Tl , ^{214}Bi and ^{222}Rn backgrounds are normalised to the nominal activities. The amplitude of the $0\nu\beta\beta$ is arbitrarily set at the limit obtained with NEMO-3. We assume a 17.5 kg.y exposure. First-order and optimised topological cuts have been applied. The ROI of [3.1;3.55] MeV is represented by two vertical dashed lines.

Process	Selection efficiencies (%)	
	Selenium	Neodymium
$0\nu\beta\beta$	25.8	25.5
$2\nu\beta\beta$	8.21	8.11
^{208}Tl	0.0846	0.0749
^{214}Bi	0.144	0.138
^{222}Rn	5.34×10^{-3}	5.34×10^{-3}

Table 3.7: Selection efficiencies in the full energy range [0;4] MeV, for ^{82}Se and ^{150}Nd sources. First-order and optimised topological cuts have been applied.

the source material by Coulombian interactions, which is particularly true for low energy electrons. We used the same Radon simulations as for ^{82}Se sources, neglecting possible effects due to a higher proton number.

In Tab. 3.8 we give the expected number of background events in the optimised ROI [3.1;3.55] MeV. The selection efficiency of the $0\nu\beta\beta$ decay in this energy range is also given. Although the $2\nu\beta\beta$ half-life of the ^{150}Nd is lower than that of the ^{82}Se the number of $2\nu\beta\beta$ events in the ROI remains low. Indeed, thanks to the Coulombian effects described above, this process has a limited contribution at high energy. The high energy of transition $Q_{\beta\beta} = 3.36$ MeV of ^{150}Nd implies that the contributions of ^{214}Bi and ^{222}Rn are very small, or even zero. The $2\nu\beta\beta$ and ^{208}Tl events are therefore the majority contributors to the background.

Process	Selenium [2.7;3.15] MeV	Neodymium [3.1;3.55] MeV
$\epsilon_{0\nu}$ (%)	14.4	10.3
$2\nu\beta\beta$	0.39	0.28
^{208}Tl	0.044	0.029
^{214}Bi	0.053	5.6×10^{-4}
^{222}Rn	0.20	0.0

Table 3.8: Expected number of background events in the optimised region of interests, for ^{82}Se and ^{150}Nd sources. The $2\nu\beta\beta$ half-life taken as $T_{1/2}^{2\nu} = 9.39 \times 10^{19}$ y, and the nominal background activities are considered. We assume a 17.5 kg.y exposure. First-order and optimised topological cuts have been applied.

The SuperNEMO demonstrator, with 7 kg of material and 2.5 years of data acquisition, would achieve a $T_{1/2}^{0\nu} > 2.2 \times 10^{24}$ y sensitivity, one order of magnitude higher than the best limit ever reached. The corresponding limit on the effective neutrino mass is $\langle m_{\beta\beta} \rangle = [0.15 - 0.50]$ eV. This is a better result than for ^{82}Se sources, as the ^{150}Nd has a higher $G^{0\nu}$ factor.

- révéfier chiffers dans tableau car coupures lâches Pint et vertex ajoutées
- relire conclusion sur comparaison nb bdf attendu Se/Nd
- peut-être (sans doute) pour le ^{150}Nd , on pourrait alléger les spécifications sur le ^{214}Bi et le Radon. Pour cela il faudrait connaître la composition des bruits de fond dans le RoI optimisé pour le ^{150}Nd .

3.7 The final detector sensitivity

The ultimate goal of the SuperNEMO demonstrator is to show that the NEMO technology is scalable to probe unprecedented half-life on the $0\nu\beta\beta$ decay. The final detector would consist in building 20 modules similar to the demonstrator.

We suppose the specified activities of $\mathcal{A}^{\text{Tl}} = 2 \mu\text{Bq/kg}$, $\mathcal{A}^{\text{Bi}} = 10 \mu\text{Bq/kg}$ and $\mathcal{A}^{\text{Rn}} = 0.15 \text{ mBq/m}^3$ are reached. The simulations with an uniform magnetic field are used. With an exposure of 500 kg.y, the SuperNEMO final detector should reach a sensitivity $T_{1/2}^{0\nu} > 5.4 \times 10^{25}$ y, with ^{82}Se sources. The expected number of background events in the region of interest [2.75;3.1] MeV being high enough, optimised topological cut-offs were applied, with $P_{\text{int}} > 4 \%$ and $|\Delta Z| < 80 \text{ mm}$ (samely for $|\Delta Y|$). The corresponding sensitivity to the neutrino mass is $\langle m_{\beta\beta} \rangle = [0.079 - 0.15]$ eV.

3.8 Conclusion

Assuming the target background levels are reached, the SuperNEMO demonstrator, running for two and half years with 7 kg of ^{82}Se would be able to set a limit on the $0\nu\beta\beta$ process $T_{1/2}^{0\nu} > 5.6 \times 10^{24}$ years, translating into a

limit on the neutrino effective mass $\langle m_{\beta\beta} \rangle < [0.25 - 0.48]$ eV (depending on the Nuclear Matrix Elements)⁶. Latest measurements of source activities show that the specified background level for ^{208}Tl isotope is not reached, although it is improved in average by a factor of 2, compared to NEMO-3. To offset this, topological selections, designed to reject non-internal and non-simultaneous $2e$ events, have been optimised. The calorimeter time resolution plays an important role on the cut-off efficiencies, especially on the internal probability. Future studies would investigate the influence of this resolution on the final limit set on $T_{1/2}^{0\nu}$. Finally, assuming the worst activity is reached for the internal Bismuth isotope, we would set a limit on the $T_{1/2}^{0\nu} > 3.6 \times 10^{24}$ years, a decrease of 35 % compared to the specified activities result. This corresponds to $\langle m_{\beta\beta} \rangle < [0.31 - 0.59]$ eV. Finally, the limit on $T_{1/2}^{0\nu}$ could be enhanced by using a multivariate analysis, similarly to what is done in other double beta decay experiments, taking advantage of the several topological variables offered by SuperNEMO.

Recent studies have shown that the 25 Gauss magnetic field would be distorted by detector materials, especially the calorimeter magnetic shields. In this context, we studied the influence on the demonstrator sensitivity of the presence of magnetic field. Switching-off the field would decrease the sensitivity, but can be compensated by the topological cut-offs, useful with such a level of background.. Finally, the absence of magnetic field would slightly increase the limit set on the sensitivity to $T_{1/2}^{0\nu} > 3.7 \times 10^{24}$ years, taking into account the measured activities, and $\langle m_{\beta\beta} \rangle < [0.30 - 0.58]$ eV. Simulations with a mapped field shown that the signal and background selection efficiencies would be degraded by a non-uniform field. A more complete study would also take into account the possible impacts on the calorimeter energy resolution.

Like its predecessor, the SuperNEMO demonstrator was designed to study several isotopes, such as the ^{150}Nd . Assuming the target background activities are reached for ^{150}Nd sources, the SuperNEMO demonstrator would achieve a $T_{1/2}^{0\nu} > 2.2 \times 10^{24}$ years, and $\langle m_{\beta\beta} \rangle < [0.15 - 0.50]$ eV.

Finally, assuming we reach the target background levels, the SuperNEMO final detector would achieve an unprecedented limit of $T_{1/2}^{0\nu} > 5.4 \times 10^{25}$ years for ^{82}Se sources, and $T_{1/2}^{0\nu} > 2.4 \times 10^{25}$ years for ^{150}Nd sources. To complete these results, the SuperNEMO collaboration would study the influence on the sensitivity of external backgrounds, coming from detector materials as well as the laboratory.

- écrire commentaires tab bdf

⁶The real mass of isotope is 6.23 kg, then to achieve a 17.5 kg.y exposure, the demonstrator should run a little more than two years and a half.

Bibliography

- [1] M. Agostini et al. Probing majorana neutrinos with double- β decay. *Science* 365, 1445, 2019.
- [2] S.I. Alvis et al. Search for neutrinoless double-beta decay in ^{76}Ge with 26 kg-yr of exposure from the majorana demonstrator. *Phys. Rev. C*, 100, 2019.
- [3] O. Azzolini et al. First result on the neutrinoless double- β decay of ^{82}Se with cupid-0. *Phys. Rev. Lett.*, 120:232502, Jun 2018.
- [4] C. Alduino et al. First results from cuore: A search for lepton number violation via $0\nu\beta\beta$ decay of ^{130}Te . *Phys. Rev. Lett.*, 120:132501, Mar 2018.
- [5] J. B. Albert et al. Search for neutrinoless double-beta decay with the upgraded exo-200 detector. *Phys. Rev. Lett.*, 120:072701, Feb 2018.
- [6] A. Gando et al. Search for majorana neutrinos near the inverted mass hierarchy region with kamland-zen. *Phys. Rev. Lett.*, 117:082503, Aug 2016.
- [7] Nucleid database.
- [8] R. Arnold et al. Probing new physics models of neutrinoless double beta decay with supernemo. *Eur. Phys. J. C*, 2010.
- [9] S. Clavez. *Development of reconstruction tools and sensitivity of the SuperNEMO demonstrator*. PhD thesis, Université Paris Sud, 2017.
- [10] Garrido X. Bongrand M. Hamamatsu 8" pmt test in magnetic shield. Internal presentation, 2014.
- [11] Gomez-Cadenas et al. Physics case of supernemo with ^{82}Se source. Internal presentation, 2008.
- [12] R. Arnold et al. Final results on ^{82}Se double beta decay to the ground state of ^{82}Kr from the nemo-3 experiment. *Eur. Phys. J. C*, 2018.
- [13] Tretyak V.I. Ponkratenko O.A. and Zdesenko Yu.G. The event generator decay4 for simulation of doublebeta processes and decay of radioactive nuclei. *Phys. At. Nucl.*, 63:1282–1287, Jul 2000.

- [14] R. Arnold et al. Results of the search for neutrinoless double- β decay in ^{100}mo with the nemo-3 experiment. *Phys. Rev. D*, 2015.
- [15] Perrot F. Radiopurity measurements for 8" pmts and preliminary budget for the sn demonstrator. Internal presentation, 2017.
- [16] Cousins D. Feldman G. A unified approach to the classical statistical analysis of small signals. *Phys.Rev.*, pages 3873–3889, 1999.
- [17] J. Kotila and F. Iachello. Phase-space factors for double- β decay. *Phys. Rev. C*, 85:034316, Mar 2012.
- [18] J. Menéndez et al. Disassembling the nuclear matrix elements of the neutrinoless $\beta\beta$ decay. *Nuclear Physics A*, 818(3):139 – 151, 2009.
- [19] Y. Iwata et al. Large-scale shell-model analysis of the neutrinoless $\beta\beta$ decay of ^{48}Ca . *Phys. Rev. Lett.*, 116:112502, Mar 2016.
- [20] J. Barea, J. Kotila, and F. Iachello. $0\nu\beta\beta$ and $2\nu\beta\beta$ nuclear matrix elements in the interacting boson model with isospin restoration. *Phys. Rev. C*, 91:034304, Mar 2015.
- [21] J. Hyvärinen and J. Suhonen. Nuclear matrix elements for $0\nu\beta\beta$ decays with light or heavy majorana-neutrino exchange. *Phys. Rev. C*, 91:024613, Feb 2015.
- [22] F. Šimkovic et al. $0\nu\beta\beta$ and $2\nu\beta\beta$ nuclear matrix elements, quasiparticle random-phase approximation, and isospin symmetry restoration. *Phys. Rev. C*, 87:045501, Apr 2013.
- [23] Tomás R. Egidio J. Luis Vaquero López N., Rodríguez. Shape and pairing fluctuation effects on neutrinoless double beta decay nuclear matrix elements. *Phys. Rev. Lett.*, 111:142501, Sep 2013.
- [24] J. M. Yao, L. S. Song, K. Hagino, P. Ring, and J. Meng. Systematic study of nuclear matrix elements in neutrinoless double- β decay with a beyond-mean-field covariant density functional theory. *Phys. Rev. C*, 91:024316, Feb 2015.
- [25] P. K. Rath, R. Chandra, K. Chaturvedi, P. K. Raina, and J. G. Hirsch. Uncertainties in nuclear transition matrix elements for neutrinoless $\beta\beta$ decay within the projected-hartree-fock-bogoliubov model. *Phys. Rev. C*, 82:064310, Dec 2010.
- [26] Dong-Liang Fang, Amand Faessler, Vadim Rodin, and Fedor Šimkovic. Neutrinoless double- β decay of deformed nuclei within quasiparticle random-phase approximation with a realistic interaction. *Phys. Rev. C*, 83:034320, Mar 2011.
- [27] Loaiza P. Source foils measurement with bipo. Internal presentation, 2017.

- [28] Xin Ran Liu. Radon mitigation strategy and results for the supernemo experiment. IoP APP / HEPP Conference, 2018.
- [29] A. Chapon. *Mesure des processus de double désintégration bêta du Mo vers l'état excité 0_1^+ du Ru dans l'expérience Nemo3, Programme de R&D SuperNEMO : mise au point d'un détecteur BiPo pour la mesure de très faibles contaminations de feuilles sources*. PhD thesis, Université Caen Basse-Normandie, 2011.
- [30] Snow S. A magnetic field map for the tracker. Internal presentation, 2015.
- [31] A. Pin. *Recherche de la nature du neutrino via la décroissance double bêta sans émission de neutrinos. Caractérisation et optimisation du calorimètre SuperNEMO et impact sur la recherche de la décroissance du 82Se* . Développement du premier prototype LiquidO. PhD thesis, Université Bordeaux-Gradignan, 2020.
- [32] A. H. Wapstra G. Audi. The 1995 update to the atomic mass evaluation. *Nucl. Phys. A*, 595:409–480, feb 1995.
- [33] Dassié D. et al. Two-neutrino double- β decay measurement of ^{100}Mo . *Phys. Rev. D*, 51:2090–2100, Mar 1995.
- [34] R. Arnold et al. Measurement of the $2\nu\beta\beta$ decay half-life of ^{150}Nd and a search for $0\nu\beta\beta$ decay processes with the full exposure from the nemo-3 detector. *Phys. Rev. D*, 94, oct 2016.