

Integer Polynomials

Andres Buritica Monroy

1 Basic facts

For all of these, let $p(x) = \sum_{i=0}^n a_i x^i$ be a polynomial with integer coefficients.

We say that $p(x)$ is *divisible* by a polynomial $q(x)$ if there is a polynomial $r(x)$ (not necessarily with integer coefficients) such that $p(x) = q(x)r(x)$. We also write $q(x) \mid p(x)$.

- (Division algorithm) For any polynomial q there are unique polynomials f and r such that $p(x) = q(x)f(x) + r$ and $\deg R < \deg q$.
If q has integer coefficients, then f and r have rational coefficients. If q has a leading coefficient of ± 1 , then f and r have integer coefficients.
- (Factor theorem) If a is a root of p , then $x - a \mid p(x)$.
- (Remainder theorem) If $p(a) = c$ then $x - a \mid p(x) - c$.
- (*) If a and b are integers, then $a - b \mid p(a) - p(b)$.
- (Finite differences) If c is a constant and p is a polynomial with leading coefficient ax^n , then $p(x) - p(x - c)$ is a polynomial with leading coefficient nax^{n-1} .
- (Rational Root Theorem) If y and z are integers with $\gcd(y, z) = 1$ such that $p(y/z) = 0$ then $y \mid a_0$ and $z \mid a_n$.

2 Problems

1. Prove that every nonconstant integer polynomial has a composite number in its image.
2. Do there exist two quadratics $ax^2 + bx + c$ and $(a + 1)x^2 + (b + 1)x + (c + 1)$ which both have two integer roots?
3. Let P be an integer polynomial such that if $P(x)$ is an integer then x is rational. Prove that P is linear.
4. Find all polynomials $P(x)$ with integer coefficients such that if $m \mid n$ then $P(m) \mid P(n)$.
5. Prove that for any two distinct polynomials P and Q with coefficients in $\{0, 1, \dots, 9\}$, either $P(-2) \neq Q(-2)$ or $P(-5) \neq Q(-5)$.

6. Let $P(x)$ and $Q(x)$ be polynomials with integer coefficients such that the leading coefficient of $P(x)$ is 1. Suppose that $P(n)^n$ divides $Q(n)^{n+1}$ for infinitely many positive integers n . Prove that $P(n)$ divides $Q(n)$ for infinitely many positive integers n .
7. Let $n \geq 2$ be an integer and $P(x)$ be a polynomial with nonnegative integer coefficients satisfying $P(1) = 1$ and $x^n P(1/x) = P(x)$ for all x . Prove that there exist infinitely many pairs x, y of positive integers such that $x|P(y)$ and $y|P(x)$.
8. Given are positive integers a, b satisfying $a \geq 2b$. Does there exist a polynomial $P(x)$ of degree at least 1 with coefficients from the set $\{0, 1, 2, \dots, b-1\}$ such that $P(b) \mid P(a)$?

3 Homework

1. Prove that if P is a polynomial with integer coefficients and leading term $a_0 n^k$ such that $m \mid P(n)$ for all n , then $m \mid k!a_0$.
2. Prove that for every polynomial $P(x)$ of degree at least 2 with integer coefficients, there is an infinite arithmetic progression of integers which does not contain $P(k)$ for any integer k .
3. Is there a polynomial f of degree 2023 with integer coefficients such that

$$f(n), f(f(n)), f(f(f(n))), \dots$$

are pairwise relatively prime for any integer n ?