

Hensel's Lemma, quadratic residues

Andres Buritica Monroy

Once again, p is a prime and n is a positive integer throughout.

1 Hensel's Lemma

Let

$$P(x) = a_0x^0 + \cdots + a_nx^n$$

be a polynomial with integer coefficients. We define the *derivative*

$$P'(x) = a_1x^0 + 2a_2x^1 + 3a_3x^2 + \cdots + na_nx^{n-1}.$$

Let r be an integer such that $P(r) \equiv 0 \pmod{n}$ but $\gcd(P'(r), n) = 1$. Prove that for any positive integer m , there is a unique $s \pmod{n^m}$ such that $s \equiv r \pmod{n}$ and $P(s) \equiv 0 \pmod{n^m}$.

The case where $n = p$ is prime is most common. In this case, the condition $P(r) \equiv 0 \pmod{p}$ and $P'(r) \not\equiv 0 \pmod{p}$ is equivalent to r being a single root of $P \pmod{p}$. That is, $(x - r) \mid P(x)$ but $(x - r)^2 \nmid P(x)$.

2 Quadratic residues

If x can be written as y^2 for $y \in \mathbb{Z}_n^*$, then we say that x is a *quadratic residue (QR)* mod n . Note that 0 is not a QR mod n .

Prove that x is a QR mod n iff both

- $x \equiv 1 \pmod{\gcd(8, n)}$, and
- for each odd prime $p \mid n$, x is a QR mod p .

Hence, we now restrict ourselves to considering QRs mod p . Define the *Jacobi symbol*

$$\left(\frac{a}{p}\right) = \begin{cases} 0 & p \mid a \\ 1 & a \text{ is a QR mod } p \\ -1 & \text{otherwise} \end{cases}$$

- (Euler's criterion) Prove that $\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod{p}$. Hence, $\left(\frac{a}{p}\right) \left(\frac{b}{p}\right) = \left(\frac{ab}{p}\right)$.
- (Gauss' Lemma) Let $a \in \mathbb{Z}_p^*$, and let $S \subseteq \mathbb{Z}_p^*$ such that $x \in S \iff -x \notin S$. Let $T = \{ay : y \in S\}$. Then $\left(\frac{a}{p}\right) = (-1)^{|T \setminus S|}$.

- Find $\left(\frac{2}{p}\right)$ and $\left(\frac{-1}{p}\right)$.

Finally, there is quadratic reciprocity which we won't prove today. Let p and q be distinct odd primes. Then,

$$\left(\frac{q}{p}\right)\left(\frac{p}{q}\right) = (-1)^{\frac{(p-1)(q-1)}{4}}.$$

3 Problems

1. Let p be an odd prime and let a be an integer with $\gcd(a, p) = 1$. Prove that

$$\sum_{n=1}^p \left(\frac{n^2 + a}{p}\right) = -1.$$

2. Let k and n be positive integers such that n is odd. Prove that there is an integer a such that $a^{32} \equiv (n+1)^3 \pmod{n^k}$.
3. Prove that for any prime p and positive integer a with $p \nmid a$ there are at least $p-1$ solutions in \mathbb{Z}_p to $x^2 + y^2 \equiv a \pmod{p}$.
4. Let p be prime and let a and b be positive integers such that $p \nmid b$. Prove that there exists a positive integer n such that $p^a \mid n^n - b$.
5. If $p > 3$ is a prime such that $\varphi(p-1) > \frac{p-1}{3}$, prove that there are two consecutive generators mod p .
6. Let P be a nonconstant polynomial with integer coefficients. Prove that for any integer m there exist an integer n and a prime p such that $p^m \mid P(n)$.
7. Let a_1, a_2, a_3, \dots be a sequence of integers, such that for any positive integers n and k , the quantity

$$\frac{a_n + a_{n+1} + \dots + a_{n+k-1}}{k}$$

is always the square of an integer. Prove that all a_i s are equal.

8. Find all completely multiplicative functions $f : \mathbb{N} \rightarrow \mathbb{Z}$ such that for all $a, b \in \mathbb{N}$, at least two of $f(a)$, $f(b)$, $f(a+b)$ are equal.

4 Homework

1. Let $k = 2^{2^n} + 1$ for some positive integer n . Prove that k is prime if and only if $k \mid 3^{(k-1)/2} + 1$.
2. Find all positive integers k such that for all positive integers n , there exist a prime p and positive integers x and y for which $\gcd(x, y) = 1$ and $p^n \mid \frac{x^k - y^k}{x - y}$.
3. Let $p > 3$ be a prime and let a, b, c be integers with $a \neq 0$. Suppose that $ax^2 + bx + c$ is a perfect square for p consecutive integers x . Prove that $p \mid b^2 - 4ac$.