

Gastric Cancer

Lymphadenectomy

Retrospective data from Japan in the 1980's suggested superior survival after extended lymphadenectomy for gastric cancer.

Extent of lymphadenectomy can be categorized:


D1: Perigastric D2: Central nodes + splenic hilum D2 α : Central nodes D3: Extended nodes (peripancreatic, para-aortic, SMV=14V)

D1 Perigastric nodes

Lymph node stations immediately adjacent to the stomach

- 1: Right cardia
- 2: Left cardia
- 3: Lesser curvature
- 4: Greater curvature
- 5: Suprapyloric
- 6: Infrapyloric

D1 Perigastric Nodes



D2 Central Nodes + splenic hilum

Lymph nodes adjacent to celiac axis:

- 12a: Left side of porta hepatis
- 8: Common hepatic artery
- 7: Left gastric artery
- 9: Celiac axis
- 11: Proximal splenic artery
- 10: Splenic hilum

D1 α Central Nodes


Lymph nodes adjacent to celiac axis:

- 12a: Left side of porta hepatis
- 8: Common hepatic artery
- 7: Left gastric artery
- 9: Celiac axis
- 11: Proximal splenic artery
- ~~10: Splenic hilum~~


D2 Central Nodes

N2 Lymph nodes (branches coeliac axis)


- 7 Nodes along root left gastric artery
- 8 Nodes along common hepatic artery
- 9 Nodes around coeliac axis
- 10 Nodes at splenic hilum
- 11 Nodes along splenic artery



Durch Trial: D2 vs D1 Lymphadenectomy




Dutch Trial: Overall Survival



Number at risk

D1	380	231	174	149	132	108	47
D2	331	191	158	138	125	110	70


Dutch Trial: D2 vs D1 Lymphadenectomy




Number at risk

D1	380	231	174	149	132	108	47
D2	331	191	158	138	125	110	70


Durch Trial: Cause of Death



Durch Trial: Disease-free Survival



Dutch Trial: D2 vs D1 Lymphadenectomy



Dutch Trial: D2 vs D1

Operative mortality higher with D2 (10% vs 4%)

More complications with D2 (43% vs 25%)

More reoperations with D2 (18% vs 8%)

Dutch Trial: Total vs Subtotal gastrectomy

Protocol did not dictate extent of gastric resection, but did require a proximal margin of 5cm if a subtotal gastrectomy performed.

No difference in survival between total vs subtotal gastrectomy

Dutch Trial: Conclusions

D2 lymphadenectomy is associated with better local control of gastric cancer than D1 node dissection, but at an increased risk of mortality and complications.

Can the toxicity of extended lymphadenectomy be reduced?

- Elimination of splenectomy (D1 α)
- Minimally-invasive techniques

Splenectomy vs Splenic Preservation

505 patients with proximal gastric adenocarcinoma not invading the greater curvature treated with total gastrectomy randomized to with or without splenectomy. Splenectomy associated with higher morbidity and larger blood loss.

No difference in 5-year survival (75.1% vs 76.4%)

(sano277?)

Sentinel Node Biopsy in Early Gastric Cancer

Senorita trial Hur Lee Kim

MAGIC Trial - Perioperative Chemotherapy

503 gastric cancer stage II adenocarcinoma of stomach, GE junction or lower esophagus

ECF Chemo → Surgery → ECF Chemot vs Surgery alone

Chemotherapy: Epirubicin, cisplatin, 5FU

Surgery 3-6 weeks after last dose of chemo Chemo 6-12 weeks after surgery

MAGIC - Perioperative Chemotherapy

Tumor Location

- Gastric 74%
- GE junction 11%
- Distal esophagus 15%

MAGIC- Perioperative Chemotherapy

Curative radical resection 79% with chemo vs. 70% (p=0.03)

Longer 5-year survival with chemo (36% vs 23%). p=0.0009

Complete chemotherapy regimen (6 doses) in only 42%

Of patients who completed preop chemotherapy and surgery, only 34% received postoperative chemotherapy.

FLOT - Perioperative Chemotherapy

7616 patients with adenocarcinoma of GE junction or stomach randomized:

ECF → Surgery → ECF vs FLOT → Surgery →

Longer survival with FLOT (median 50 months vs 35 months)

TOPGEAR - Periop Chemo + Adjuvant ChemoRT

ECF → Surgery → ECF → ChemoRT

vs

ECF → Surgery → ECF

No difference in survival

GASTRICHIP - HIPEC

Patients with peritoneal disease from gastro cancer.

Chemo → Surgery with cytoreduction → Chemo

vs

Chemo → Surgery with cytoreduction + HIPEC → Chemo

(glehen1?)

GASTRICHIP

105 patients randomized 2014 - 2018. Trial closed due to slow accrual

55 patient treatment stopped prior to cytoreductive surgery due to disease progression

HIPEC with mitomycin and cisplatin for 60min at 42°C.

Median survival 15 months in both groups (without a difference).

(glehen1?)

PERISCOPE-II

Comparison of cytoreductive surgery + HIPEC to systemic chemotherapy in patients with gastric cancer and peritoneal metastasis.

(koemans1?)

CHIMERA Trial

FLOT + HIPEC vs FLOT + Surgery in advanced gastric cancer

PREVENT

Diffuse-type gastric and GE junction adenocarcinoma:

FLOT → Gastrectomy + HIPEC → FLOT vs

FLOT → Gastrectomy → FLOT

(gotze1?)

STOPGAP-II

Gastric and GE junction cancers with carcinomatosis

Chemo (3-6 month without progression) → laparoscopy with
PCI Staging. Intraop randomization:


Intraperitoneal Taxol

vs

Additional chemotherapy

Senthil

STOPGAP-II Schema



STOPGAP-II Schema

