Question	Answer	Marks	Partial Marks
3(c)(i)	*D* in correct position	**2**	**B1** for *D* 9.6 cm from *C* **B1** for *D* 100° from *C*
3(c)(ii)	64	**2**	**M1** for 96 ÷ 1.5 oe
3(c)(iii)	60	**2**	**M1** for 96 ÷ 8 × 5 oe
4(a)	Correct pattern drawn	**1**	
4(b)(i)	17 11	**2**	**B1** for 17 in correct position If 0 scored **SC1** for *their* 17 – 6 correct
4(b)(ii)	Subtract 6 oe	**1**	
4(c)(i)	$4n + 7$ oe final answer	**2**	**B1** for $4n + k$ or $jn + 7$, $j \neq 0$
4(c)(ii)	No, [because] 30.5 is not a whole number oe	**2**	**M1** for *their* **(c)(i)** = 129 or better
5(a)(i)	[0].66	**2**	**M1** for 1.24 × 3.5
5(a)(ii)	3500	**1**	
5(b)	11	**2**	**M1** for 10 ÷ [0].85 oe soi by 11.7[6]...
5(c)	609	**2**	**M1** for [0].87 × 700 oe
5(d)	21.4 or 21.42 to 21.43	**2**	**M1** for $\frac{18+5+23+40+28+19+17}{7}$ or 150 ÷ 7
5(e)	2 correct simultaneous equations e.g. $4a + 2p = 1.96$ and $7a + 3p = 3.24$	**B2**	Allow equations working in cents or dollars **B1** for one correct equation
	Correctly equating one set of coefficients	**M1**	**FT** *their* equations
	Correct method to eliminate one variable	**M1**	**FT** *their* equations Dependent on the coefficients being the same for one of the variables. Correct consistent use of addition or subtraction using *their* equations
	[apple =] 0.3[0]	**A1**	
	[plum =] 0.38	**A1**	If **M0M0** scored **SC1** for 2 correct answers given or **SC1** for 2 values satisfying one of *their* two original equations.
6(a)	Six hundred (and) two thousand (and) forty seven	**1**	
6(b)(i)	Any multiple of 14	**1**	

© UCLES 2018

Question	Answer	Marks	Partial Marks
6(b)(ii)	3136	**1**	
6(b)(iii)	47	**1**	
6(b)(iv)	1	**1**	
6(c)	156	**2**	**M1** for (12 =) 2, 2, 3 and (78 =) 2, 3, 13 or for 2, 2, 3, 13 or for list of multiples of 12 and 78 to at least 156 If 0 scored **SC1** for 156*k*
6(d)	6	**2**	**B1** for 2 or 3 as final answer
6(e)	$2^4 \times 3^3$ or $2 \times 2 \times 2 \times 2 \times 3 \times 3 \times 3$	**2**	**M1** for a complete factor tree with at most one error or 2, 2, 2, 2, 3, 3, 3 clearly identified as factors or **B1** for a correct product that equals 432
7(a)	−1.2, −1.5, 1.5	**2**	**B1** for 2 correct
7(b)	Correct curve	**4**	**B3FT** for 11 or 12 points correctly plotted or **B2FT** for 9 or 10 points correctly plotted or **B1FT** for 6, 7 or 8 points correctly plotted
7(c)	1.2 to 1.4	**1**	**FT** *their* **(b)**
7(d)(i)	Correct ruled line	**1**	
7(d)(ii)	(−2.6 to −2.3, −2.6 to −2.3,) and (2.3 to 2.6, 2.3 to 2.6)	**2**	**FT** $y = x$ drawn and *their* curve **B1FT** for one correct, or both *x* values correct or both *y* values correct
8(a)(i)	$\frac{4}{20}$ oe	**1**	
8(a)(ii)	0	**1**	
8(a)(iii)	$\frac{17}{20}$ oe	**1**	
8(b)(i)	9.75 or 9.746 to 9.747	**3**	**M2** for $\sqrt{12^2 - 7^2}$ soi or **M1** for $7^2 + BC^2 = 12^2$ soi
8(b)(ii)	33.5 or 33.49 or 33.492 to 33.494	**1**	**B1 FT** *their* **(b)(i)** × 2 + 14