Abbreviations

cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
nfww	not from wrong working
soi	seen or implied

Question	Answer	Marks	Partial Marks
1(a)(i)	5.84	**4**	**B1** for [0].76, 5.6[0], 5.6[0], 2.2[0] **M1** for *their* 4 costs added together **M1 dep** on **M1** for 20 – (*their* (0.76 + 2.2 + 5.6 + 5.6)) soi
1(a)(ii)(a)	$\frac{5}{100} \times 5.60$	**1**	
1(a)(ii)(b)	Correct explanation	**1**	
1(b)(i)	1.965	**3**	**M1** for figs 2 = *w* + figs(0.005 + 0.01 + 0.02) oe and **B1** for correct conversion of 2 kg = 2000 g or for 5 g, 10 g and 20 g or 35 g correctly converted to kg
1(b)(ii)	12.35 12.45	**2**	**B1** for each or **B1** for both correct but reversed
2(a)(i)	1	**1**	
2(a)(ii)	5	**1**	
2(a)(iii)	1.62 or 1.621…	**3**	**M1** for ([0 × 16] + 1 × 19 + 2 × 14 + 3 × 10 + 4 × 5 + 5 × 2) soi by 107 **M1dep** for *their* 107 ÷ 66
2(b)(i)	Eleven thousand six hundred (and) seventy-eight	**1**	
2(b)(ii)	11 700	**1**	
2(c)	71	**1**	
2(d)	1250	**2**	**M1** for $\frac{20}{0.016}$
3(a)(i)	Obtuse	**1**	
3(a)(ii)	134	**1**	
3(b)(i)	Octagon	**1**	

Question	Answer	Marks	Partial Marks
3(b)(ii)	165	**2**	**M1** for $\frac{360}{24}$ or $(24 - 2) \times 180$
3(c)(i)	132	**B2**	**B1** for angle $OBA = 24$ soi
	[Triangle AOB is] isosceles oe	**B1**	
3(c)(ii)	17 Angle [in a] semicircle is [90°]	**2**	**B1** for each
3(c)(iii)	Ruled tangent drawn at A	**1**	
4(a)(i)	55.8	**3**	**M2** for $5 \times 9 + 0.5 \times 2.4 \times 9$ oe or **M1** for 5×9 or $0.5 \times 2.4 \times 9$ oe Alternative method **M2** for $2 \times \frac{1}{2}(7.4 + 5) \times 4.5$ oe or **M1** for $\frac{1}{2}(7.4 + 5) \times 4.5$ oe
4(a)(ii)	9.09	**3**	**M2** for $0.9 \times 2.1 + 4 \times 1.5 \times 1.2$ oe or **M1** for 0.9×2.1 oe or [4 ×] 1.5×1.2 oe
4(a)(iii)	46.7 or 46.71	**1**	**FT** *their* **(a)(i)** – *their* **(a)(ii)**
4(b)	123.75 nfww	**4**	**M2** for $\frac{53}{4.5 \times 2.5}$ or **M1** for $\frac{53}{4.5}$ or 4.5×2.5 **B1FT** for 5 [tins]
5(a)(i)(a)	14	**1**	
5(a)(i)(b)	3.5[0]	**1**	
5(a)(i)(c)	[c=] $3.5d$	**1**	**FT** *their* **(a)(i)(b)**
5(a)(ii)(a)	$2 \times 4 + 5$	**1**	
5(a)(ii)(b)	[c=] $2d + 5$	**2**	**M1** for $2d + k$ or $md + 5$, $m \neq 0$
5(a)(ii)(c)	Correct ruled line	**2**	**M1** for ruled line with intercept (0, 5) or for ruled line with gradient 2 or for 2 correct points plotted or for *their* $c = 2d + 5$ correctly drawn
5(a)(ii)(d)	No, with correct reason	**1**	**FT** if two intersecting linear graphs
5(b)(i)	61	**1**	
5(b)(ii)	$5h + 8p = 80$	**2**	**M1** for $5h + 8p$