MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M Method marks, awarded for a valid method applied to the problem.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.

B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation 'dep' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
SC Special Case
nfww not from wrong working
soi seen or implied

Question	Answer	Marks	Partial Marks
1(a)(i)	7	**1**	
1(a)(ii)	49	**1**	
1(a)(iii)	7	**1**	
1(b)(i)	12	**1**	
1(b)(ii)	32	**1**	
1(b)(iii)	1	**1**	
1(b)(iv)	6	**1**	
1(c)	$3 \times 2 - (6 - 2) \div 2 = 4$	**1**	

© Cambridge University Press & Assessment 2024

Question	Answer	Marks	Partial Marks
1(d)	1020	**2**	**B1** for 1020k as final answer or **M1** for [30 =] 2 × 3 × 5 **and** [68 =] 2 × 2 × 17 or for [30 =] 2 × 15 **and** [68 =] 2 × 34 or 2 correct factor trees or correct tables or correct Venn diagram or a list of multiples of both 30 and 68 with at least their first 3 correct or 2 × 2 × 3 × 5 × 17 oe
2(a)(i)	2a final answer	**1**	
2(a)(ii)	$2x^2 - 7x$ final answer	**2**	**B1** for $2x^2$ or $-7x$ in final answer or for $2x^2 - 7x$ seen then spoilt.
2(b)	74	**2**	**B1** for 49 or 25
2(c)(i)	[m =] $\frac{T-20}{35}$ final answer	**2**	**M1** for $T - 20 = 35m$ or $\frac{T}{35} = m + \frac{20}{35}$
2(c)(ii)	1.8	**2**	**FT** *their* **(c)(i)** for 2 marks or 1 mark **M1** for (83 – 20) ÷ 35
2(d)	Correctly eliminates one variable	**M1**	Making the coefficients the same for one of the variables and correct consistent use of addition or subtraction using their equations alternative substitution method. M1 for correct rearrangement of one equation to make either x or y the subject and correct substitution of their rearrangement into 2nd equation.
	[x =] 3	**A1**	If A0 scored **SC1** for 2 values satisfying one of the original equations.
	[y =] −1.5	**A1**	
3(a)	Trapezium	**1**	
3(b)(i)	7.5	**1**	
3(b)(ii)	11 to 11.4	**1**	
3(c)(i)	Translation $\begin{pmatrix} 9 \\ -7 \end{pmatrix}$	**2**	**B1** for each
3(c)(ii)	Reflection $y = -3$ oe	**2**	**B1** for each

© Cambridge University Press & Assessment 2024