| Question | Answer | Marks | Partial Marks |
|---|---|---|---|
| 1 | rhombus | **1** | |
| 2 | triangular prism | **1** | |
| 3 | [$A$ =] 3.2<br>[$B$ =] 5.6 | **3** | **B2** for one correct or for both correct but reversed<br>or **M1** for $\frac{2.4}{7-4}\times k$ where $k$ = 1, 4, 7 or 11 oe<br>or for $\frac{x}{x+2.4}=\frac{4}{7}$ oe |
| 4 | 88 | **2** | **M1** for $\frac{1}{2}(10+12)\times 8$ oe |
| 5 | 250 | **1** | |
| 6 | 149 or 148.7 to 148.8… | **3** | **M1** for $\frac{1}{3}\times\pi\times 4^2\times 12$ oe<br>**M1** for 0.74 × *their* volume |
| 7 | $[m=]\frac{y-c}{x}$ or $\frac{y}{x}-\frac{c}{x}$<br>final answer | **2** | **M1** for $y - c = mx$ or $c - y = -mx$ or $\frac{y}{x}=m+\frac{c}{x}$ |
| 8 | 5.02 cao | **1** | |
| 9 | Correct elimination of one variable | **M1** | |
| | [$w$ =] 4<br>[$y$ =] –1 | **A2** | **A1** for one correct<br><br>If A0 scored, **SC1** for answers satisfying one of the original equations |
| 10 | 17.5[0] cao | **3** | **M1** for $12n + 9(n - 10) = 277.50$ oe<br><br>**M1** dep on *their* equation using $n$ and $n - 10$ for simplifying *their* equation correctly to $an = b$ |
| 11 | 27.8[0] | **2** | **M1** for $X\times\left(\frac{100-15}{100}\right)=23.63$ oe |
| 12 | 58 nfww | **2** | **M1** for 15.5 oe or 13.5 oe seen |
| 13(a) | 8 | **2** | **M1** for $\frac{9}{6}=\frac{12}{PR}$ oe |