Question	Answer	Marks	Partial Marks
27	[x =] 3, [y =] 7 [x =] 8, [y =] 22	**6**	**M2** for $x^2 - 11x + 24$ [= 0] oe simplified or **M1** for $x^2 - 8x + 22 = 3x - 2$ oe or better **M2** for correct method to solve *their* three-term quadratic e.g. $(x - 8)(x - 3)$ [= 0] or **M1** for $x(x - 3) - 8(x - 3)$ or $x(x - 8) - 3(x - 8)$ or $(x + a)(x + b)$ where $ab = 24$ or $a + b = -11$ **B1** for $x = 3$ and $x = 8$ or $y = 7$ and $y = 22$ or one correct pair If B0 scored and at least two method marks scored **SC1** for correct substitution of both of *their* x-values into $y + 2 = 3x$ or $y = x^2 - 8x + 22$
28	$\frac{2x+3}{x+7}$ final answer	**4**	**B2** for $(2x + 3)(x - 7)$ or **B1** for $2x(x - 7) + 3(x - 7)$ or $x(2x + 3) - 7(2x + 3)$ or $(2x + a)(x + b)$ where $ab = -21$ or $a + 2b = -11$ **B1** for $(x + 7)(x - 7)$
29	[R =] $\frac{x}{31}$	**5**	**M1** for $\pi (2R)^2$ oe **M1** for $2 \times \pi \times 2R \times 5R$ oe **M1** for $\frac{4\pi(2R)^2}{2}$ oe **M1** for $\pi \times R \times x + \pi \times R^2$ oe