Question	Answer	Marks	Partial Marks
19(a)	$11\sqrt{2}$ final answer	**2**	**B1** for $4\sqrt{2}$ or $7\sqrt{2}$
19(b)	$\sqrt{2}-1$ final answer	**2**	**M1** for $\times \dfrac{\sqrt{2}-1}{\sqrt{2}-1}$
20	$2\frac{2}{7}$ or $\frac{16}{7}$ oe	**3**	**M2** for $y = \dfrac{16}{\sqrt{x}}$ or for $y\sqrt{49} = 8\sqrt{4}$ oe or **M1** for $y = \dfrac{k}{\sqrt{x}}$
21(a)	$\dfrac{39}{x+1}$ or $\left(\dfrac{22}{x}+2\right)$	**1**	
21(b)	$\dfrac{39}{x+1} - \dfrac{22}{x} = 2$ oe OR $(x+1)\left(\dfrac{22}{x}+2\right) = 39$ oe	**M1**	
	$39x - 22(x+1) = 2x(x+1)$ oe OR $22 + 2x + \dfrac{22}{x} + 2 = 39$	**M1**	**FT** *their* **M1** if two fractions with *their* linear denominators May still be with common denominator **FT** *their* **M1** if product of 2 pairs of brackets = 39
	$2x^2 - 15x + 22 = 0$	**A1**	Brackets removed and terms collected with no errors or omissions
21(c)	$(2x-11)(x-2)$ [= 0]	**M2**	**M1** for $(2x+a)(x+b)$ [= 0] where $ab = 22$ or $a + 2b = -15$ or $x(2x-11) - 2(2x-11)$ [= 0] or $2x(x-2) - 11(x-2)$ [= 0]
	$\dfrac{11}{2}$ oe and 2	**A1**	
	[h =] 4 and [h =] 11	**A2**	**A1** for each

© Cambridge University Press & Assessment 2022

Question	Answer	Marks	Partial Marks
22	$\frac{16}{\sqrt{3}}$ or $\frac{16\sqrt{3}}{3}$	4	**B1** for $\cos 30 = \frac{\sqrt{3}}{2}$ **M2** for $\frac{8}{\cos 30}$ or **M1** for $\frac{8}{x} = \cos 30$
23	$\frac{25 - x}{(x - 4)(x + 3)}$ oe final answer	3	**M1** for denominator $(x – 4)(x + 3)$ or $x^2 – x – 12$ **B1** for $3(x + 3) – 4(x – 4)$ seen
24(a)	$(x – 2)^2 + 3$	2	**M1** for $(x – 2)^2$ seen
24(b)	(2, 3)	1	**FT** *their* **(a)**
25	$\frac{15}{2}$ oe	3	**M2** for $\frac{9}{x} = \frac{6}{5}$ oe or $x^2 = \frac{25}{36} \times 9^2$ oe or better or **M1** for $\sqrt{\frac{36}{25}}$ or $\sqrt{\frac{25}{36}}$ oe or better or for $\left(\frac{9}{x}\right)^2 = \frac{36}{25}$ or $\left(\frac{x}{9}\right)^2 = \frac{25}{36}$ oe or better
26(a)	Correct sketch and –2, 0 and 3 indicated on x-axis with graph going beyond intersections with x-axis  	3	**B2** for correct sketch without labels or **B1** for any positive cubic with two turning points
26(b)	$x^3 – x^2 – 6x$	3	**B2** for $x(x^2 + 2x – 3x – 6)$ or $(x^2 + 2x)(x – 3)$ or $(x + 2)(x^2 – 3x)$ or **B1** for $x^2 + 2x – 3x – 6$ or $x^2 + 2x$ or $x^2 – 3x$

© Cambridge University Press & Assessment 2022