

# STA623 - Bayesian Data Analysis - Practical 3 (Solutions)

Marc Henrion

7 September 2022

## Practical 3

### Notation

- $X, Y, Z$  - random variables
- $x, y, z$  - measured / observed values
- $\bar{X}, \bar{Y}, \bar{Z}$  - sample mean estimators for  $X, Y, Z$
- $\bar{x}, \bar{y}, \bar{z}$  - sample mean estimates of  $X, Y, Z$
- $\hat{T}, \hat{t}$  - given a statistic  $T$ , estimator and estimate of  $T$
- $P(A)$  - probability of an event  $A$  occurring
- $f_X(\cdot), f_Y(\cdot), f_Z(\cdot)$  - probability mass / density functions of  $X, Y, Z$ ; sometimes  $p_X(\cdot)$  etc. rather than  $f_X(\cdot)$
- $p(\cdot)$  - used as a shorthand notation for pmfs / pdfs if the use of this is unambiguous (i.e. it is clear which is the random variable)
- $X \sim F$  -  $X$  distributed according to distribution function  $F$
- $E[X], E[Y], E[Z], E[T]$  - the expectation of  $X, Y, Z, T$  respectively

### Exercise 1

Show that the Bayes estimator  $\hat{\theta}_B$  for the quadratic loss function  $\mathcal{C}(\theta - \hat{\theta}) = (\theta - \hat{\theta})^2$  is given by the posterior mean. In other words, show that:

$$E[\theta|y] = \arg \min_{\hat{\theta}} \int_{\mathcal{Y}} \int_{\Theta} \mathcal{C}(\theta - \hat{\theta}) p(\theta, y) d\theta dy$$

### Exercise 1 (Solution)

We saw in lectures that we can use the multiplication rule and then optimise the inner integral only:

$$\hat{\theta}_B = \arg \min_{\hat{\theta}} \int \mathcal{C}(\theta - \hat{\theta}) p(\theta|y) d\theta \tag{1}$$

$$= \arg \min_{\hat{\theta}} \int (\theta - \hat{\theta})^2 p(\theta|y) d\theta \tag{2}$$

To find the minimum, we solve

$$\frac{d}{d\hat{\theta}} \int (\theta - \hat{\theta})^2 p(\theta|y) d\theta = 0 \quad (3)$$

$$\iff \int 2(\theta - \hat{\theta})(-1)p(\theta|y) d\theta = 0 \quad (4)$$

$$\iff \int (\theta - \hat{\theta})p(\theta|y) d\theta = 0 \quad (5)$$

$$\iff \int \theta p(\theta|y) d\theta = \int \hat{\theta} p(\theta|y) d\theta \quad (6)$$

Note that  $\int \hat{\theta} p(\theta|y) d\theta = \hat{\theta} \int p(\theta|y) d\theta = \hat{\theta}$  since the posterior distribution for  $\theta$  is a probability distribution and needs to integrate to 1.

Therefore we see that

$$\hat{\theta}_B = \int \theta p(\theta|y) d\theta$$

This is the posterior mean  $E[\theta|y] = \int \theta p(\theta|y) d\theta$ .

## Exercise 2

Suppose  $\pi \sim \text{Beta}(2, 3)$  and  $Y_1, \dots, Y_n \sim_{\text{iid}} \text{Bernoulli}(\pi)$ . Further suppose we observe data  $y_1, \dots, y_n$  with  $n = 25, k = \sum_i y_i = 16$ .

Find the following:

- posterior distribution  $p(\pi|k)$  and plot it, comparing it to the prior distribution
- posterior predictive distribution  $p(\tilde{y}|y_1, \dots, y_n)$
- the 95% quantile-based Bayesian confidence interval for  $\pi$
- the 95% HPD interval

Further, compute:

- $P(\pi > 0.5|k)$
- For the following 2 hypotheses:  $H_1 : \pi \in [0.3, 0.5], H_2 : \pi \in [0.5, 0.7]$ , compute the prior and posterior odds and calculate the Bayes factor.

## Exercise 2 (Solution)

### Posterior distribution

Recall:

$$\begin{cases} \text{prior } \Pi & \sim \text{Beta}(a, b) \\ \text{likelihood } Y|\Pi & \sim \text{Bin}(n, \pi) \end{cases}$$

$$\Rightarrow \text{posterior } \Pi|Y = k \sim \text{Beta}(a + k, b + n - k)$$

Here:

- $a = 2$
- $b = 3$
- $n = 25$
- $k = 16$

Hence, we have a  $\text{Beta}(2+16, 3+9) = \text{Beta}(18, 12)$  distribution.


Alternatively, you can also see that

$$\begin{aligned} p(\pi|k = 16) &\propto \pi^{2-1}(1-\pi)^{3-1}\pi^{16}(1-\pi)^{25-16} = \pi^{17}(1-\pi)^{11} \\ &\Rightarrow \Pi|Y = k \sim \text{Beta}(18, 12) \end{aligned}$$

```
xx<-seq(0,1,length=100)
yyPr<-dbeta(xx,2,3)
yyPo<-dbeta(xx,18,12)

plot(type="l",xx,yyPo,lwd=3,col="mediumorchid",
      xlab=expression(pi),ylab="density",cex.lab=2,cex.axis=2)
lines(xx,yyPr,lwd=3,lty=2,col="steelblue")

legend(x="topleft",bty="n",lwd=2,col=c("steelblue","mediumorchid"),
       lty=c(2,1),legend=c("prior","posterior"),cex=2)
```



### Posterior predictive distribution

$$p(\tilde{Y} = 1|y_1, \dots, y_n) = \int p(\tilde{y}, \pi|y_1, \dots, y_n) d\pi \quad (7)$$

$$= \int p(\tilde{y}|\pi) p(\pi|y_1, \dots, y_n) d\pi \quad (8)$$

$$= \int \pi \frac{\Gamma(30)}{\Gamma(18)\Gamma(12)} \pi^{17} (1-\pi)^{11} d\pi \quad (9)$$

$$= \frac{\Gamma(30)\Gamma(19)}{\Gamma(18)\Gamma(31)} \int \frac{\Gamma(31)}{\Gamma(19)\Gamma(12)} \pi^{18} (1-\pi)^{11} d\pi \quad (10)$$

$$= \frac{\Gamma(30)\Gamma(19)}{\Gamma(18)\Gamma(31)} = \frac{18}{30} = 3/5 = 0.6 \quad (11)$$

From this it follows that  $p(\tilde{Y} = 0|y_1, \dots, y_n) = 1 - \frac{3}{5} = \frac{2}{5} = 0.4$ .

### Credible intervals

We have:

- $q_{0.025; \text{Beta}(18,12)} = 0.4226$
- $q_{0.975; \text{Beta}(18,12)} = 0.7648$

Therefore the 95% quantile based Bayesian confidence interval for  $\pi$  is given by  $[0.42, 0.76]$

```
qbeta(c(0.025,0.975),18,12)
```

```
## [1] 0.4226046 0.7647598
```

For the HPD we are going to use the function `hdi()` from the R package `HDInterval`:

```
library(HDInterval)
hdp<-hdi(qbeta,0.95,shape1=18,shape2=12)
print(hdp)
```

```
##      lower      upper
## 0.4273464 0.7690367
## attr("credMass")
## [1] 0.95
```

From this we find that the 95% HDP interval for  $\pi$  is given by  $[0.43, 0.77]$

We have

$$P(\pi > 0.5 | Y = k) = \int_{0.5}^1 p(\pi | k) d\pi \quad (12)$$

$$= \int_{0.5}^1 \frac{\Gamma(30)}{\Gamma(18)\Gamma(12)} \pi^{17} (1 - \pi)^{11} d\pi \quad (13)$$

$$= \approx 0.87 \quad (14)$$

```
integrate(dbeta, lower=0.5, upper=1, shape1=18, shape2=12)
```

```
## 0.8675346 with absolute error < 2.8e-09
```

Alternatively we could have worked with the posterior cdf:

```
1-pbeta(0.5, 18, 12)
```

```
## [1] 0.8675346
```

**Bayes factor**

Prior odds

$$\frac{P(H_1)}{P(H_2)} = \frac{\int_{0.3}^{0.5} p_{\beta(3,2)}(\pi) d\pi}{\int_{0.5}^{0.7} p_{\beta(3,2)}(\pi) d\pi} = 1.48$$

Prosterior odds

$$\frac{P(H_1 | k)}{P(H_2 | k)} = \frac{\int_{0.3}^{0.5} p_{\beta(18,12)}(\pi) d\pi}{\int_{0.5}^{0.7} p_{\beta(18,12)}(\pi) d\pi} = 0.18$$

```
priorOdds<-
  integrate(dbeta, 0.3, 0.5, shape1=2, shape2=3)$value/integrate(dbeta, 0.5, 0.7, shape1=2, shape2=3)$value
#priorOdds<-
# (pbeta(0.5, 2, 3)-pbeta(0.3, 2, 3))/(pbeta(0.7, 2, 3)-pbeta(0.5, 2, 3)) # alternative way of doing this
priorOdds
```

```
## [1] 1.482517
```

```
posteriorOdds<-
  integrate(dbeta, 0.3, 0.5, shape1=18, shape2=12)$value/integrate(dbeta, 0.5, 0.7, shape1=18, shape2=12)$value
posteriorOdds
```

```
## [1] 0.1789838
```

Bayes Factor

$$BF = \frac{\text{posterior odds}}{\text{prior odds}} = 0.18/1.48 = 0.12$$

```
bayesFactor<-posteriorOdds/priorOdds  
bayesFactor
```

```
## [1] 0.1207296
```

## Exercise 3

Suppose  $\lambda \sim \text{Gamma}(5, 2)$  and  $Y_1, \dots, Y_n \sim_{\text{iid}} \text{Poisson}(\lambda)$ . Further suppose we observe data  $y_1, \dots, y_n$  with  $n = 18, k = \sum_i y_i = 40$ .

Find the following:

- posterior distribution  $p(\lambda|y_1, \dots, y_n)$  and plot it, comparing it to the prior distribution
- posterior predictive distribution  $p(\tilde{y}|y_1, \dots, y_n)$
- the 95% quantile-based Bayesian confidence interval for  $\lambda$
- the 95% HPD interval

Further, compute:

- $P(\lambda \leq 1|y_1, \dots, y_n)$
- For the following 2 hypotheses:  $H_1 : \lambda \in [0.75, 1.25], H_2 : \lambda \in [1.75, 2.25]$ , compute the prior and posterior odds and calculate the Bayes factor.

## Exercise 3 (Solution)

Posterior distribution

$$p(\lambda|y_1, \dots, y_n) \propto \lambda^{5-1} e^{-2\lambda} \lambda^{\sum_i y_i} e^{-n\lambda} \quad (15)$$


$$\propto \lambda^{5+40-1} e^{-(2+18)\lambda} \quad (16)$$

$$\Rightarrow \Lambda|y_1, \dots, y_n \sim \Gamma(45, 20) \quad (17)$$

```
xx<-seq(0,8,length=100)
yyPr<-dgamma(xx,shape=5,rate=2)
yyPo<-dgamma(xx,shape=45,rate=20)

plot(type="l",xx,yyPo,lwd=3,col="mediumorchid",
      xlab=expression(pi),ylab="density",cex.lab=2,cex.axis=2)
lines(xx,yyPr,lwd=3,lty=2,col="steelblue")

legend(x="topleft",bty="n",lwd=2,col=c("steelblue","mediumorchid"),
       lty=c(2,1),legend=c("prior","posterior"),cex=2)
```



### Posterior predictive distribution

From Practical 1&2, Exercise 4, we know:

$$\tilde{Y}|y_1, \dots, y_n \sim \text{NegBin}(45, 20/21)$$

### Credible intervals

We have:

- $q_{0.025;\Gamma(45,20)} = 1.6412$
- $q_{0.975;\Gamma(45,20)} = 2.9534$

Therefore the 95% quantile based Bayesian confidence interval for  $\pi$  is given by  $[1.64, 2.95]$

```
qgamma(c(0.025,0.975),shape=45,rate=20)
```

```
## [1] 1.641165 2.953397
```

For the HPD we are going to use the function `hdi()` from the R package `HDInterval`:

```
library(HDInterval)
hdp<-hdi(qgamma,0.95,shape=45,rate=20)
print(hdp)
```

```
##   lower   upper
## 1.611342 2.917067
## attr(,"credMass")
## [1] 0.95
```

From this we find that the 95% HDP interval for  $\lambda$  is given by  $[1.61, 2.92]$

We have



$$P(\lambda \leq 1|y_1, \dots, y_n) = \int_0^1 p(\lambda|y_1, \dots, y_n) d\lambda \quad (18)$$

$$= \int_0^1 \frac{20^{45}}{\Gamma(45)} \lambda^{44} e^{-20\lambda} d\lambda \quad (19)$$

$$= 1.06 \cdot 10^{-6} \quad (20)$$

```
integrate(dgamma,lower=0,upper=1,shape=45,rate=20)
```

```
## 1.060263e-06 with absolute error < 4.4e-13
```

**Bayes factor**

Prior odds

$$\frac{P(H_1)}{P(H_2)} = \frac{\int_{0.75}^{1.25} p_{\gamma(5,2)}(\lambda) d\lambda}{\int_{1.75}^{2.25} p_{\Gamma(5,2)}(\lambda) d\lambda} = 0.2513$$

Prosterior odds

$$\frac{P(H_1|y_1, \dots, y_n)}{P(H_2|y_1, \dots, y_n)} = \frac{\int_{0.75}^{1.25} p_{\gamma(45,20)}(\lambda) d\lambda}{\int_{1.75}^{2.25} p_{\Gamma(45,20)}(\lambda) d\lambda} = 0.0004$$

```
priorOdds<-
```

```
  integrate(dgamma,0.75,1.25,shape=5,rate=2)$value/integrate(dgamma,1.25,2.25,shape=5,rate=2)$value
priorOdds
```

```
## [1] 0.2513296
```

```
posteriorOdds<-
```

```
  integrate(dgamma,0.75,1.25,shape=45,rate=20)$value/integrate(dgamma,1.25,2.25,shape=45,rate=20)$value
posteriorOdds
```

```
## [1] 0.0003851655
```

Bayes Factor

$$BF = \frac{\text{posterior odds}}{\text{prior odds}} = 0.004/0.2513 = 0.0015$$

```
bayesFactor<-posteriorOdds/priorOdds
bayesFactor
```

```
## [1] 0.001532511
```

[end of STA623 BDA Practical 3]