

# STA623 - Bayesian Data Analysis - Practical 5

Marc Henrion

2024-11-01

## Practical 5

### Notation

- $X, Y, Z$  - random variables
- $x, y, z$  - measured / observed values
- $\bar{X}, \bar{Y}, \bar{Z}$  - sample mean estimators for  $X, Y, Z$
- $\bar{x}, \bar{y}, \bar{z}$  - sample mean estimates of  $X, Y, Z$
- $\hat{T}, \hat{t}$  - given a statistic  $T$ , estimator and estimate of  $T$
- $P(A)$  - probability of an event  $A$  occurring
- $f_X(\cdot), f_Y(\cdot), f_Z(\cdot)$  - probability mass / density functions of  $X, Y, Z$ ; sometimes  $p_X(\cdot)$  etc. rather than  $f_X(\cdot)$
- $p(\cdot)$  - used as a shorthand notation for pmfs / pdfs if the use of this is unambiguous (i.e. it is clear which is the random variable)
- $X \sim F$  -  $X$  distributed according to distribution function  $F$
- $E[X], E[Y], E[Z], E[T]$  - the expectation of  $X, Y, Z, T$  respectively

## Exercise 1

Fit the model from Practical 3, Exercise 3 using JAGS and the `rjags` package. Use this as the data from the sampling model:

$$y = (1, 3, 2, 3, 0, 2, 6, 4, 4, 1, 1, 3, 2, 3, 1, 1, 3, 0)$$

Inspect the trace plot and plot the posterior distribution.

Compute the posterior mean and the quantile-based 95% Bayesian confidence interval.

## Exercise 2

Generate the following data

```
N<-100
x<-rnorm(N)
z<-2-4*x
p<-1/(1+exp(-z))
y<-rbinom(n=N,size=1,prob=p)

dat<-list(N=N,x=x,y=y)
```

Use R and JAGS to fit a Bayesian logistic regression model to these data:

$$g(E[Y|X]) = \beta_0 + \beta_1 X$$

where  $g(\pi) = \log(\pi/(1 - \pi))$ .

Compute the Gelman-Rubin convergence statistic and inspect trace plots and autocorrelations for the samples from the posterior distributions.

Compute the posterior mean, median, a 95% quantile-based confidence interval and a 95% highest posterior density confidence interval.

Compute the effective sample sizes for  $\beta_0, \beta_1$ .

[end of STA623 BDA Practical 5]