

Solutions to some exercises from Walter Rudin's
Functional Analysis

gitcordier

July 24, 2020

Contents

1	Topological Vector Spaces	1
1.1	Exercise 7. Metrizability & number theory	2

Chapter 1

Topological Vector Spaces

1.1 Exercise 7. Metrizable & number theory

Let be X the vector space of all complex functions on the unit interval $[0, 1]$, topologized by the family of seminorms

$$p_x(f) = |f(x)| \quad (0 \leq x \leq 1).$$

This topology is called the topology of pointwise convergence. Justify this terminology. Show that there is a sequence $\{f_n\}$ in X such that (a) $\{f_n\}$ converges to 0 as $n \rightarrow \infty$, but (b) if $\{\gamma_n\}$ is any sequence of scalars such that $\gamma_n \rightarrow \infty$ then $\{\gamma_n f_n\}$ does not converge to 0. (Use the fact that the collection of all complex sequences converging to 0 has the same cardinality as $[0, 1]$.) This shows that metrizable cannot be omitted in (b) of Theorem 1.28.

Proof. Our justification consists in proving that τ -convergence and pointwise convergence are the same one. To do so, remark first that the family of the seminorms p_x is separating. By [1.37], the collection \mathcal{B} of all finite intersections of the sets

$$V^{(x,k)} \triangleq \{p_x < 2^{-k}\} \quad (x \in [0, 1], k \in \mathbf{N}) \quad (1.1)$$

is then a local base for a topology τ on X . Given $\{f_n : n = 1, 2, 3, \dots\}$, we set

$$\text{off}(U) \triangleq \sum_{n=1}^{\infty} [f_n \notin U] \quad (U \in \tau), \quad (1.2)$$

with the convention $\text{off}(U) = \infty$ whether the sum has no finite support. So,

$$\sum_{i=1}^m \text{off}(U^{(i)}) = \sum_{n=1}^{\infty} \sum_{i=1}^m [f_n \notin U^{(i)}] \geq \text{off}(U^{(1)} \cap \dots \cap U^{(m)}) \quad (1.3)$$

We first assume that $\{f_n\}$ τ -converges to some f in X , *i.e.*

$$\text{off}(f + V) < \infty \quad (V \in \mathcal{B}). \quad (1.4)$$

The special cases $V = V^{(x,k)}$ mean the pointwise convergence of $\{f_n\}$. Conversely, assume that $\{f_n\}$ does not τ -converges to any g in X , *i.e.*

$$\forall g \in X, \exists V^{(g)} \in \mathcal{B} : \text{off}(g + V^{(g)}) = \infty. \quad (1.5)$$

Given g , $V^{(g)}$ is then an intersection $V^{(x^{(1)}, k^{(1)})} \cap \dots \cap V^{(x^{(m)}, k^{(m)})}$. Thus

$$\sum_{i=1}^m \text{off}(g + V^{(x^{(i)}, k^{(i)})}) \stackrel{(1.3)}{\geq} \text{off}(g + V^{(g)}) \stackrel{(1.5)}{=} \infty. \quad (1.6)$$

One of the sum $\text{off}(g + V^{(x^{(i)}, k^{(i)})})$ must then be ∞ . This implies that convergence of f_n to g fails at point x_i . g being arbitrary, we so conclude that f_n does not converge pointwise. We have just proved that τ -convergence is a rewording of pointwise convergence. We now aim to prove the second part. From now on, k , n and p run on \mathbf{N}_+ . Let $\text{dyadic}(x)$ be the usual dyadic expansion of a real number x , so that $\text{dyadic}(x)$ is an aperiodic binary sequence **iff** x is irrational. Define

$$f_n(x) \triangleq \begin{cases} 2^{-\sum_{k=1}^n \text{dyadic}(x)_k} & (x \in [0, 1] \setminus \mathbf{Q}) \\ 0 & (x \in [0, 1] \cap \mathbf{Q}) \end{cases} \quad (1.7)$$

so that $f_n(x) \xrightarrow{n \rightarrow \infty} 0$ and take scalars γ_n such that $\xrightarrow{n \rightarrow \infty} \infty$, *i.e.* at fixed p , γ_n is greater than 2^p for almost all n . Next, choose $n^{(p)}$ among those *almost all* n that are large enough to satisfy

$$n^{(p-1)} - n^{(p-2)} < n^{(p)} - n^{(p-1)} \quad (1.8)$$

(start with $n^{(-1)} = n^{(0)} = 0$) and so obtain

$$2^p < \gamma_{n^{(p)}} : 0 < n^{(p)} - n^{(p-1)} \xrightarrow{p \rightarrow \infty} \infty. \quad (1.9)$$

The indicator χ of $\{n^{(1)}, n^{(2)}, \dots\}$ is then aperiodic, *i.e.*

$$x^{(\gamma)} \triangleq \sum_{k=1}^{\infty} \chi_k 2^{-k} \quad (1.10)$$

is irrational. Consequently,

$$\text{dyadic}(x^{(\gamma)})_{-k} = \chi_k. \quad (1.11)$$

We now easily see that

$$\chi_1 + \dots + \chi_{n^{(p)}} = p, \quad (1.12)$$

which, combined with (1.7), yields

$$f_{n^{(p)}}(x^{(\gamma)}) = 2^{-p}. \quad (1.13)$$

Finally,

$$\gamma_{n^{(p)}} f_{n^{(p)}}(x^{(\gamma)}) > 1. \quad (1.14)$$

We have so established that the subsequence $\{\gamma_{n^{(p)}} f_{n^{(p)}}\}$ does not tend pointwise to 0, hence neither does the whole sequence $\{\gamma_n f_n\}$. In other words, (b) holds, which is in violent contrast with [1.28]: X is then not metrizable. So ends the proof. \square