

# PASHUSETU: VETERINARY TRIAGE SUMMARY

*AI-Assisted Bovine Disease Assessment & Triage*

|                         |                       |                         |         |
|-------------------------|-----------------------|-------------------------|---------|
| <b>Owner / Farmer:</b>  | Ramesh Kumar          | <b>Cattle Tag / ID:</b> | #1025   |
| <b>Assessment Date:</b> | 17 Aug 2026, 01:22 PM | <b>Language:</b>        | English |



## Clinical Assessment & First-Aid Guidelines

### [Obs] 1. Verification & Clinical Observations

- Subject Confirmation: The image clearly depicts a bovine animal, likely a Zebu breed (indicated by the hump and dewlap), standing in a dry, dusty environment.
- Physical Lesions: The most striking feature is the presence of numerous, discrete, raised nodules scattered across the animal's skin. These lesions are visible on the neck, shoulder, flank, and head.
- Lesion Characteristics: The nodules appear to be firm and vary in size. Some lesions show dark, necrotic centers or scabbing, indicating tissue death or healing. The skin around the lesions appears thickened.
- General Condition: The animal appears somewhat emaciated (thin), suggesting a chronic condition or poor nutritional status alongside the illness.

### [Dx] 2. Suspected Condition & Confidence

- Disease Name: Lumpy Skin Disease (LSD)
- Confidence Level: High

### [Markers] 3. Diagnostic Markers

- Nodular Dermatitis: The hallmark sign is the presence of multiple, firm, raised nodules (papules) on the skin.
- Distribution: The lesions are widespread, affecting the head, neck, and body.
- Necrosis/Scabbing: Several nodules appear to have central necrosis (black spots) or are forming scabs, which is typical as LSD lesions progress.
- Emaciation: The visible ribs and thin frame suggest the animal has been suffering for some time or has a concurrent

metabolic issue.

#### [Care] 4. Immediate First-Aid & Farm Management

- Isolation: Immediately quarantine the affected animal from the rest of the herd. LSD is highly contagious and spread by vectors (flies, mosquitoes, ticks).
- Wound Care: Gently clean the open lesions or scabs with an antiseptic solution (like Povidone-Iodine) to prevent secondary bacterial infections.
- Supportive Treatment:
  - Administer broad-spectrum antibiotics to prevent secondary infections.
  - Provide anti-inflammatory drugs (NSAIDs) to reduce fever and pain.
  - Ensure adequate hydration and nutrition (high-quality fodder) to help the animal recover from emaciation.
- Vector Control: Apply insecticides to the animal and the surrounding environment to reduce the population of biting insects that transmit the virus.

---

Disclaimer: AI triage assistant only. Always consult a licensed veterinarian for definitive treatment.

*Disclaimer: PashuSetu AI provides preliminary automated triage support only. Always consult a licensed veterinary doctor for laboratory confirmation, prescription antibiotics, and definitive therapy.*