

PASHUSETU: VETERINARY TRIAGE SUMMARY

AI-Assisted Bovine Disease Assessment & Triage

Owner / Farmer:	Ramesh Kumar	Cattle Tag / ID:	#1254
Assessment Date:	17 Aug 2026, 01:30 PM	Language:	English



Clinical Assessment & First-Aid Guidelines

[Obs] 1. Verification & Clinical Observations

- Confirmation: The image clearly depicts a bovine animal, specifically a Zebu-type cattle (likely a crossbreed or indigenous breed common in South Asia/Africa) characterized by a hump and dewlap.
- Physical Lesions: The most striking observation is the presence of numerous, discrete, raised nodules and papules scattered across the entire body surface. These lesions are visible on the neck, flank, legs, and face.
- Lesion Characteristics: Many of the nodules appear dark, suggesting necrosis or scabbing. Some look like firm, round bumps, while others appear to be healing scabs. The distribution is generalized rather than localized to one area.
- Other Observations: The animal appears to be standing on dry, dusty ground. There is no obvious discharge from the eyes or nose visible in these specific frames, though the muzzle area is affected by the skin lesions.

[Dx] 2. Suspected Condition & Confidence

- Disease Name: Lumpy Skin Disease (LSD)
- Confidence Level: High

[Markers] 3. Diagnostic Markers

- Generalized Nodules: The presence of firm, discrete skin nodules (papules) that vary in size is the hallmark sign of LSD.
- Necrotic Centers: The dark appearance of many spots indicates the center of the nodules may be necrotic (dying tissue), which is typical as the disease progresses.
- Widespread Distribution: The lesions are not confined to one area but cover the neck, body, and limbs, which is

characteristic of the viral spread in LSD.

- Breed Susceptibility: The animal appears to be a Zebu breed, which is highly susceptible to this poxvirus.

[Care] 4. Immediate First-Aid & Farm Management

- Quarantine: Immediately isolate the affected animal from the rest of the herd. LSD is highly contagious and spreads via vectors like mosquitoes, flies, and ticks.
- Hygiene & Disinfection: Disinfect the area where the animal is kept. Use standard farm disinfectants to kill potential vectors.
- Topical Care: Apply antiseptic sprays or iodine solutions to the open lesions to prevent secondary bacterial infections. Keep the wounds clean and dry.
- Fly Control: Implement strict fly control measures (spraying insecticides on the animal and surroundings) to prevent flies from landing on the open sores, which can cause myiasis (fly strike) and spread the virus further.
- Supportive Care: Ensure the animal has access to clean water and high-quality feed to support its immune system. Monitor for fever and appetite loss.

Disclaimer: AI triage assistant only. Always consult a licensed veterinarian for definitive treatment.

Disclaimer: PashuSetu AI provides preliminary automated triage support only. Always consult a licensed veterinary doctor for laboratory confirmation, prescription antibiotics, and definitive therapy.