

STA 250/MTH 342 – Intro to Mathematical Statistics

Lecture 14

A brief recap

- ▶ Inference using Bayes' Theorem allows us to summarize our knowledge about unknown parameters after observing the data probabilistically (the ideal inference).
- ▶ But it requires treating the parameters as random variables and represent our *a priori* knowledge probabilistically as well.
- ▶ Sampling theory doesn't require this, but inference based on it must give up the ideal, and target at more limited inferential goals.
- ▶ We have looked at one particularly important example of such goals—the (point and interval) *estimation* problem.
- ▶ In the next few weeks we will focus on another important example—the *hypothesis testing* problem.

What is hypothesis testing?

- ▶ Estimation is about pinning down the underlying values of unknown parameters from a potentially *large (even infinite) number* of possibilities.
- ▶ In the simplest hypothesis testing setting, we ask a *dichotomous* question: If the unknown parameter can take two different values (or two different sets of values), which of the two should be inferred from the data?
- ▶ In other words, among two *hypotheses* about the parameter value, which should be accepted and which rejected?
- ▶ Let us look at three examples to have a feel of how such problems may arise.

Example I: Zip code recognition

- ▶ The parameter θ here is the underlying true number from “0” to “9”.
- ▶ The data is a 200×300 black/white pixels. How many possible values can the data be?
- ▶ $2^{200 \times 300}$ possible values!
- ▶ A probability model is a specification of the probability distribution $p(x|\theta)$ on these $2^{200 \times 300}$ values.

Testing a hypothesis

- ▶ For a single observed pattern X that could have arisen from either “0” or “6”, which should be assigned? (Draw a figure.)
- ▶ Under a Bayesian perspective, this decision seems straightforward. Let us assign prior probability on θ to characterize our *a priori* knowledge about its value.
- ▶ Apply Bayes’ theorem to get the posterior probabilities

$$P(\theta = \text{“0”}|x) \quad \text{and} \quad P(\theta = \text{“6”}|x).$$

- ▶ Based on this probabilistic summary an assignment can be made.
- ▶ More generally, introduce a loss function, $L(\theta, a)$ and Bayes rule is

$$\delta^*(x) = \operatorname{argmin}_a E(L(\theta, a) | x).$$

Verify: the above rule corresponds to the 0-1 loss

$$L(\theta, a) = \mathbf{1}(\theta \neq a).$$

- ▶ A Bayesian hypothesis testing problem takes exactly the same form as an estimation problem!

What if we want to adopt the sampling viewpoint?

- ▶ We can only use the model: $p(x|\theta)$, not the prior.
- ▶ One possibility: Assign $\theta = \text{"0"}$ or "6" based on whether $p(x|\theta) > 0.5$.
- ▶ This is problematic as $p(x|\theta)$ may be >0.5 (or <0.5) for both "0" and "6" !
- ▶ How about assign "0" or "6" based on whether

$$p(x|\theta = \text{"0"}) > p(x|\theta = \text{"6"})?$$

In other words, let us *compare the likelihood* at the two θ values and pick the one with larger likelihood.

- ▶ If we repeat the experiment many times, what would happen if $\theta = 0$ versus if $\theta = 6$.
- ▶ The idea of comparing the likelihoods under the two hypotheses seems to make sense intuitively.

Example II: Testing the quality of a lot

A computer company purchases many different electronic components from suppliers for making a laptop.

- ▶ Want to make sure that the components are of good quality in terms of life time.
- ▶ The quality of the components in each lot is usually similar, and one can model the life-time X of a component from a given lot to be an $\text{Exponential}(\lambda)$ random variable. So the expected life-time for the components in this lot is $E(X) = 1/\lambda$.
- ▶ Suppose a lot is considered good if $\lambda \leq 1.0$ and bad if $\lambda > 1.0$.
- ▶ How do we judge whether a lot is good or bad?
- ▶ Note that in this problem we are deciding between two *sets* of values for λ , instead of two specific values as in the previous example.

- ▶ Obviously we can't test every single component in a lot.
- ▶ A common strategy is to randomly *sample* a few, say n , components from the lot, and measure their life-time

$$X_1, X_2, \dots, X_n.$$

- ▶ How do we judge whether this lot is good or bad based on these life-times?
- ▶ An intuitive idea: How about we calculate the sample average life-time \bar{X} and see whether it is large or small?
- ▶ Is it enough to just compare \bar{X} with 1.0?
- ▶ *If we repeat the experiment many times*, what would happen if H_0 is true versus if H_1 is true?

Example III: Contingency tables

- ▶ In the 1880's, Francis Galton carried out a study on whether men and women choose their spouse on the basis of height.
- ▶ Do tall men tend to marry tall women, do short women tend to marry short men, or do people choose spouses regardless of each other's height?
- ▶ He was studying the heritability of height, and would like to know how he should take the correlation between parental heights into the analysis.
- ▶ He collected the following data set.

Galton's data

		Wife:		
		Tall	Medium	Short
Husband:	Tall	18	28	14
	Medium	20	51	28
	Short	12	25	9

- ▶ Question: Do these data support or refute the *hypothesis of independence* of spouses' heights?
- ▶ What is the sampling model here? What are the unknown parameters? How is the hypothesis formulated in terms of the unknown parameters?

Testing simple hypotheses

- ▶ Suppose we observe data $\mathbf{X} = (X_1, X_2, \dots, X_n)$ from some distribution $f(\mathbf{x}|\theta)$ or $p(\mathbf{x}|\theta)$.
- ▶ What is a *simple hypothesis*?
- ▶ A *simple hypothesis* is one that completely specifies the sampling distribution of the data $f(\mathbf{x}|\theta)$.
- ▶ For example, in the zipcode recognition example, if we know what

$$f(\mathbf{x}|\theta = \text{"0"}) \quad \text{and} \quad f(\mathbf{x}|\theta = \text{"6"})$$

are exactly then $\theta = \text{"0"}$ and $\theta = \text{"6"}$ are both simple hypotheses.

- ▶ In contrast, in the lot testing example, $\lambda \leq 1.0$ and $\lambda > 1.0$ do not completely specifies the distribution of $f(\mathbf{x}|\lambda)$.
- ▶ This kind of hypotheses, which specify the distribution $f(\mathbf{x}|\lambda)$ to be one of a collection of probability distributions, are called *composite* hypotheses.
- ▶ We will get to testing composite hypotheses later.
- ▶ Is $\lambda = 2.5$ is a simple or composite hypothesis?

Another example of simple versus composite hypotheses.

- ▶ Suppose our data are i.i.d. observations from a $N(\mu, \sigma^2)$ distribution.
- ▶ First consider the case when σ^2 is known.
- ▶ Is $\mu = 5$ a simple or composite hypothesis?
- ▶ Now what if σ^2 is unknown?

How about the the example of testing independence on a contingency table?

How do we compare two simple hypotheses

- ▶ This is the simplest situation for hypothesis testing.
- ▶ The two simple hypotheses can be formally treated as

$$H_0 : \theta = \theta_0 \quad \text{versus} \quad H_1 : \theta = \theta_1$$

where each of θ_0 and θ_1 completely specifies $f(\mathbf{x}|\theta)$.

- ▶ We make a choice between these two possibilities based on the data \mathbf{X} we observe.
- ▶ Let us consider this as a binary decision problem—reject or accept H_0 .
- ▶ When \mathbf{X} take values in a set of possible values \mathcal{R} , we will *reject* θ_0 and choose θ_1 .
- ▶ Otherwise, we accept, or do not reject, θ_0 .

The rejection region

- ▶ Therefore a test, i.e. a *decision rule* for choosing one of the two hypotheses based on the data, can be specified by the rejection region \mathcal{R} .

If \mathbf{X} is not in the rejection region \mathcal{R} decide $\theta = \theta_0$.

If \mathbf{X} is in the rejection region \mathcal{R} decide $\theta = \theta_1$.

- ▶ So constructing a test boils down to choosing the corresponding rejection region \mathcal{R} . That is, choosing the data values corresponding to each decision.

Examples of rejection regions

- ▶ Zipcode example. Consider a 200×300 black/white grid. Total of 2^{60000} possible \mathbf{X} values.
- ▶ There are numerous— 2^{60000} —different ways to define a rejection region.
- ▶ (Draw a figure.)

Examples of rejection regions

- ▶ Consider the lot testing example, where the data are continuous.
- ▶ The sample space (i.e. the collection of possible data values) is infinite.
- ▶ Infinite number of possible rejection regions.
- ▶ In particular, one can define rejection regions based on common statistics—such as sample mean, sample median and sample variance.

- ▶ This same formulation applies to testing composite hypotheses too.
- ▶ For the lot testing example, one possible rejection region may be

$$\mathcal{R} = \{(X_1, X_2, \dots, X_n) : \bar{X} < 1.0.\}$$

Another may be that

$$\mathcal{R} = \{(X_1, X_2, \dots, X_n) : \text{sample median}(X) < 1.0.\}$$

Yet another may be that

$$\mathcal{R} = \{(X_1, X_2, \dots, X_n) : \text{sample variance}(X) < 1.0.\}$$

The question is

- ▶ How do we choose the rejection region?
- ▶ It turns out that there is a general procedure with which one can construct good tests.
- ▶ The idea is related to the maximum likelihood principle we used for the estimation problem.
- ▶ Let's start with the testing of simple vs simple hypothesis.

Constructing good tests: comparing likelihoods

- ▶ How about we compare the likelihoods under the two hypotheses.
- ▶ Choose the hypothesis with the higher likelihood. That is

$$\mathcal{R} : \quad \text{Those } \mathbf{x} \text{ values such that } \frac{f(\mathbf{x}|\theta_1)}{f(\mathbf{x}|\theta_0)} > 1.$$

- ▶ The “best” test is very much along this line!

The likelihood ratio test

- ▶ The *likelihood ratio statistic*

$$\frac{f(\mathbf{X}|\theta_1)}{f(\mathbf{X}|\theta_0)}$$

measures the relative evidence for the data under the two hypotheses.

- ▶ A rejection region based on this statistic is

$$\frac{f(\mathbf{X}|\theta_1)}{f(\mathbf{X}|\theta_0)} > K$$

for some constant K .

- ▶ Tests with rejection regions of this form are called *likelihood ratio tests*.
- ▶ It turns out that these tests are the “best” test to use (under certain criteria).
- ▶ Next time we will introduce a few notions to make precise what “good”, and “best” mean.