

**Contribution of the Late Positive Potential to Emotional Memory: A Joint Neuro-Cognitive
Approach to Memory Prediction**

Robin Hellerstedt¹, Jordan Gunn², and Deborah Talmi²

¹University

²University of Cambridge

Author Note

Contribution of the Late Positive Potential to Emotional Memory: A Joint Neuro-Cognitive Approach to Memory Prediction

Introduction

Retrieved-context theory ([Howard & Kahana, 2002](#)) describes the state-of-the-art in terms of the rules that govern the chances that an experience we have now will come to mind later. It is supported by extensive behavioural ([Healey & Kahana, 2016](#); [Lohnas et al., 2015](#); [Sederberg et al., 2008](#)) and neural ([Folkerts et al., 2018](#); [Kragel et al., 2021](#); [Manning et al., 2011](#); [Polyn et al., 2005](#)) evidence (for a recent review, see [Lohnas & Healey, 2021](#)). According to this theory, items become associated with a gradually-drifting internal context during encoding. This association functions bidirectionally: retrieving an item reinstates its associated context, which can then cue further recall. This process accounts for the tendency to consecutively recall items that were studied nearby in time (temporal contiguity effects) and, more broadly, items that shared similar encoding contexts. Polyn et al. ([2009](#)) extended retrieved-context theory to nontemporal dimensions of context, including pre-experimental semantic associations and source features that capture shared encoding conditions. For example, items presented in the same modality share a source context that promotes their co-recall.

Within this framework, emotional Context-Maintenance and Retrieval ([Talimi et al., 2019](#)) (eCMR) extends CMR by introducing an emotional dimension of context and by allowing emotion to modulate encoding strength. The first mechanism captures emotional similarity among items; the second captures preferential attention to emotional stimuli. The basis for emotional similarity is that people process emotional experiences differently than neutral ones in terms of appraisal ([Sander et al., 2005](#)), attention ([Pourtois et al., 2013](#)), physiology ([Stephens et al., 2010](#)) and feelings. Because emotional items share these processing characteristics, they are more similar to one another than to neutral items ([Riberto et al., 2022](#)). Emotionality is represented as a source context dimension. Emotional items share a common emotional source context and neutral items share a distinct neutral source context, so that recalling an item from either category promotes further recall of items from the same category. This feature is used not only to describe

which items are similar to each other, but also to model the cognitive repercussions of their unique features.

It is known that emotional items enjoy preferential processing during encoding (Anderson, 2005; MacKay et al., 2004; Pourtois et al., 2013; Schupp et al., 2006). eCMR simulates this by scaling the learning rate for associations from context to item features (M^{CF}) during encoding, so that emotional items form stronger bindings to their encoding context. The degree of extra processing is governed by the value of parameter Φ_{emot} . When it equals 1, emotional and neutral items receive the same encoding strength. When it is greater than 1, emotional items are modelled as receiving extra processing during encoding. Constraining Φ_{emot} with an independent neural measure, rather than treating it as a free parameter, would provide a more principled account of individual differences in emotional memory and a stronger test of the theory (Turner et al., 2016). An excellent candidate is the late positive potential (LPP), an event-related potential (ERP) component that increases in amplitude when participants process emotionally significant visual stimuli (Cuthbert et al., 2000; Hajcak et al., 2010; Schupp et al., 2000, 2006). In this project, our aim was to examine the suitability of the LPP for neuro-cognitive modelling of memory recall. We pursued this aim in three steps: by establishing individual-level reliability of the LPP under memory-experiment conditions, by testing whether LPP variation predicts emotional memory, and by investigating how best to incorporate the neural measure into the model.

The LPP is broadly characterised as reflecting affective stimulus significance (Hajcak & Foti, 2020; Olofsson et al., 2008), and its emotional modulation has been observed at the group level across a range of stimuli and presentation durations. However, for the LPP to serve as a useful candidate marker of emotional encoding, it must also differentiate emotional from neutral processing at the individual level. Recently, Schupp and Kirmse (2021) examined individual-level sensitivity to emotional content using a case-by-case approach across three studies with different emotional stimulus categories. They observed that the LPP was larger for high- compared to low-arousing stimuli in 98% of individual-level tests, and a complementary specificity analysis confirmed that this effect was driven by emotional content rather than low-level stimulus

differences. Because eCMR attributes the emotional encoding advantage to preferential processing during study, and the LPP is sensitive to the enhanced processing of motivationally significant stimuli, these findings motivate testing whether LPP variation can usefully constrain Φ_{emot} .

For the LPP to serve this constraining role, however, the same individual-level reliability must hold under conditions typical of memory experiments. Although emotional modulation of the LPP has been observed at the group level in intentional-encoding paradigms followed by free recall (Barnacle et al., 2018), it is not yet known whether comparable individual-level sensitivity is preserved. Participants in Schupp and Kirmse (2021)'s study encoded briefly-presented pictures passively. By contrast, participants in typical memory recall experiments are aware of an upcoming memory test. Intentional encoding instructions could dilute the emotional modulation of the LPP in several ways. Participants may shift attention from processing each stimulus in isolation to comparing stimuli or organising them for later recall. They may also adopt encoding strategies that reduce processing differences between emotional and neutral items, or engage in rehearsal that redistributes attention more evenly across stimulus types. The potential impact of encoding mode is supported by evidence that task demands modulate emotional ERP effects (Schindler & Straube, 2020) and that encoding instructions alter the organisation of recall (Healey et al., 2019). Our first aim, therefore, was to replicate Schupp and Kirmse (2021)'s findings using a setup more typical in memory research. In addition to testing whether the LPP's emotional modulation survives intentional encoding, a replication would increase confidence in the generality of the findings. Relative to Schupp and Kirmse (2021), the present study uses a different stimulus set, a longer presentation duration (2 s vs. 150 ms), and a larger sample ($n = 38$).

Beyond individual-level reliability, our second aim was to examine whether the emotional modulation of the LPP predicts the emotional modulation of memory. Attention and memory are typically coupled, such that when all else is held equal, increased attention during encoding increases participants' ability to recall attended stimuli subsequently (Chun & Turk-Browne, 2007). Increased attention to emotional stimuli during encoding is considered a key driver of

enhanced memory for these stimuli (Mather & Sutherland, 2011; Talmi, 2013). Accordingly, previous studies report that emotional arousal increased both LPP and memory (Dolcos & Cabeza, 2002). Yet direct tests of whether the LPP–memory relationship is consistent within or between subjects are scarce (Fields, 2023). Indeed, Barnacle et al. (2018) found that the emotional modulation of the LPP was equally strong in pure and mixed lists under intentional encoding conditions, despite the fact that emotional memory enhancement was context-dependent. These results suggest a functional dissociation between attention-related LPP modulation and the downstream memory advantage. Understanding how LPP variation relates to memory within a mechanistic model may help bridge this gap. At the between-subjects level, individuals with larger emotion-dependent LPP differences may exhibit larger emotion-dependent memory differences. At the within-subject level, individuals may recall best those specific lists where encoding was accompanied by a larger LPP. The dissociation between within- and between-subject effects in ERPs was recently demonstrated for the left parietal old-new effect, where robust within-subject effects were observed despite null between-subject effects (MacLeod & Donaldson, 2017). Although the emotional modulation of the LPP shows good internal consistency (Moran et al., 2013) and is reliable within individuals across testing sessions (Weinberg et al., 2021), it is unknown whether this reliability extends to predicting memory differences. If trial-level LPP variation tracks encoding-related processing relevant to Φ_{emot} , the model predicts a positive association between the emotional modulation of the LPP and the emotional recall advantage, at the within-subject level, the between-subjects level, or both (Figure 1).

In addition, we wanted to investigate whether the relation between the LPP and emotion enhanced memory is better captured on the single-trial level or when averaged over all the items in a list to decide at what level we should add the LPP to the eCMR. It could for example be that the LPP is too noisy at the single-trial level to be useful, or that the loss of variability in LPP amplitude between items after averaging leads to a weaker relationship between the LPP and emotion enhanced memory. It was therefore important to examine at what level the relationship between the LPP and subsequent memory is stronger to inform our modelling decisions.

Establishing that the LPP is a reliable and behaviorally relevant signal at the individual and trial level is a necessary precondition for using it as a model input. The question of whether it actually improves model predictions was tested separately at the population level. We used these results to constrain a process model of memory encoding and retrieval. CMR captures temporal-context encoding and retrieval but has no emotion mechanism, and so provides a floor. CMR+LPP tests whether the LPP can serve as a substitute for category labels: it uses the same single-context architecture as CMR but scales encoding strength by trial-level LPP amplitude without any emotional context dynamics. eCMR adds an emotional context layer and emotion-modulated encoding strength, using only category labels. LPP-eCMR translates the emotion-specific LPP effect identified by the GLMM into the model's per-item encoding-strength term. The comparison tests whether trial-level LPP can substitute for category labels and whether embedding the LPP in a mechanistic retrieval model improves prediction beyond what category labels alone provide. We also examine whether the resulting model generates qualitatively distinct predictions about neural-behavioral relationships. We assess each model's population-level predictive validity using pooled fits and examine subject-wise fits secondarily to characterize individual heterogeneity.

The four models cross two dimensions: the presence or absence of an emotional context layer (which uses category labels to distinguish emotional from neutral items) and the use of trial-level LPP to modulate encoding strength. CMR+LPP and eCMR each have 10 free parameters, differing only in whether emotional memory is driven by a continuous neural signal or categorical source features. eCMR and LPP-eCMR share identical architecture, differing only in how LPP enters the encoding-strength term, so differences in predictive performance between them are attributable to the LPP term alone. Comparing alternatives that differ in retrieval dynamics would require output-order information that the present data lack.

Method

This secondary data analysis is based on data from Zarubin et al. (2020). A more detailed description of the methods in the original experiment is provided in paper.

Participants

The forty included participants from the mixed condition in Zarubin et al. (2020) were re-analysed. One participant was excluded for missing data from three lists and one participant was excluded due to having fewer than 16 trials in one of the subsequent memory conditions. The final sample consisted of 38 participants (28 females, $M_{\text{age}} = 20$, $SD_{\text{age}} = 1.07$). Gender and age data was missing for one of the participants. All participants provided written informed consent and the experiment was approved by Wofford College Institutional Review Board (Zarubin et al., 2020).

Materials

There were 22 pictures in each list and nine lists in the whole experiment. The first two pictures were used as fillers to control for primacy effects and were excluded from data analysis. The pictures were selected from the International Affective Picture System (Lang et al., 2005), the Geneva Affective Picture Database (Dan-Glauser & Scherer, 2011), the Emotional Picture Set (Wessa et al., 2010), the image pool of Talmi et al. (2007) and Google Images. The pictures were either of negative valence or neutral valence. The neutral condition was divided into semantically unrelated and semantically related neutral pictures in the original study, but they were collapsed in this secondary data analysis to increase the number of trials in the neutral condition.

Procedure

The participants studied 22 pictures shown consecutively on the screen for 2000 ms with a 4000 ms inter-trial interval, intended to reduce emotional carryover effects (Talmi & McGarry, 2012). The presentation order of both the lists and the images within lists was randomised. A one minute arithmetic distracter task followed immediately after the study task to reduce rehearsal of the pictures and recency effects. A verbal recall test followed the distracter task. The participants were instructed to recall images only from the list that they had just seen. The participants had three minutes to recall the images from the study phase, but could stop sooner if they had no more items to report.

EEG pre-processing

Raw EEG files were downloaded from the open science framework page for the Zarubin et al. (2020) study. The EEG was pre-processed with the EEGLAB toolbox in MATLAB (Delorme & Makeig, 2004). The data was re-referenced offline to the average of the left and right mastoid electrodes and filtered between 0.1 and 40Hz with a band pass FIR filter. The data was time-locked to the stimulus and segmented into epochs from –2000 ms to 3000 ms post stimulus onset. The epochs were baseline corrected using a pre-stimulus time window from –200 ms to 0 ms. Customized threshold functions from the FASTER toolbox (Nolan et al., 2010) were used to detect and reject bad channels and bad epochs based on participant-level z-transformed values over trials and electrodes that exceeded ± 3 . Independent component analysis was used to correct for EOG artefacts, by manually removing components classified as blinks, horizontal or vertical eye-movements. Bad channels which were deleted prior to ICA were interpolated after the ICA cleaning. In addition, bad channels within single epochs were detected and interpolated with a criterion of ± 3 SD. All participants contributed at least 16 trials in each of the four ERP conditions. The mean number of trials was 30.7 ($SD = 6.08$) in the negative remembered condition, 26.1 ($SD = 5.54$) in the negative forgotten condition, 48 ($SD = 15.33$) in the neutral remembered condition and 66.5 ($SD = 16.93$) in the neutral forgotten condition after pre-processing. Trials that were rejected during EEG preprocessing were excluded from the behavioural analyses as well to ensure that the EEG and behavioural data contained the same trials and to increase the chance of finding a relation between the two measures.

Behavioural Data Analysis

We investigated whether there was an emotion enhanced memory effect by comparing the proportion of negative items recalled and the proportion of neutral items recalled with a paired t-test.

EEG Data Analysis

Focal ERP Analysis. We first analysed the electrodes and time windows where the LPP is typically maximal in the literature. Averaged EEG amplitude was extracted from four

centroparietal electrodes, Cz, CP1, CP2 and Pz, in an early time window between 400–1000 ms and a late time window between 1000–2000 ms post stimulus onset. We extracted the EEG amplitudes separately for subsequently remembered and subsequently forgotten trials for both the negative and the neutral condition. All mixed effects model analyses were conducted using the lme4 package in R (Bates et al., 2015).

We first performed a linear mixed effects model to analyse if ERP amplitude was predicted by emotion and subsequent memory. In this model we used ERP amplitude as the dependent variable, Emotion (Negative vs Neutral) and Subsequent memory (Remembered vs Forgotten) as fixed effects, and random intercept + random slope effects of participant. The formula was:

$$\text{ERP amplitude} \sim 1 + \text{Emotion} \times \text{Memory} + (1 + \text{Emotion} \times \text{Memory} \mid \text{Participant})$$

The random effect structure needed to be simplified to only include a random intercept effect of participant in the late LPP time window due to a warning of singular fit. The statistical significance of the fixed effects and their interaction was assessed using χ^2 Wald test as implemented in the car package (Fox & Weisberg, 2019). Significant interactions were followed-up using pairwise contrasts as implemented in the emmeans package in R (Lenth, 2024) applying a Benjamini & Hochberg (false discovery rate, FDR) correction for multiple comparisons.

Global ERP Analysis. We complemented the focal analysis with a global data-driven analysis including all data points between 0 and 2000 ms, the time window when the stimulus was presented. The reason for including this analysis is that analysing ERP data from a small group of electrode sites and time windows may overlook effects at other electrode sites and time windows. In this analysis we controlled for multiple comparisons using cluster-based permutation tests as implemented in the FieldTrip toolbox (Maris & Oostenveld, 2007; Oostenveld et al., 2011). The first step of this analysis is to perform *t*-tests at every ERP data sample. Significant samples, uncorrected at $\alpha = .05$, that were neighbours in time or space, including at least two electrodes, were grouped into clusters. The cluster level *t*-value was computed by calculating the sum of the

t -values from all samples included in the cluster. Finally, we compared the cluster level t -value in the observed data to the size of the maximal cluster in 5000 permutations, for which the datapoints were randomly swapped between conditions within participants. The cluster level Monte-Carlo p -value was calculated as the proportion of the permutations for which the maximum cluster was larger than the clusters in the observed data. The exact spatio-temporal distribution of the clusters should be interpreted with caution given that the correction for multiple comparisons are applied at the cluster level rather than at the individual sample level ([Sassenhagen & Draschkow, 2019](#)).

Bootstrap Analysis. To be useful for the purpose of model constraint, the LPP needs to be reliable within subjects. We therefore performed a bootstrap analysis to assess the reliability of the Negative–Neutral difference within each participant’s data. The rationale of this analysis is that negative stimuli should elicit a more positive LPP than neutral stimuli independent of which specific trials contributed to calculating the ERPs.

The number of trials was first equated for the two conditions with undersampling. The data included in this analysis was extracted from the same centro-parietal electrode sites as in the focal ERP analysis, i.e. Cz, CP1, CP2 and Pz. Bootstrap resampling from single trials was used to create a distribution of 1000 resampled ERPs per condition. A sliding time window was applied to detect the most positive LPP peak (the mean amplitude of the 100 ms time window between 400–1000 ms for the early LPP and between 1000–2000 ms for the late LPP). Difference scores were calculated by subtracting the amplitude in the neutral condition from the amplitude in the negative condition for each bootstrap sample. Next, the proportion of positive bootstrap differences were calculated for the two time-windows based on the rationale that participants with robust LPP effects should have a large proportion of positive bootstrapped differences.

We interpreted the emotion enhanced LPP as being reliable if a proportion of 0.9 or more of the bootstrapped differences were positive because this cutoff has been used in similar analyses in the concealed information test literature (for example [Rosenfeld, 2020](#)).

Relationship between the LPP and Emotion Enhanced Memory. The ERPs were z -transformed using the mean and the standard deviation of a pre-stimulus interval from –200 to 0

ms. This was done on a single-trial level and separately for all electrodes. Next, the z-transformed LPP was extracted from the same centroparietal electrodes as in the focal ERP analysis for both the early, 400–1000 ms, and the late, 1000–2000 ms, time window.

In the between subjects correlation, we calculated the average LPP and memory performance across the 9 lists and computed a negative–neutral difference score in both measures. These measures were then correlated using Bayesian Pearson correlation.

Besides this between subject correlation we also calculated a within-subject correlation. In this analysis, we correlated the emotion enhanced LPP with the emotion enhanced memory across the 9 lists and calculated the average correlation coefficient across lists for each participant with Pearson correlation. Next, we performed a Bayesian one-sample t -test against zero to test if there was a correlation.

Finally, we investigated the relation between memory performance and the LPP on the single-trial level with a binomial generalized linear mixed-effect model where we investigated if Subsequent memory was predicted by Emotion and LPP amplitude. In this model we used Subsequent memory as the dependent variable and included Emotion (Negative vs Neutral) and LPP amplitude as fixed effects and random intercept + random slope effects of Participant. The model formula was:

$$\text{Memory} \sim 1 + \text{Emotion} \times \text{LPP} + (1 + \text{Emotion} \times \text{LPP} \mid \text{Participant})$$

eCMR Specification

Building on the behavioural and ERP results reported above, we specify the eCMR family used to test mechanistic accounts of how the LPP can modulate encoding within the eCMR framework (Talimi et al., 2019). Where LPP is included, it enters as an item-level modulator of emotional encoding strength.

We use the full eCMR architecture, which maintains two parallel context layers — a temporal context c^T and an emotional context c^E — with separate bidirectional association matrices for each pathway. Items are represented by one-hot feature vectors f_i . Each item also

carries a binary emotion feature vector s_i that codes emotional category: $s_i = [1, 0]$ for emotional items and $s_i = [0, 1]$ for neutral items. The emotional context includes feature dimensions corresponding to the emotional and neutral categories, so emotional and neutral items cue different emotional-context states during encoding and retrieval. The eCMR-based variants share this dual-context architecture; they differ in whether and how LPP modulates encoding strength within the emotional pathway.

The temporal pathway includes feature-to-context associations M^{FC} (items to temporal context) and context-to-feature associations M^{CF} (temporal context to items). The emotional pathway includes analogous associations M_{emot}^{FC} (items to emotional context) and M_{emot}^{CF} (emotional context to items).

Pre-experimental memory is initialized as

$$M_{pre(ij)}^{FC} = \begin{cases} 1 - \gamma & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

$$M_{pre(ij)}^{CF} = \begin{cases} \delta & \text{if } i = j \\ \alpha & \text{if } i \neq j \end{cases}$$

Pre-experimental item-to-context associations in M_{emot}^{FC} are initialized so that each item cues its fixed emotion feature vector s_i with weight $(1 - \gamma)$: emotional items cue the emotional unit and neutral items cue the neutral unit. M_{emot}^{CF} is initialized to zero, so emotional context-to-item associations are acquired only through encoding.

During encoding, temporal contextual input is retrieved as $c_i^{IN} = M^{FC} f_i$ and normalized to unit length. Temporal context updates according to

$$c_i^T = \rho_i c_{i-1}^T + \beta_{enc} c_i^{IN}$$

$$\rho_i = \sqrt{1 + \beta_{enc}^2 \left[\left(c_{i-1}^T \cdot c_i^{IN} \right)^2 - 1 \right]} - \beta_{enc} \left(c_{i-1}^T \cdot c_i^{IN} \right)$$

Emotional context drifts toward the feature cued by the presented item via

$$c_i^{E,IN} = M_{emot}^{FC} f_i:$$

$$c_i^E = \rho_i^E c_{i-1}^E + \beta_{emot} c_i^{E,IN}$$

with ρ_i^E computed analogously. Both emotional and neutral items update emotional context, pushing it toward the feature associated with that item.

Feature-to-context learning proceeds identically for both pathways:

$$\Delta M^{FC} = \gamma f_i c_i^{T\top}, \quad \Delta M_{emot}^{FC} = \gamma f_i c_i^{E\top}$$

The primacy-graded attention term is:

$$\phi_i = \phi_s e^{-\phi_d(i-1)} + 1$$

Following Talmi et al. (2019, Eq. 11), context-to-feature learning differs between pathways. The temporal pathway uses primacy only:

$$\Delta M^{CF} = \phi_i \cdot c_i^T f_i^\top$$

The emotional pathway is additionally scaled by the per-item emotional encoding strength $\phi_{emot,i}$:

$$\Delta M_{emot}^{CF} = \phi_i \cdot \phi_{emot,i} \cdot c_i^E f_i^\top$$

The parameterization of $\phi_{emot,i}$ defines each model variant (see below). Negative values of $\phi_{emot,i}$ are clamped to 0.

At recall onset, both contexts are shifted toward their respective pre-list states with drift rate β_{start} . At each recall step, temporal and emotional contexts jointly cue item activations:

$$A_i = \left(M^{CF} c^T \right)_i + \left(M_{emot}^{CF} c^E \right)_i$$

$$P(i) = \frac{A_i^{\tau_c}}{\sum_k A_k^{\tau_c}}$$

After recall, the retrieved item’s context is reinstated in both pathways via M^{FC} and M_{emot}^{FC} , each integrated with drift rate β_{rec} . We do not model recall termination explicitly and instead condition each simulation on the number of items recalled on that list. We make this scope choice because the present model comparison is aimed at the structure of item selection within recalled sets, whereas adding a stopping rule would introduce an additional submodel that is not central to that comparison. This is nevertheless a simplification: in free recall, stopping may depend on retrieval state, and the present analysis does not attempt to model that dependency.

Model Variants

We compare four models that cross two factors: the emotional context architecture (absent vs. present) and the information source for encoding modulation (LPP vs. category labels). The model architecture represents a binary emotional-vs-neutral distinction but does not represent pre-experimental semantic similarity among studied items. Semantically related and unrelated neutral items were therefore collapsed (see Materials), which may compress theoretically meaningful semantic variation. Let e_i be an indicator that equals 1 for emotional items and 0 for neutral items, and let L_i denote the trial-mean-centered Early LPP amplitude for item i .

The base model, CMR, has 9 free parameters and includes temporal context only. No emotional context layer is included and no emotion parameters are estimated.

CMR+LPP, with 10 free parameters, uses the same single-context architecture as CMR but adds one parameter κ_L scaling trial-level Early LPP amplitude into encoding strength:

$$\phi_{emot,i} = \kappa_L \cdot L_i$$

The encoding modulation applies to all items regardless of category and operates through the temporal context-to-feature pathway M^{CF} . This model tests whether a continuous neural measure of encoding-related processing, without any category labels or emotional context dynamics, can account for emotional memory effects.

eCMR, also with 10 free parameters, takes a different approach: rather than using continuous neural measures, it parameterizes emotion-modulated encoding strength with a category-based term ϕ_{emot} within the emotional context architecture described above:

$$\phi_{emot,i} = \phi_{emot} \cdot e_i$$

The model distinguishes emotional from neutral items via category labels but has no access to trial-level neural measures.

LPP-eCMR, with 11 free parameters, is a hypothesis-driven extension in which one parameter κ_{EL} scales the Early LPP's contribution to emotional encoding strength:

$$\phi_{emot,i} = \phi_{emot} \cdot e_i + \kappa_{EL} \cdot L_i \cdot e_i$$

This specification operationalizes the earlier GLMM result that Early LPP amplitude predicted recall for emotional but not neutral items ($\beta = .14, p < .001$ for emotional; $\beta = .01, p = .488$ for neutral). The e_i term therefore restricts the LPP's influence to emotional items, and no main-effect LPP term is included. A matched variant in which LPP is allowed to affect all items was not fit here. The emotional-only restriction is therefore a hypothesis-driven modelling choice motivated by the earlier analysis, not a conclusion established by the model comparison itself.

The four models form two nested pairs: CMR nests under CMR+LPP (setting $\kappa_L = 0$), and eCMR nests under LPP-eCMR (setting $\kappa_{EL} = 0$). CMR+LPP and eCMR are not nested because they differ in architecture (single vs. dual context) and information source (LPP vs. category labels). With 10 free parameters each, the comparison between CMR+LPP and eCMR tests whether trial-level LPP can replace category labels at equal parameter cost.

Evaluation Approach

We evaluated each model by how well it predicts the unordered set of items recalled from each list (not recall order), because our data is missing recall order information and because our primary behavioral outcomes and benchmarks target item selection rather than order-dependent

dynamics. Let S_t denote the set of items recalled on list t , and let r range over recall sequences that contain exactly the items in S_t (in any order). We define the *set likelihood* as

$$P(S_t | \theta) = \sum_{r: r \text{ yields } S_t} P(r | \theta),$$

where θ are the model parameters.

Evaluating this sum exactly is infeasible because the number of sequences that yield a given set grows rapidly with $|S_t|$. We therefore approximate $P(S_t | \theta)$ by Monte Carlo sampling: for each list we draw K random permutations $r_1, \dots, r_K$ of S_t , compute $P(r_k | \theta)$, and estimate the sum as $|S_t|! \frac{1}{K} \sum_{k=1}^K P(r_k | \theta)$. Because the factorial factor does not depend on θ , we maximize the (unscaled) mean permutation likelihood $\frac{1}{K} \sum_k P(r_k | \theta)$, which is proportional to the set likelihood and yields identical fits and ΔAIC comparisons.

We maximize the objective with differential evolution (Storn & Price, 1997). Because the optimizer is stochastic, we run three independent fits per model variant and retain the best-performing fit. We compare models using AIC computed from the approximate set log-likelihood.

The primary model comparisons use pooled fits, in which a single shared parameter vector is fit to all 342 lists across 38 subjects. Pooled fits target population-level predictive validity. They also avoid the small-sample problem that arises in subject-wise fits: with only 9 lists per subject and 9–11 free parameters, the standard AICc correction (Burnham & Anderson, 2002) is unavailable because $n \leq k + 1$. Subject-wise (no-pooling) fits are reported secondarily to characterize heterogeneity. Unlike pooled fits, which ask whether a single shared parameter vector predicts recall sets well at the population level, subject-wise fits ask how model performance varies when each participant is allowed their own parameter vector (Gelman & Hill, 2006). For nested model pairs in the subject-wise analysis, occasional optimizer failures can make the more flexible model fit worse than its simpler parent for a given subject, even though the models are nested. When that occurred, we replaced the more flexible model’s negative log-likelihood with the parent model’s value before computing pairwise comparisons. Pairwise ΔAIC with 95% CIs

are reported for the key subject-wise contrasts.

We assess qualitative adequacy by simulating each fitted model on the study lists while matching the number of items recalled on each list and plotting benchmark summaries. This set-likelihood approach evaluates models on their ability to reproduce the full recalled sets, providing a stringent test of mechanistic hypotheses.

Results

Behavioural Results

Replicating the original analysis, a paired t-test showed that memory performance was significantly higher for negative items ($M = 0.54$, $SD = 0.1$) compared to neutral items ($M = 0.42$, $SD = .14$), $t(37) = 6.18$, $p < .001$.

For the record. If we run a one-way repeated measures ANOVA with proportion remembered as the dependent variable and condition (Negative vs Category vs Neutral) as the independent variable, we get a significant effect $F(1,65) = 42.85$, $p < .001$, partial eta squared = 0.54. Follow up pairwise comparisons shows that Negative items were remembered more than both category items ($t(37) = 2.46$, $p = .01$) and neutral items ($t(37) = 9.41$, $p < .001$). Memory performance was also higher for category items (semantically related neutral items) than for neutral items (semantically unrelated neutral items; $t(37) = 7.29$, $p < .001$). The pattern of results was hence Negative > Category > Neutral.

EEG Results

Focal ERP Analysis. A linear mixed effect model was conducted to investigate if Emotion (Negative vs Neutral) and Subsequent memory (Remembered vs Forgotten) predicted ERP amplitude was conducted separately for the early and the late LPP time window.

In the early time window, there was a main effect of Emotion ($\beta = .10$, $\chi^2(1) = 6.84$, $p = .009$) and a main effect of Subsequent memory ($\beta = .25$, $\chi^2(1) = 26.70$, $p < .001$) and a significant Emotion x Subsequent memory interaction ($\beta = .20$, $\chi^2(1) = 14.70$, $p < .001$). Follow-up pairwise contrasts showed that the LPP amplitude was higher for subsequently remembered than for subsequently forgotten negative items ($z = -5.61$, $p < .001$), while there was no difference in LPP

amplitude for subsequently remembered and forgotten neutral items ($z = -.88, p = .379$).

In the late time window, there was a main effect of Emotion ($\beta = .14, \chi^2(1) = 12.72, p < .001$), indicating that the LPP amplitude was higher for negative than for neutral items. There was no significant main effect of Subsequent memory ($\beta = .08, \chi^2(1) = 2.13, p = .144$). There was a trend for a significant Emotion x Subsequent memory interaction ($\beta = .09, \chi^2(1) = 3.06, p = .080$).

Global ERP Analysis. We first analysed the main effect of Emotion (Negative vs Neutral trials) and found a positive cluster ($p < .001$) which was significant between approximately 380-2000ms with a widespread distribution. In other words, there was an emotionally enhanced LPP effect which roughly overlapped with the early and late time windows analysed in the focal analysis.

Replicating the focal analysis, there was also a main effect of subsequent memory with more positive ERPs for remembered compared to forgotten trials ($p < .001$) with a widespread spatial distribution between around 210-1240ms.

In addition, there was an Emotion x Subsequent Memory interaction between 360-1240ms ($p < .001$) with a widespread distribution, indicating that the negative items subsequent memory effect was larger than the neutral items subsequent memory effect. In addition, there was a borderline significant widespread cluster ($p = .029$) between 1240-1390ms. Follow-up cluster permutation tests showed that for negative items, remembered items were related to more positive going ERPs between 220-1420ms ($p < .001$). This effect was also widespread. In contrast, there was no significant subsequent memory (all $ps > .19$). ERP effect for neutral items.

Overall, the global analysis replicated the focal analysis and shows that these results hold up in more conservative tests with correction for multiple comparisons. The analysis also shows that the time windows and electrodes analysed in the focal analysis capture the maximum of the LPP effects well in this data set.

Bootstrap Reliability. We next performed a bootstrap analysis to test if the emotional LPP effect was reliable within subjects. This analysis showed that the early LPP was reliable in 73.7% of the participants while the late LPP effect was reliable in 50% of the participants.

LPP and Emotion Enhanced Memory. For the LPP to be useful as a model constraint, it needs to relate to memory performance at a relevant level of analysis. We therefore examined that relation at the between-subject, within-subject, and single-trial levels in turn. We first investigated whether the emotionally enhanced LPP amplitude (negative minus neutral items) correlated with emotionally enhanced memory (negative minus neutral items) between subjects. Surprisingly, the Bayesian Pearson correlation showed moderate evidence for no correlation for both the early time window ($r = -.018$, $BF_{01} = 4.925$) and the late time window ($r = -.028$, $BF_{01} = 4.885$).

Next, we investigated the correlation between emotion enhanced LPP amplitude and emotion enhanced memory within subjects. For each subject, we correlated the emotion enhanced LPP with the emotion enhanced memory across the 9 lists. We then calculated the average correlation coefficient for each subject and tested it against zero using a one-sample Bayesian t -test. There was moderate support for no correlation in both the early time window ($t(37) = -.26$, $p = .795$, $BF_{01} = 7.658$) and the late time window ($t(37) = -1.26$, $p = .215$, $BF_{01} = 3.698$).

Finally, we conducted binomial generalized mixed effects models to test if Emotion and LPP amplitude predicted memory separately for the two time windows. In the early time window, there was a main effect of Emotion ($\beta = .21$, $\chi^2(1) = 22.68$, $p < .001$), a main effect of LPP ($\beta = .08$, $\chi^2(1) = 18.87$, $p < .001$) and an Emotion x LPP interaction ($\beta = .06$, $\chi^2(1) = 11.20$, $p < .001$). The interaction was followed-up by running the same analysis separately for negative and neutral items. For negative items, there was a significant positive relationship between early LPP amplitude and memory ($\beta = .14$, $\chi^2(1) = 25.43$, $p < .001$). In contrast, there was no significant relationship between early LPP amplitude and memory for neutral items ($\beta = .01$, $\chi^2(1) = .48$, $p = .488$). In the late time window, there was a main effect of Emotion ($\beta = .23$, $\chi^2(1) = 66.58$, $p < .001$), showing once again that memory was higher for negative than neutral items, but importantly there was neither a main effect of LPP amplitude ($\beta = .02$, $\chi^2(1) = 1.98$, $p = .160$) nor any interaction between Emotion and LPP amplitude ($\beta = .02$, $\chi^2(1) = 2.44$, $p = .118$).

In sum, the analyses show that the early LPP is more reliable and related to emotion-enhanced memory on a single-trial level. Based on this, we will use single-trial data

from the early LPP time window to constrain the eCMR model.

eCMR Model Comparison

Pooled Analysis. Table 2 summarizes the primary pooled model comparison, in which a single parameter vector per model was fit to all 342 lists. eCMR dominated both single-context models ($\Delta\text{AIC} = 67.26$ over CMR, 51.46 over CMR+LPP), supporting the eCMR specification tested here over the single-context alternatives. CMR+LPP improved on CMR ($\Delta\text{AIC} = 15.80$), indicating that the LPP carries a population-level signal about encoding strength even without an emotional context layer. However, a single-context model driven by LPP did not substitute for the eCMR specification tested here: with 10 free parameters each, eCMR substantially outperformed CMR+LPP.

LPP-eCMR achieved the best fit overall, improving on eCMR by $\Delta\text{AIC} = 34.16$. Supplementing category labels with trial-level LPP improved pooled predictive fit.

Subject-Wise Analysis. Separate subject-wise (no-pooling) fits were also examined. Because each subject contributed only 9 lists for models with 9–11 free parameters, these comparisons are methodologically fragile and are not treated as the primary inferential basis; they are reported to characterize heterogeneity. Nesting flooring was required for 23 of 152 subject–model entries.

The broad ordering of models was similar, but the LPP-related gains were weak and heterogeneous at the subject level. eCMR outperformed CMR (mean $\Delta\text{AIC} = 4.06$ [1.36, 6.77]). However, mean ΔAIC was 0.33 [−0.77, 1.44] for CMR+LPP over CMR and 0.30 [−0.70, 1.29] for LPP-eCMR over eCMR, with both CIs including zero.

Qualitative Diagnostics. We assessed qualitative adequacy with two benchmark diagnostics (Figure 11; Figure 12; Figure 13). The first targets the emotional enhancement of memory established in the Behavioural Results. Category-specific serial position curves (Figure 12) show that CMR, which lacks an emotional context layer, predicts nearly identical recall for negative and neutral items. CMR+LPP produces a faint separation because LPP amplitude correlates weakly with emotional category, but the effect is far weaker than what the

eCMR-based specifications achieve. Both eCMR and LPP-eCMR reproduce the empirical separation: their shared emotional context ensures that negative items promote one another's recall, lifting the negative SPC above the neutral baseline.

A second diagnostic targets the LPP–recall link established in the EEG Results. Empirically, recalled emotional items had higher Early LPP amplitudes than unrecalled emotional items; no such difference existed for neutral items. eCMR cannot reproduce this pattern: it assigns identical encoding strength to all emotional items regardless of their LPP amplitude, so simulated LPP distributions for recalled and unrecalled emotional items overlap. Both CMR+LPP and LPP-eCMR reproduce the separation, since both use LPP to modulate encoding strength. Higher LPP leads to stronger encoding and higher recall probability (Figure 13). The critical difference is that only LPP-eCMR also captures the category-level recall structure visible in the SPC diagnostic. This recalled-versus-unrecalled Early LPP separation was not directly fit: the models were optimized to predict recalled sets, not this neural-behavioral contrast. In that sense, LPP-eCMR generalizes beyond the fitting target, although the model form was informed by prior empirical results. LPP-eCMR captures a mechanistic link (LPP $\rightarrow$ encoding strength $\rightarrow$ recall probability) that produces the observed neural-behavioral association as a consequence of its architecture, while also reproducing the categorical emotional memory advantage.

The LPP thus improves pooled predictive fit and generates mechanistic predictions that were not used as fitting targets. However, it does not substitute for the eCMR specification tested here: CMR+LPP remains far behind eCMR, and subject-wise gains are small and heterogeneous under current data constraints. The value of incorporating trial-level neural measures lies in the population-level predictive improvement and in generating qualitatively distinct predictions about neural-behavioral relationships. One possible interpretation is that trial-to-trial Early LPP variation is more behaviorally meaningful for emotional items because it more strongly indexes affective significance or motivated attention during encoding. A weaker relationship for neutral items could still exist, but under the present data it may be too small or too noisy to detect.

Discussion

References

- Anderson, A. K. (2005). Affective influences on the attentional dynamics supporting awareness. *Journal of Experimental Psychology: General*, 134(2), 258–281.
<https://doi.org/10.1037/0096-3445.134.2.258>
- Barnacle, G. E., Tsivilis, D., Schaefer, A., & Talmi, D. (2018). Local context influences memory for emotional stimuli but not electrophysiological markers of emotion-dependent attention. *Psychophysiology*, 55(4), e13014. <https://doi.org/10.1111/psyp.13014>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information-theoretic approach* (2nd ed.). Springer. <https://doi.org/10.1007/b97636>
- Chun, M. M., & Turk-Browne, N. B. (2007). Interactions between attention and memory. *Current Opinion in Neurobiology*, 17(2), 177–184. <https://doi.org/10.1016/j.conb.2007.03.005>
- Cuthbert, B. N., Schupp, H. T., Bradley, M. M., Birbaumer, N., & Lang, P. J. (2000). Brain potentials in affective picture processing: Covariation with autonomic arousal and affective report. *Biological Psychology*, 52(2), 95–111.
[https://doi.org/10.1016/S0301-0511\(99\)00044-7](https://doi.org/10.1016/S0301-0511(99)00044-7)
- Dan-Glauser, E. S., & Scherer, K. R. (2011). The geneva affective picture database (GAPED): A new 730-picture database focusing on valence and normative significance. *Behavior Research Methods*, 43(2), 468–477. <https://doi.org/10.3758/s13428-011-0064-1>
- Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, 134(1), 9–21. <https://doi.org/10.1016/j.jneumeth.2003.10.009>
- Dolcos, F., & Cabeza, R. (2002). Event-related potentials of emotional memory: Encoding pleasant, unpleasant, and neutral pictures. *Cognitive, Affective, & Behavioral Neuroscience*, 2(3), 252–263. <https://doi.org/10.3758/CABN.2.3.252>

- Fields, E. C. (2023). The P300, the LPP, context updating, and memory: What is the functional significance of the emotion-related late positive potential? *International Journal of Psychophysiology*, 192, 43–52. <https://doi.org/10.1016/j.ijpsycho.2023.08.005>
- Folkerts, S., Rutishauser, U., & Howard, M. W. (2018). Human episodic memory retrieval is accompanied by a neural contiguity effect. *The Journal of Neuroscience*, 38(17), 4200–4211. <https://doi.org/10.1523/JNEUROSCI.2312-17.2018>
- Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (3rd ed.). Sage Publications.
- Gelman, A., & Hill, J. (2006). *Data analysis using regression and multilevel/hierarchical models*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511790942>
- Hajcak, G., & Foti, D. (2020). Significance?... Significance! Empirical, methodological, and theoretical connections between the late positive potential and P300 as neural responses to stimulus significance: An integrative review. *Psychophysiology*, 57(7), e13570. <https://doi.org/10.1111/psyp.13570>
- Hajcak, G., MacNamara, A., & Olvet, D. M. (2010). Event-related potentials, emotion, and emotion regulation: An integrative review. *Developmental Neuropsychology*, 35(2), 129–155. <https://doi.org/10.1080/87565640903526504>
- Healey, M. K., & Kahana, M. J. (2016). A four-component model of age-related memory change. *Psychological Review*, 123(1), 23–69. <https://doi.org/10.1037/rev0000015>
- Healey, M. K., Long, N. M., & Kahana, M. J. (2019). Contiguity in episodic memory. *Psychonomic Bulletin & Review*, 26(3), 699–720. <https://doi.org/10.3758/s13423-018-1537-3>
- Howard, M. W., & Kahana, M. J. (2002). A distributed representation of temporal context. *Journal of Mathematical Psychology*, 46(3), 269–299. <https://doi.org/10.1006/jmps.2001.1388>
- Kragel, J. E., Ezzyat, Y., Lega, B. C., Sperling, M. R., Worrell, G. A., Gross, R. E., Jobst, B. C., Sheth, S. A., Zaghoul, K. A., Stein, J. M., & Kahana, M. J. (2021). Distinct cortical systems reinstate the content and context of episodic memories. *Nature Communications*, 12, 4444.

<https://doi.org/10.1038/s41467-021-24393-1>

Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (2005). *International affective picture system (IAPS): Affective ratings of pictures and instruction manual*. University of Florida.

Lenth, R. V. (2024). *Emmeans: Estimated marginal means, aka least-squares means*.

<https://CRAN.R-project.org/package=emmeans>

Lohnas, L. J., & Healey, M. K. (2021). The role of context in episodic memory: Behavior and neurophysiology. In *Psychology of learning and motivation* (Vol. 75, pp. 157–199). Elsevier.

<https://doi.org/10.1016/bs.plm.2021.06.003>

Lohnas, L. J., Polyn, S. M., & Kahana, M. J. (2015). Expanding the scope of memory search: Modeling intralist and interlist effects in free recall. *Psychological Review*, 122(2), 337–363.

<https://doi.org/10.1037/a0039036>

MacKay, D. G., Shafto, M., Taylor, J. K., Marian, D. E., Abrams, L., & Dyer, J. R. (2004).

Relations between emotion, memory, and attention: Evidence from taboo Stroop, lexical decision, and immediate memory tasks. *Memory & Cognition*, 32(3), 474–488.

<https://doi.org/10.3758/BF03195840>

MacLeod, C. A., & Donaldson, D. I. (2017). Investigating the functional utility of the left parietal ERP old/new effect: Brain activity predicts within but not between participant variance in episodic recollection. *Frontiers in Human Neuroscience*, 11, 580.

<https://doi.org/10.3389/fnhum.2017.00580>

Manning, J. R., Polyn, S. M., Baltuch, G. H., Litt, B., & Kahana, M. J. (2011). Oscillatory patterns in temporal lobe reveal context reinstatement during memory search. *Proceedings of the National Academy of Sciences*, 108(31), 12893–12897.

<https://doi.org/10.1073/pnas.1015174108>

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, 164(1), 177–190.

<https://doi.org/10.1016/j.jneumeth.2007.03.024>

Mather, M., & Sutherland, M. R. (2011). Arousal-biased competition in perception and memory.

Perspectives on Psychological Science, 6(2), 114–133.

<https://doi.org/10.1177/1745691611400234>

Moran, T. P., Jendrusina, A. A., & Moser, J. S. (2013). The psychometric properties of the late positive potential during emotion processing and regulation. *Brain Research*, 1516, 66–75.

<https://doi.org/10.1016/j.brainres.2013.04.018>

Nolan, H., Whelan, R., & Reilly, R. B. (2010). FASTER: Fully automated statistical thresholding for EEG artifact rejection. *Journal of Neuroscience Methods*, 192(1), 152–162.

<https://doi.org/10.1016/j.jneumeth.2010.07.015>

Olofsson, J. K., Nordin, S., Sequeira, H., & Polich, J. (2008). Affective picture processing: An integrative review of ERP findings. *Biological Psychology*, 77(3), 247–265.

<https://doi.org/10.1016/j.biopsycho.2007.11.006>

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011, 156869. <https://doi.org/10.1155/2011/156869>

Polyn, S. M., Natu, V. S., Cohen, J. D., & Norman, K. A. (2005). Category-specific cortical activity precedes retrieval during memory search. *Science*, 310(5756), 1963–1966.

<https://doi.org/10.1126/science.1117645>

Polyn, S. M., Norman, K. A., & Kahana, M. J. (2009). A context maintenance and retrieval model of organizational processes in free recall. *Psychological Review*, 116(1), 129–156.

<https://doi.org/10.1037/a0014420>

Pourtois, G., Schettino, A., & Vuilleumier, P. (2013). Brain mechanisms for emotional influences on perception and attention: What is magic and what is not. *Biological Psychology*, 92(3), 492–512.

<https://doi.org/10.1016/j.biopsycho.2012.02.007>

Riberto, M., Paz, R., Pobric, G., & Talmi, D. (2022). The neural representations of emotional experiences are more similar than those of neutral experiences. *The Journal of Neuroscience*,

42(13), 2772–2785. <https://doi.org/10.1523/JNEUROSCI.1490-21.2022>

Rosenfeld, J. P. (2020). P300 in detecting concealed information and deception: A review.

- Psychophysiology*, 57(7), e13362. <https://doi.org/10.1111/psyp.13362>
- Sander, D., Grandjean, D., & Scherer, K. R. (2005). A systems approach to appraisal mechanisms in emotion. *Neural Networks*, 18(4), 317–352. <https://doi.org/10.1016/j.neunet.2005.03.001>
- Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology*, 56(6), e13335. <https://doi.org/10.1111/psyp.13335>
- Schindler, S., & Straube, T. (2020). Selective visual attention to emotional pictures: Interactions of task-relevance and emotion are restricted to the late positive potential. *Psychophysiology*, 57(9), e13585. <https://doi.org/10.1111/psyp.13585>
- Schupp, H. T., Cuthbert, B. N., Bradley, M. M., Cacioppo, J. T., Ito, T., & Lang, P. J. (2000). Affective picture processing: The late positive potential is modulated by motivational relevance. *Psychophysiology*, 37(2), 257–261. <https://doi.org/10.1111/1469-8986.3720257>
- Schupp, H. T., Flaisch, T., Stockburger, J., & Junghöfer, M. (2006). Emotion and attention: Event-related brain potential studies. In *Progress in brain research* (Vol. 156, pp. 31–51). [https://doi.org/10.1016/S0079-6123\(06\)56002-9](https://doi.org/10.1016/S0079-6123(06)56002-9)
- Schupp, H. T., & Kirmse, U. M. (2021). Case-by-case: Emotional stimulus significance and the modulation of the EPN and LPP. *Psychophysiology*, 58(4), e13766. <https://doi.org/10.1111/psyp.13766>
- Sederberg, P. B., Howard, M. W., & Kahana, M. J. (2008). A context-based theory of recency and contiguity in free recall. *Psychological Review*, 115(4), 893–912. <https://doi.org/10.1037/a0013396>
- Stephens, C. L., Christie, I. C., & Friedman, B. H. (2010). Autonomic specificity of basic emotions: Evidence from pattern classification and cluster analysis. *Biological Psychology*, 84(3), 463–473. <https://doi.org/10.1016/j.biopsycho.2010.03.014>
- Storn, R., & Price, K. (1997). Differential evolution – a simple and efficient heuristic for global optimization over continuous spaces. *Journal of Global Optimization*, 11(4), 341–359. <https://doi.org/10.1023/A:1008202821328>

- Talmi, D. (2013). Enhanced emotional memory: Cognitive and neural mechanisms. *Current Directions in Psychological Science*, 22(6), 430–436.
<https://doi.org/10.1177/0963721413498893>
- Talmi, D., Lohnas, L. J., & Daw, N. D. (2019). A retrieved context model of the emotional modulation of memory. *Psychological Review*, 126(4), 455–485.
<https://doi.org/10.1037/rev0000132>
- Talmi, D., Luk, B. T. C., McGarry, L. M., & Moscovitch, M. (2007). The contribution of relatedness and distinctiveness to emotionally-enhanced memory. *Journal of Memory and Language*, 56(4), 555–574. <https://doi.org/10.1016/j.jml.2007.01.002>
- Talmi, D., & McGarry, L. M. (2012). Accounting for immediate emotional memory enhancement. *Journal of Memory and Language*, 66(1), 93–108. <https://doi.org/10.1016/j.jml.2011.07.009>
- Turner, B. M., Rodriguez, C. A., Norcia, T. M., McClure, S. M., & Steyvers, M. (2016). Why more is better: Simultaneous modeling of EEG, fMRI, and behavioral data. *NeuroImage*, 128, 96–115. <https://doi.org/10.1016/j.neuroimage.2015.12.030>
- Weinberg, A., Correa, K. A., Stevens, E. S., & Shankman, S. A. (2021). The emotion-elicited late positive potential is stable across five testing sessions. *Psychophysiology*, 58(11), e13904.
<https://doi.org/10.1111/psyp.13904>
- Wessa, M., Kanske, P., Neumeister, P., Bode, K., Heissler, J., & Schönfelder, S. (2010). EmoPics: Subjektive und psychophysiologische evaluationen neuen bildmaterials für die klinisch-bio-psychologische forschung. *Zeitschrift für Klinische Psychologie Und Psychotherapie, Supplement*(1/11), 77.
- Zarubin, V. C., Phillips, T. K., Robertson, E., Bolton Swafford, P. G., Bunge, T., Aguiard, D., Martsberger, C., & Mickley Steinmetz, K. R. (2020). Contributions of arousal, attention, distinctiveness, and semantic relatedness to enhanced emotional memory: An event-related potential and electrocardiogram study. *Affective Science*, 1(3), 172–185.
<https://doi.org/10.1007/s42761-020-00012-y>

Table 1*Parameters and structures specifying eCMR.*

Symbol	Meaning
c^T	temporal context
c^E	emotional context
f_i	item feature vector (one-hot)
s_i	emotion feature vector (2-unit localist)
M^{FC}, M^{CF}	temporal pathway associations
$M_{emot}^{FC}, M_{emot}^{CF}$	emotional pathway associations
β_{enc}	temporal encoding drift rate
β_{emot}	emotional encoding drift rate
β_{start}	start-list drift rate
β_{rec}	recall drift rate
α	shared support in M^{CF}
δ	self-support in M^{CF}
γ	learning rate for ΔM^{FC}
ϕ_s	primacy scale
ϕ_d	primacy decay
τ_c	choice sensitivity
$\phi_{emot,i}$	emotional encoding strength
L_i	list-centered Early LPP
κ_L, λ_L	LPP main-effect slope, offset
$\kappa_{EL}, \lambda_{EL}$	LPP-by-emotion slope, offset

Table 2

Pooled model comparison. k = free parameters; $-LL$ = negative log-likelihood for a single shared parameter vector fit to all 342 lists across 38 subjects; ΔAIC relative to best model.

Model	k	$-LL$	AIC	ΔAIC
CMR	9	8317.75	16653.51	+101.42
CMR+LPP	10	8308.85	16637.71	+85.62
eCMR	10	8283.13	16586.25	+34.16
LPP-eCMR	11	8265.05	16552.09	0 (best)

Table 3

Subject-wise model comparison. k = free parameters; NLL = negative log-likelihood (mean $\pm$ SE across 38 participants).

Model	k	Mean NLL ($\pm$ SE)
CMR	9	215.01 $\pm$ 15.63
CMR+LPP	10	213.84 $\pm$ 15.73
eCMR	10	211.98 $\pm$ 16.31
LPP-eCMR	11	210.83 $\pm$ 16.46


Figure 1*Early LPP ERP waveforms (400–1000 ms)*


Figure 2*Early LPP topography and ERP differences*


Figure 3*Late LPP ERP waveforms (1000–2000 ms)*


Figure 4*Global ERP analysis: cluster summary*


Figure 5*Global ERP analysis: detailed cluster results*

Figure 6

Bootstrap reliability of the emotional LPP effect





Figure 7*Between-subjects LPP–memory correlation overview*

Figure 8

Early LPP between-subjects scatter plot





Figure 9*Late LPP between-subjects scatter plot*

Figure 10

Within-subjects LPP–memory correlation




Figure 11

Empirical benchmark diagnostics. Left: category-specific serial position curves showing the negative-above-neutral recall advantage. Right: Early LPP amplitude by recall status for negative items showing the recalled/unrecalled separation.




Figure 12

Simulated category-specific serial position curves. Top: CMR (left), CMR+LPP (right). Bottom: eCMR (left), LPP-eCMR (right). CMR predicts nearly identical curves; CMR+LPP produces a faint separation via indirect LPP-category correlation; eCMR and LPP-eCMR reproduce the full negative-above-neutral separation (compare Figure 11).




Figure 13

Simulated Early LPP amplitude by recall status for negative items. Top: CMR (left), CMR+LPP (right). Bottom: eCMR (left), LPP-eCMR (right). CMR+LPP and LPP-eCMR reproduce the recalled/unrecalled separation; CMR and eCMR cannot (compare Figure 11).

