

Using Statistical Dynamics to Explain Features of Anomalous Statistics in shallow water waves with Abrupt Depth Changes

A. Mayda, N. Moore (Di Qi?)

I) Experiments (B.S.M., 2018)

(Deterministic)

II) (Model for I) Initial KdV + Scattered KdV after abrupt depth change (ADC)

Ref. R.S. Johnson, ... Math Theory of Water Waves. Cam.U. Press (1997)

Read pgs 268-277

III) Statistical Dynamics Model for I) based on II)

Read ① Mayda + Wang book, Cam.U. Press (2006), Chapter 6

Read ② Abranov, Kovacic, Mayda, Comm. Pure Appl. Math (2003) 1-46

Read ③ Bajars, Frank, Leinukhler, Nonlinearity (2013) 1945-1973

Only Read 1948-1952 and 1956-1957 (Fig. 1)

Plan I)-II Detailed Notes

May 8/1/13

I Deterministic Dynamics Model with Abrupt Depth Change

Model Assumptions

Assume 1) Nearly unidirectional, periodic narrow channel $2\pi L$, $L_0 = L^{-1} \ll 1$

2) Role of paddle wheel forcing looking a little downstream

(A) is to generate stat soln with $M=0$, E_0
 where E_0 is function of paddle strength at $t=0$

3) $|K| \leq 1$ fixed truncation $u_x \equiv u$

4) time $t \Leftrightarrow$ downstream coordinate, abrupt depth change at $t=t_0$
 (ADC) $t_0 \rightarrow T_{ADC}$

Use notation from 7/16/18 for non-lin dynamics

1) Dynamics Before ADC, $0 \leq t \leq t_0$, the upstream dynamics

$$1) \quad u_t + E_0^{1/2} \lambda^{3/2} u u_x + \lambda^{3/2} u_{xxx} = 0, \quad u(x, t)|_{t=0} = u_0(x)$$

(2) u , 2π periodic, $M = \int_0^{2\pi} u dx = 0$, $E = \frac{1}{2} \int_0^{2\pi} u^2 dx = 1$ } These conserved quantities in time

3) HAMILTONIAN $H = E_0^{1/2} \lambda^{3/2} H_3 - \lambda^{3/2} H_2$ } Hamiltonian

4) $H_3 = \int_0^{2\pi} \frac{1}{6} u^3 dx$, $H_2 = \int_0^{2\pi} \frac{1}{2} u^2 dx$

(part > t_0)

(1) Dynamics After ADC at t_0 , the downstream dynamics

Note use 7/10/18 with 7/16/18 as rough guidelines

1) $\tilde{u}_+(x, t) = D_+^{1/4} u(x, t)$ where u solves B(1), with depth $0 < D_+ < 1$

2) $(\tilde{u}_+)_t + D_+^{-2/2} E_0^{1/2} \lambda^{3/2} \tilde{u}_+ \tilde{u}_{+x} + \lambda^{3/2} \tilde{u}_{+xxx} = 0$; ADC, lower energy
 stronger nonlinearity
 weaker dispersion

3) Set $u_+(x, t) \equiv D_+^{-1/4} \tilde{u}_+(x, t)$

$$p_H \rightarrow 2$$

$$3) a_+ = D_+^{\frac{1}{4}} u_+; \tilde{u}_+ + D_+^{\frac{1}{2}} \int_0^{\frac{1}{2}} \lambda^{\frac{3}{2}} a a_H + D_+^{\frac{3}{2}} \lambda^{\frac{3}{2}} \tilde{u}_{HH}$$

$$4) (a)_+ + D_+^{\frac{13}{4}} \int_0^{\frac{1}{2}} \lambda^{\frac{3}{2}} u_+(u)_H + D_+^{\frac{3}{2}} \lambda^{\frac{3}{2}} u_{,HHH} = 0$$

$$Mu_+(0) = 0, E(u_+) = 1 \text{ for } t_0 \leq t \leq +\infty$$

5) Hamiltonian, H_+ in outgoing state

$$H_+ = D_+^{\frac{13}{4}} \int_0^{\frac{1}{2}} \lambda^{\frac{3}{2}} H_3 = D_+^{\frac{3}{2}} \lambda^{\frac{3}{2}} H_2 \text{ for } t_0 \leq t \leq +\infty, \text{ Conserved}$$

H_+ more weight on 3rd moment, H_3 by factor $D_+^{\frac{13}{4}}$ } favors 3rd moment
 less weight on H_2 " " $D_+^{\frac{3}{2}}$ } after T_{ADC}
 $0 < D_+ < 1$, for

Statistical Behavior: Before ADC

nearly Gaussian, Model by $\theta \approx 0$, zero inv temp in Mixed Ensemble (FN of E_0)

Gives pdf of H_3, H_2 almost Explicitly at $t = T_{ADC}$ Supplies Stat Initial Data
 Note $H_3 + H_2$ comp separately for H_+ after ADC i.e. $t \geq T_{ADC}$

6) Matching Across T_{ADC}

$$u(x,t) \Big|_{t=T_{ADC}} = u(x,t) \Big|_{t=T_{ADC}}$$

B. Statistical TKDV Perspective

A) Use Micro-CAN-Energy - CAN HAMILTONIAN (AKM)

LA truncation level, 2 ADDF, S_n is 2dim unit sphere

H_n is HAMILTONIAN, Gibbs measure \mathcal{G}_0 is prob measure on S_n

$$1) \quad \mathcal{G}_0 = \frac{e^{-\theta H_n} dS_n}{\int_{S_n} e^{-\theta H_n} dS_n} \quad \begin{array}{l} \theta = 0, \text{ zero inv Temp} \\ \theta > 0, \text{ neg inv Temp} \end{array}$$

(can be \mathcal{G}_0 , inv of dynamics invariant dynamics)

B) Incoming Stat Dynamics: $-\infty < t \leq T$
AOC

From exp. Needs to have nearly Gaussian statistics, given E_0 - initial energy level obs.

2) From 1) B, use TKDV Hamiltonian from B(3)

$$H_n = E_0^{\frac{1}{2}} x^{\frac{3}{2}} H_3 - x^{\frac{3}{2}} H_2 \equiv H_3^\lambda - H_2^\lambda$$

3) Zero Inv Temp, with H_n in 2) is inv

i.e. $\mathcal{G}_0(H)$ is incoming inv measure

Two Key Prop of \mathcal{G}_0 as incoming invariant measure

9-1) For above, $\langle H_3^\lambda \rangle_{\mathcal{G}_0}$, $\langle H_2^\lambda \rangle_{\mathcal{G}_0}$ can be calc. by MC and Explicitly (B(5))

Prop: 4-2) for λ large or λ fixed and λ large, \mathcal{G}_0 is nearly Gaussian as reg in B(1)

See Fig 1 of B-F-1 and 7/16/18

1) Matching Outgoing Initial State Measures at $t = T_{ADC}$

From a) D-M-1 and II.B) above the initial value ρ^* at $T = T_{ADC}$ for outgoing incoming prob. meas, $\rho_+^*(u_+^*)$ is given by

$$1) \quad \rho_+(H) \equiv \rho_+^* \Big|_{t=T_{ADC}}$$

2) Outgoing Hamiltonian for $t > T_{ADC}$, Use 5) from DM-2

$$3) \quad H_+ = D_+^{-13} H_3^A - D_+^{3/2} H_2^A$$

3) From B4-1) above and (1-1) above, at $t = T_{ADC}$ we know

$$3) \quad \langle H_+^A \rangle_{\rho_+^*} \Big|_{t=T_{ADC}} = D_+^{13} \langle H_3^A \rangle_{\rho_0} + D_+^{3/2} \langle H_2^A \rangle_{\rho_0}$$

1) Transient and long-time Outgoing State Dynamics

1) Transient ~~low~~ Outgoing state dynamics

ρ_+^* for $t \geq T_{ADC}$ solves Liouville eq. t_{dyn} with IV in c) 1)

2) H_+ is conserved $\Rightarrow \langle H_+ \rangle_{\rho_+} = \langle H_+ \rangle_{\rho_0}$ for all $t \geq T_{ADC}$

3) Long-time Outgoing Dynamics with ergodicity converging to Outgoing Gibbs Measure

p) 3-1) $\rho_+(H_+)$ where ρ_+^* is Lagrange multiplier det. in

$$p) 3-2) \quad \langle H_+ \rangle_{\rho_0^+} = \langle H_+ \rangle_{\rho_{TADC}^+}$$

E. Projects for Outgoing Stat. Solns

- 1) Short-time PDF expansion and non-Gaussian behavior, pos skewness
for μ_{eff}
- 2) Non-Gaussian Stat. for outgoing Gibbs measure, $g(H^*)$
and stat. corr. of μ_i for P varying
- 3) Mean field behavior, central distribution, etc?

After Deterministic ADC, KdV eqn
KdV ADC Dynamics

1) Not E or L rescaled

$$u(x,0) = D_+^4 u(x,0)$$

See pg 2

$$u_t + D_+^3 u u_x + D_+^2 u_{xxx} = 0$$

u

2) E_0 and $L_0 = 1$, unit energy, 2π periodic, $M=0$

Outgoing TKdV Soln

$$\text{OTKdV)} \quad (u)_+ + D_+^3 E_0^{1/2} t^{-3/2} (\frac{1}{2} u^2)_+ + D_+^2 u_{xxx} = 0, \quad t > T_{ADC}$$

Incoming TKdV, $M \neq 0$, E_0 prescribed, $M=0$

B) $(u)_+ \equiv$ Same eqn. in A) with $D_+ \equiv 1$ $t_0 < t < T_{ADC}$, $t_0 \rightarrow -\infty$

3) Transition Matching Initial Conditions

$$\boxed{(u)_+ \Big|_{t=T_{ADC}} = D_+^4 u(x,t) \Big|_{t=T_{ADC}} \Leftrightarrow u \Big|_{t=T_{ADC}} = u \Big|_{t=T_{ADC}}}$$

Terminology TKdV \leftrightarrow Λ fixed

ADC - Abrupt Depth Change T_{ADC}

u or u - Incoming TKdV $t_0 < t < T_{ADC}$

u - Outgoing TKdV $T_{ADC} \leq t < \infty$

Detailed Steps:

Majda P1010

ADC-KdV rescaled Rascally

1) $\tau = \frac{1}{\epsilon} x - t, \quad x = \epsilon \tau = \epsilon x$

2) Before ADC, $\tau = x - t$

3) After ADC, $\tau(x) = \int_0^x D^{-1/2} x \quad \text{so} \quad \tau = D^{-1/2} x - t$

Set $\tau = D^{-1/2} x \quad \tau = D^{-1/2} (x - t), \quad x = \epsilon x$

4) pg. 272, Johnson, $H_0 = D^{-1/2} \eta_0$

Original
Derivatives

$2U_0 x + \frac{3}{D^4} H_0 H_0 + \frac{1}{5} D^2 H_0 = 0 \quad \tau \text{ derivatives here}$

5)

Change variables: $\tau = D^{-1/2} x$

$\frac{\partial}{\partial \tau} = D^{1/2} \frac{\partial}{\partial x} \Leftrightarrow \frac{d}{d\tau} = D^{1/2} \frac{d}{dx}$

$2H_0(x) + 3 \frac{D^{-1/2}}{D^4} H_0 H_0 + \frac{1}{5} \frac{D^2}{D^2} H_0 = 0$

$x \cdot \tau = x - \tau \tau, \quad x = \epsilon x, \quad \frac{\partial}{\partial \tau} = D^{1/2} \frac{\partial}{\partial x} \text{ and } \frac{d}{d\tau} = \frac{dx}{d\tau} \frac{d}{dx} = D^{1/2} \frac{d}{dx}$

rescaled KdV

6) $2(H_0)_\tau + 3 \frac{D^{-1/2}}{D^4} H_0 H_0 + \frac{1}{5} \frac{D^2}{D^2} H_0 = 0 \quad \text{Tone Rescaled KdV}$

7) $H_0 = u, \quad \frac{3}{2} u^2, \quad \text{Simple Soling}$

Downstream after T_{ADC}

Note line Rescaling - X rescaling D_+ factor

- 0) $t < T_{ADC}$, zero temp steady state, E_0, λ , Nearly Gaussian, IV
 At $t = T_{ADC}$ Gaussian pdf for $\Lambda_3(P_3, E_3, \lambda) H_3 = N(0, \sigma_0^2)$ for $\sigma_0 \ll 1$ } Mean Field II
 " " for $\Lambda_2(P_2, E_2, \lambda) H_2 = N(0, \sigma_0^2)$

2) $F \rightarrow \infty$ + ergodic Final stat predicted by TKN

3) Question: Short term stat trans state

$\langle H(P_2) \rangle$ cons

SAM - game as practice notes

Link 2) + 3)

4) If Mean Field Regime still applies

Model 2) pdf by centered I-list as in MW book

The ISBS Small Spatial Period Rescaling

$$1) \quad u_t + \frac{1}{2} u_x^2 + u_{xxx} = 0$$

$$M=0, E, H = \frac{1}{6} \left(u^3 - \frac{1}{2} u_x^2 \right) \equiv H_3 - H_2 \quad \text{cubic + Quad Parts}$$

$2\pi L$ periodic in x , L in expt

$$2) \quad \tilde{u}(x,t) \quad \left[\tilde{u} = \frac{u(x,t) \lambda^{1/2}}{E^{1/2}} \right] \quad \int_0^{2\pi} \frac{1}{2} \tilde{u}^2 = \frac{1}{2} \int_0^{2\pi} \frac{u^2(x,t) \lambda^{1/2}}{E^{1/2}} dx = 1$$

\tilde{u} Unit energy, $2\pi L \lambda$ periodic

$$3) \text{ Dynamics for } \tilde{u}, \quad u(x,t) = \frac{E^{1/2}}{\lambda} \tilde{u} = \left(\frac{E}{\lambda} \right)^{1/2} \tilde{u}$$

u_t

$$(\lambda \rightarrow L) \quad X' = L \quad 2\pi L \text{ initial period } L \text{ small, narrow channel}$$

Rescaled

$$\begin{aligned} \tilde{u} &= 0 \text{ at } \partial \Omega, t \\ \tilde{x} = \lambda x \quad u_t = u_x, \quad \frac{\partial}{\partial x} &= \frac{\partial}{\partial \tilde{x}} \\ \tilde{u}(\tilde{x}, t) &= 0 \text{ at } \partial \Omega, t \\ u_t + \frac{1}{2} u_x^2 + u_{xxx} &= 0 \\ \tilde{u}_t + (\lambda)^{-1} \tilde{u} \tilde{u}_{\tilde{x}} + \lambda^3 \tilde{u}_{\tilde{x}\tilde{x}\tilde{x}} &= 0 \end{aligned}$$

$$4) \quad H = \frac{1}{6} E \lambda^{-3/2} H_3 - \lambda^{-3} H_2, \quad \tilde{u}(\tilde{x}, t)$$

$$5) \text{ Zero Inverse Temp } G_0 = \rho^{\text{OH}} dL_n, \quad \text{zero inv temp } G_0 = \frac{dL_n}{\text{Area}(L_n)} \quad \text{indep. of } H$$

No detailed dynamics

$$A) \quad \left[\frac{1}{2} \lambda^{-3/2} \right] \times [H_3] \quad \text{zero mean "GAUSSIAN" for } L \text{ large enough } L \rightarrow 0$$

$$B) \quad \lambda^{-3} [H_2] \quad \text{neg. mean "GAUSSIAN"} \quad \left[\text{See Fig 1-B-F-L, } p=0 \right]$$

$$C) \text{ Regime, } \lambda \text{ large } \Leftrightarrow L \text{ small pdf is sum of two GAUSSIAN RV} \\ A) \text{ zero mean, Var } \approx E \lambda^{-3} \\ B) \text{ mean neg } = \lambda^{-3} \text{ and } V = \lambda^{-6}$$


$$D) \quad \text{pdf}(A+B) = \text{pdf}(A) * \text{pdf}(B) \quad \rightarrow \text{Sharply peaked GAUSSIAN with slightly neg. mean}$$

SUMMARY: Zero Temp, Gaussian ISBS

E) $D = \text{pdf } H \text{ in } \psi$ for rescaled Hamiltonian in (4)

$\langle H \rangle = -\epsilon A^3$, $\text{var}(EA^3)$, nearly Gaussian, small mean

6)



Small V_{IR} , zero mean

Very close SB-I with $\langle H \rangle \leq 0 \leftrightarrow$ zero temp Motivated SB from 6/28/18

Look at in scattered state \Rightarrow neg temp regime