

The line integral of \vec{F} along C is

$$\int_C \vec{F} \cdot d\vec{r} = \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}' dt$$

$$\vec{F}(\vec{r}(t)) = \vec{F}(x(t), y(t), z(t))$$

$$\int_C \vec{F} \cdot d\vec{r} = \int_C \vec{F} \cdot \vec{T} ds \quad \vec{T}(t) = \frac{\vec{r}'}{\|\vec{r}'\|}$$

$$\begin{aligned} \int_C \vec{F} \cdot d\vec{r} &= \int_C \vec{F} \cdot \vec{T} ds \\ &= \int_a^b \vec{F}(\vec{r}(t)) \cdot \frac{\vec{r}'(t)}{\|\vec{r}'(t)\|} \|\vec{r}'(t)\| dt \\ &= \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt \end{aligned}$$

Example 1

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y, z) = 8x^2yz\vec{i} + 5z\vec{j} - 4xy\vec{k}$ and C is the curve given by $\vec{r}(t) = t\vec{i} + t^2\vec{j} + t^3\vec{k}$, $0 \leq t \leq 1$.

$$\vec{F}(\vec{r}(t)) = 8t^2(t^2)(t^3)\vec{i} + 5t^3\vec{j} - 4t(t^2)\vec{k} = 8t^7\vec{i} + 5t^3\vec{j} - 4t^3\vec{k}$$

$$\vec{r}'(t) = \vec{i} + 2t\vec{j} + 3t^2\vec{k}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = (8t^7 + 5t^3 - 4t^3) \cdot (1 + 2t + 3t^2) = 8t^7 + 10t^4 - 12t^5$$

$$= \int_0^1 (8t^7 + 10t^4 - 12t^5) dt = [t^8 + 2t^5 - 2t^6]_0^1 = 1$$

Example 2

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y, z) = xz\vec{i} - yz\vec{k}$ and C is the line segment from $(-1, 2, 0)$ to $(3, 0, 1)$.

$$\vec{r}(t) = (1-t)\langle -1, 2, 0 \rangle + t\langle 3, 0, 1 \rangle = \langle 4t-1, -2t+2, t \rangle$$

$$\vec{r}'(t) = \langle 4, -2, 1 \rangle$$

$$\vec{F}(\vec{r}(t)) = (4t-1)(t)\vec{i} - (-2t+2)(t)\vec{k} = (4t^2-t)\vec{i} - (-2t^2+2t)\vec{k}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = 4(4t^2 - t) - (-2t^2 + 2t) = 18t^2 - 6t$$

$$= \int_0^1 18t^2 - 6t \, dt = [6t^3 - 3t^2]_0^1 = 3$$

Given the vector field $\vec{F}(x, y, z) = P\vec{i} + Q\vec{j} + R\vec{k}$ and the curve C parameterized by $\vec{r} = x(t)\vec{i} + y(t)\vec{j} + z(t)\vec{k}$, $a \leq t \leq b$

$$\begin{aligned} \int_C \vec{F} \cdot d\vec{r} &= \int_a^b \left(P\vec{i} + Q\vec{j} + R\vec{k} \right) \cdot \left(x'\vec{i} + y'\vec{j} + z'\vec{k} \right) dt \\ &= \int_a^b Px' + Qy' + Rz' \, dt \\ &= \int_a^b Px' \, dt + \int_a^b Qy' \, dt + \int_a^b Rz' \, dt \\ &= \int_C P \, dx + \int_C Q \, dy + \int_C R \, dz \\ &= \int_C P \, dx + Q \, dy + R \, dz \\ &= \int_{-C} \vec{F} \cdot d\vec{r} = - \int_C \vec{F} \cdot d\vec{r} \end{aligned}$$

1.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = y^2\vec{i} + (3x - 6y)\vec{j}$ and C is the line segment from $(3, 7)$ to $(0, 12)$.

$$\vec{r}(t) = (1 - t)\langle 3, 7 \rangle + t\langle 0, 12 \rangle = \langle 3 - 3t, 7 + 5t \rangle$$

$$\vec{r}'(t) = \langle -3, 5 \rangle$$

$$\vec{F}(\vec{r}(t)) = (7 + 5t)^2\vec{i} + (3(3 - 3t) - 6(7 + 5t))\vec{j} = (7 + 5t)^2\vec{i} + (-33 - 39t)\vec{j}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = -3(7 + 5t)^2 - 5(33 + 39t)$$

$$\begin{aligned} &= \int_0^1 -3(7 + 5t)^2 - 5(33 + 39t) \, dt = \left[-\frac{(7 + 5t)^3}{5} - 165t - \frac{195t^2}{2} \right]_0^1 \\ &= -\frac{12}{5} - 165 - \frac{195}{2} - \left(-\frac{1}{5}7^3 \right) = -\frac{1079}{2} \end{aligned}$$

2.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = (x + y)\vec{i} + (1 - x)\vec{j}$ and C is the portion of $\frac{x^2}{4} + \frac{y^2}{9} = 1$ that is in the 4th quadrant with the counter clockwise rotation.

$$\vec{r}(t) = \langle 2 \cos t, 3 \sin t \rangle$$

$$\vec{r}'(t) = \langle -2 \sin t, 3 \cos t \rangle$$

$$\vec{F}(\vec{r}(t)) = (2 \cos t + 3 \sin t)\vec{i} + (1 - 2 \cos t)\vec{j}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = (2 \cos t + 3 \sin t)(-2 \sin t) + (1 - 2 \cos t)(3 \cos t) = -2 \sin 2t + 3 \cos t - 6$$

$$= \int_{\frac{3\pi}{2}}^{2\pi} -2 \sin 2t + 3 \cos t - 6 \, dt = [\cos 2t + \sin t - 6t]_{\frac{3\pi}{2}}^{2\pi} = 5 - 3\pi$$

3.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = y^2\vec{i} + (x^2 - 4)\vec{j}$ and C is the portion of $y = (x - 1)^2$ from $x = 0$ to $x = 3$.

$$\vec{r}(t) = \langle t, (t - 1)^2 \rangle$$

$$\vec{r}'(t) = \langle 1, 2(t - 1) \rangle$$

$$\vec{F}(\vec{r}(t)) = (t - 1)^4\vec{i} + (t^2 - 4)\vec{j}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = (t - 1)^4 + 2(t - 1)(t^2 - 4) = (t - 1)^4 + 2t^3 - 2t^2 - 8t + 8$$

$$= \int_0^3 (t - 1)^4 + 2(t - 1)(t^2 - 4) \, dt = \left[\frac{1}{5}(t - 1)^5 + \frac{1}{2}t^4 - \frac{2}{3}t^3 - 4t^2 + 8t \right]_0^3 = \frac{171}{10}$$

4.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y, z) = e^{2x}\vec{i} + z(y+1)\vec{j} + z^3\vec{k}$ and C is given by $\vec{r}(t) = t^3\vec{i} + (1-3t)\vec{j} + e^t\vec{k}$ for $0 \leq t \leq 2$.

$$\vec{r}'(t) = 3t^2\vec{i} - 3\vec{j} + e^t\vec{k}$$


$$\vec{F}(\vec{r}(t)) = e^{2t^3}\vec{i} + e^t(2-3t)\vec{j} + e^{3t}\vec{k}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = e^{2t^3}(3t^2) + e^t(2-3t)(-3) + e^{3t}(e^t) = 3t^2e^{2t^3} - 3e^t(2-3t) + e^{4t}$$

$$= \int_0^2 3t^2e^{2t^3} - 3e^t(2-3t) + e^{4t} dt = \left[\frac{1}{2}e^{2t^3} - 3e^t(5-3t) + \frac{1}{4}e^{4t} \right]_0^2 = \frac{57}{4} + 3e^2 + \frac{1}{4}e^8 + \frac{1}{2}e^{16}$$

5.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = 3y\vec{i} + (x^2 - y)\vec{j}$ and C is the upper half of the circle centered at the origin of radius 1 with counter clockwise rotation and the portion of $y = x^2 - 1$ from $x = -1$ to $x = 1$. See the example below.



$$C_1 : \vec{r}(t) = \langle \cos t, \sin t \rangle \quad 0 \leq t \leq \pi$$

$$C_1 : \vec{r}'(t) = \langle -\sin t, \cos t \rangle$$

$$C_1 : \vec{F}(\vec{r}(t)) = 3 \sin t \vec{i} + (\cos^2 t - \sin t) \vec{j}$$

$$C_1 : \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = -3 \sin^2 t + \cos^3 t - \cos t \sin t$$

$$= -\frac{3}{2}(1 - \cos 2t) + \cos t(1 - \sin^2 t) - \cos t \sin t$$

$$\begin{aligned}
&= -\frac{3}{2}(1 - \cos 2t) + \cos t(1 - \sin^2 t) - \frac{1}{2} \sin 2t \\
&= \int_0^\pi -\frac{3}{2}(1 - \cos 2t) + \cos t - \cos t \sin^2 t - \frac{1}{2} \sin 2t \, dt \\
&= \left[-\frac{3}{2}\left(t - \frac{1}{2} \sin 2t\right) + \sin t - \frac{1}{3} \sin^3 t + \frac{1}{4} \cos 2t \right]_0^\pi = -\frac{3\pi}{2}
\end{aligned}$$

$$\vec{r}'(t) = 3t^2 \vec{i} - 3\vec{j} + e^t \vec{k}$$

$$\vec{F}(\vec{r}(t)) = e^{2t^3} \vec{i} + e^t(2 - 3t) \vec{j} + e^{3t} \vec{k}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = e^{2t^3}(3t^2) + e^t(2 - 3t)(-3) + e^{3t}(e^t) = 3t^2 e^{2t^3} - 3e^t(2 - 3t) + e^{4t}$$

$$= \int_0^2 3t^2 e^{2t^3} - 3e^t(2 - 3t) + e^{4t} \, dt = \left[\frac{1}{2} e^{2t^3} - 3e^t(5 - 3t) + \frac{1}{4} e^{4t} \right]_0^2 = \frac{57}{4} + 3e^2 + \frac{1}{4} e^8 + \frac{1}{2} e^{16}$$

$$C_2 : \vec{r}(t) = \langle t, t^2 - 1 \rangle \quad -1 \leq t \leq 1$$

$$C_2 : \vec{r}'(t) = \langle 1, 2t \rangle$$

$$C_2 : \vec{F}(\vec{r}(t)) = 3(t^2 - 1) \vec{i} + \vec{j}$$

$$C_2 : \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = 3(t^2 - 1) + 2t$$

$$= \int_{-1}^1 3(t^2 - 1) + 2t \, dt = [t^3 - 3t + t^2]_{-1}^1 = -4$$

6.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = (x + y)\vec{i} + (1 - x)\vec{j}$ and C is the portion of $\frac{x^2}{4} + \frac{y^2}{9} = 1$ that is in the 4th quadrant with the counter clockwise rotation.

$$\vec{r}(t) = \langle t, (t - 1)^2 \rangle$$

$$\vec{r}'(t) = \langle 1, 2(t - 1) \rangle$$

$$\vec{F}(\vec{r}(t)) = (t-1)^4 \vec{i} + (t^2-4) \vec{j}$$

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = (t-1)^4 + 2(t-1)(t^2-4) = (t-1)^4 + 2t^3 - 2t^2 - 8t + 8$$

$$= \int_0^3 (t-1)^4 + 2(t-1)(t^2-4) \, dt = \left[\frac{1}{5}(t-1)^5 + \frac{1}{2}t^4 - \frac{2}{3}t^3 - 4t^2 + 8t \right]_0^3 = \frac{171}{10}$$

7.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = (6x - 2y)\vec{i} + x^2\vec{j}$ for each of the following curves.

a)

C is the line segment from $(6, -3)$ to $(0, 0)$ followed by the line segment from $(0, 0)$ to $(6, 3)$.

b)

C is the line segment from $(6, -3)$ to $(6, 3)$.

8.

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = 3\vec{i} + (xy - 2x)\vec{j}$ for each of the following curves.

a)

C is the upper half of the circle centered at the origin of radius 4 with counter clockwise rotation.

b)

C is the upper half of the circle centered at the origin of radius 4 with clockwise rotation. Evaluate \int_C

$\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F}(x, y) = (x + y)\vec{i} + (1 - x)\vec{j}$ and C is the portion of $x^2 + y^2 = 1$ that is in the 4th quadrant with the counter clockwise rotation. \int_C

$\int_C \vec{F} \cdot d\vec{r} = \int_C$

$\int_C \vec{F} \cdot d\vec{r} = \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) \, dt$ where $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j} + z(t)\vec{k}$, $a \leq t \leq b$ the line integral is,

\int_C

$\rightarrow F = P \rightarrow i + Q \rightarrow j + R \rightarrow k$ and the curve C parameterized by $\rightarrow r(t) = x(t) \rightarrow i + y(t) \rightarrow j + z(t) \rightarrow k$, the line integral is,

$$\int_C \rightarrow F \cdot d\rightarrow r = \int_C (P \rightarrow i + Q \rightarrow j + R \rightarrow k) \cdot (x'(t) \rightarrow i + y'(t) \rightarrow j + z'(t) \rightarrow k) dt$$

$$= \int_C P x'(t) dt + \int_C Q y'(t) dt + \int_C R z'(t) dt =$$

$$\int_C P dx + \int_C Q dy + \int_C R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$= \int_C P dx + Q dy + R dz$$

$$f(x) = \sqrt{1 + \sqrt{x}}$$

$$f'(x) = (1 + (x)^{\frac{1}{2}})^{\frac{1}{2}}$$

$$= \frac{1}{2\sqrt{1 + \sqrt{x}}} \cdot \frac{1}{2\sqrt{x}} = \frac{1}{4\sqrt{x}\sqrt{1 + \sqrt{x}}}$$

$$F(x) = \int_0^{\sin x} \frac{t^2}{t^2 - 1} dt$$

$$F'(x) = \frac{\sin^2 x}{\sin^2 x - 1} \cdot \frac{d}{dx} \sin x$$

$$= -\frac{\sin^2 x}{1 - \sin^2 x} \cdot \cos x$$

$$= -\frac{\sin^2 x}{\cos^2 x} \cdot \cos x$$

$$= -\frac{\sin^2 x}{\cos x}$$

$$y - y_0 = m(x - x_0)$$

$$m = \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{3t^2}{2t} = \frac{3t}{2}$$

$$m(t = 5) = \frac{15}{2}$$

$$x(5) = 25 \quad y(5) = 125$$

$$y - 125 = \frac{15}{2}(x - 25) = y = 125 + \frac{15}{2}(x - 25)$$

4. (2 valores) Calcule uma (apenas uma) das seguintes primitivas

$$\int \frac{x^3}{1+x^8} dx \quad \int (x+2)e^x dx$$

$$\int \frac{x^3}{1+x^8} dx = \frac{1}{4} \int \frac{4x^3}{1+(x^4)^2} dx = \frac{1}{4} \arctan x^4$$

$$\int = (x+2)e^x dx = \int xe^x + 2e^x dx$$

$$= \int xe^x dx + \int 2e^x dx$$

$$\int xe^x dx = xe^x - e^x$$

$$= \int xe^x dx + \int 2e^x dx = (x+1)e^x$$

5. (2 valores) Calcule um (apenas um) dos seguintes integrais

$$\int_{-\pi/2}^{\pi/2} \cos^2 \theta \sin \theta d\theta \quad \int_1^3 \log(x^3) dx$$

$$\int_{-\pi/2}^{\pi/2} \cos^2 \theta \sin \theta d\theta = \frac{1}{3} [-\cos^3 \theta]_{-\pi/2}^{\pi/2} = \frac{4x^3}{1+(x^4)^2} dx = 0$$

$$\int_1^3 \ln x^3 dx = 3 \int_1^3 \ln x dx$$

$$f = \ln x \quad g' = 1$$

$$f' = \frac{1}{x} \quad g = x$$

$$3 \int_1^3 \ln x \, dx = [x \ln x]_1^3 - \int_1^3 1 = 3x \ln x - 3x = 9 \ln 3 - 6$$

6. (2 valores) Determine a solução da equação diferencial $\frac{dx}{dt} = 2 - x$ com condição inicial $x(0) = 1$.

$$\dot{y} = 2 - x \quad x(0) = 1$$

$$Y_c = c_1 e^{-t}$$

$$2 - x = \Leftrightarrow x = 2$$

$$c_1 e^0 + 2 = 3 \Leftrightarrow c_1 = 1$$

$$Y_c = e^{-t} + 2$$

8. (2 valores) Calcule o limite

$$\lim_{x \rightarrow \pi} \frac{1 + \cos x}{x - \pi}$$

Método 1:

$$f(x) = \cos x$$

$$f'(\pi) = \frac{f(x) - f(\pi)}{x - \pi}$$

$$f'(\pi) = -\sin \pi = 0$$

Método 2:

$$\lim_{x \rightarrow \pi} = \frac{1 + \cos x}{x - \pi} = \frac{\frac{d}{dx}(1 + \cos x)}{\frac{d}{dx}(x - \pi)} = 0$$

$$\ddot{y} + 9y = \sin(\pi t)$$

$$y_c(t) = c_1 \cos(3t) + c_2 \sin(3t)$$

$$z = b \sin(\pi t)$$

$$z' = \pi b \cos(\pi t)$$

$$z'' = -\pi^2 b \sin(\pi t)$$

$$-\pi^2 b \sin(\pi t) + 9bt \sin(\pi t) = \sin(\pi t)$$

$$\cancel{b \sin(\pi t)}(-\pi^2 + 9) = \cancel{\sin(\pi t)}$$

$$b = \frac{1}{-\pi^2 + 9}$$

$$y_c(t) + y_p(t) = c_1 \cos(3t) + c_2 \sin(3t) + \frac{1}{-\pi^2 + 9} \sin(\pi t)$$

$$\ddot{y} + 9y = \sin(3t)$$

Método 1:

$$y_c(t) = c_1 \cos(3t) + c_2 \sin(3t)$$

$$z = bt \cos(3t)$$

$$z' = -3bt \sin(3t) + b \cos(3t)$$

$$z'' = -6b \sin(3t) - 9bt \cos(3t)$$

$$-6b \sin(3t) - 9bt \cos(3t) + 9bt \cos(3t) = \sin(3t)$$

$$-6b \sin(3t) = \sin(3t)$$

$$b = -\frac{1}{6}$$

$$y_c(t) + y_p(t) = c_1 \cos(3t) + c_2 \sin(3t) - \frac{1}{6} t \cos(3t)$$

Método 2:

$$y_c(t) = c_1 \cos(3t) + c_2 \sin(3t)$$

$$y_1(t) = \cos(3t) \quad y_2(t) = \sin(3t)$$

$$W = \begin{vmatrix} \cos(3t) & \sin(3t) \\ -3\sin(3t) & 3\cos(3t) \end{vmatrix} = 3$$

$$Y_p(t) = -y_1 \int \frac{y_2 g(t)}{W(y_1, y_2)} dt + y_2 \int \frac{y_1 g(t)}{W(y_1, y_2)} dt$$

$$Y_p(t) = -\frac{\cos(3t)}{3} \int \sin(3t) \sin(3t) dt + \frac{\sin(3t)}{3} \int \cos(3t) \sin(3t) dt$$

$$\sin^2(3t) = \frac{1 - \cos(6t)}{2}$$

$$Y_p(t) = -\frac{\cos(3t)}{3} \int \sin^2(3t) dt + \frac{\sin(3t)}{3} \cdot -\frac{\sin^2(3t)}{6}$$

$$Y_p(t) = -\frac{\cos(3t)}{3} \int \frac{1}{2} - \frac{1}{2} \cos(6t) dt + \frac{\sin(3t)}{3} \left(-\frac{1}{12} + \frac{1}{12} \cos(6t) \right)$$