

Exercise 8, Discrete Mathematics for Bioinformatics

Sascha Meiers, Martin Seeger

Winter term 2011/2012

8.1 Linear Optimization

$$2x_1 + 3x_2 = \min \Leftrightarrow -2x_1 - 3x_2 = \max. \quad (1)$$

$$3x_1 + 6x_2 \leq 7 \Leftrightarrow 3x_1 + 6x_2 + x_3 = 7, \quad x_3 \geq 0. \quad (2)$$

$$x_1 \text{ free} \Leftrightarrow x_1 = x_4 - x_5, \quad x_4 \geq 0, \quad x_5 \geq 0.$$

Insert the previous equation into (2):

$$6x_2 + x_3 + 3x_4 - 3x_5 = 7, \quad x_3 \geq 0, \quad x_4 \geq 0, \quad x_5 \geq 0.$$

Finally,

$$2x_1 + 2x_2 = 5 \Leftrightarrow 2x_2 + 2x_4 - 2x_5 = 5,$$


$$-2x_1 - 3x_2 = -3x_2 - 2x_4 + 2x_5 = \max.$$

Summarizing,

$$\max \left\{ (-3, 0, -2, 2) \begin{pmatrix} x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} \mid \begin{pmatrix} 6 & 1 & 3 & -3 \\ 2 & 0 & 2 & -2 \end{pmatrix} \begin{pmatrix} x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}, \begin{pmatrix} x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} \geq 0 \right\}.$$

8.2 Linear Optimization

The feasible region is shown in the inequality plot below. The red arrow is the vector of the first optimization function, the blue arrow the one belonging to the second one. Both optimization goals are maximized in the vertex labeled with the check mark.



8.3 Linear Optimization

- a) For $c^T = (-1, -1)$ we have to maximize $-x_1 - x_2$, i.e. minimize $x_1 + x_2$.
In the feasible region, this is the case for $(x_1, x_2) = (0, 0)$.
- b) For $c^T = (0, -1)$ we have to maximize $-x_2$, i.e. minimize x_2 .
In the feasible region, this is the case for $(x_1, x_2) = (\lambda, 0)$, $\lambda \in [0, 1]$.
- c) For $c^T = (-1, 0)$ we have to maximize $-x_1$, i.e. minimize x_1 .
In the feasible region, this is the case for $(x_1, x_2) = (0, \lambda)$, $\lambda \in [0, \infty[$.
- d) For $c^T = (1, 1)$ we have to maximize $x_1 + x_2$.
Since e.g. the half-line $x_1 - x_2 = 0$, $x_1, x_2 \geq 0$ lies in the feasible region, and as the target function increases strictly along this half-line, there is no optimal solution.

If the following constraint is added, the problem becomes infeasible:

$$x_1 - x_2 \geq 2.$$

8.4 Profit Optimization

The linear program contains six inequalities:

$$\max\left\{\begin{pmatrix} 20 & 25 \end{pmatrix} \cdot \begin{pmatrix} A \\ B \end{pmatrix} \mid \begin{pmatrix} 1 & 0 \\ -1 & 0 \\ 0 & 1 \\ 0 & -1 \\ 1 & -2 \\ 1 & 2 \end{pmatrix} \cdot \begin{pmatrix} A \\ B \end{pmatrix} \leq \begin{pmatrix} 60 \\ 0 \\ 50 \\ 0 \\ 0 \\ 120 \end{pmatrix}\right\}$$

Again, we solve it graphically, leading to the optimal solution of \$1950 profit per day. It is reached with a daily production of $A = 60$ and $B = 30$ refrigerator units.

