



Math for the people, by the people.

## Bautin's theorem

Canonical name	BautinsTheorem
Date of creation	2013-03-22 14:28:46
Last modified on	2013-03-22 14:28:46
Owner	Daume (40)
Last modified by	Daume (40)
Numerical id	5
Author	Daume (40)
Entry type	Theorem
Classification	msc 34C07

There are at most three limit cycles which can appear in the following quadratic system

$$\begin{aligned}\dot{x} = p(x, y) &= \sum_{i+j=0}^2 a_{ij} x^i y^j \\ \dot{y} = q(x, y) &= \sum_{i+j=0}^2 b_{ij} x^i y^j\end{aligned}$$

from a singular point, if its type is either a focus or a center.

In 1939 N.N. Bautin claimed the above result and in 1952 submitted the proof [?]. [?]

## References

- [GAV] Gaiko, A., Valery: Global Bifurcation Theory and Hilbert's Sixteenth Problem. Kluwer Academic Publishers, London, 2003.
- [BNN1] Bautin, N.N.: On the number of limit cycles appearing from an equilibrium point of the focus or center type under varying coefficients. Matem. SB., 30:181-196, 1952. (written in Russian)
- [BNN2] Bautin, N.N.: On the number of limit cycles appearing from an equilibrium point of the focus or center type under varying coefficients. Translation of the American Mathematical Society, 100, 1954.