



planetmath.org

Math for the people, by the people.

inverse Gudermannian function

Canonical name	InverseGudermannianFunction
Date of creation	2013-03-22 19:06:28
Last modified on	2013-03-22 19:06:28
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	5
Author	pahio (2872)
Entry type	Definition
Classification	msc 33B10
Classification	msc 26E05
Classification	msc 26A48
Classification	msc 26A09
Synonym	inverse Gudermannian
Related topic	HyperbolicFunctions
Related topic	AreaFunctions
Related topic	MercatorProjection
Related topic	EulerNumbers2
Related topic	DualityOfGudermannianAndItsInverseFunction

Since the real Gudermannian function gd is strictly increasing and forms a bijection from \mathbb{R} onto the open interval $(-\frac{\pi}{2}, \frac{\pi}{2})$, it has an inverse function

$$\text{gd}^{-1}: (-\frac{\pi}{2}, \frac{\pi}{2}) \rightarrow \mathbb{R}.$$

The function gd^{-1} is denoted also **aragd**.

If $x = \text{gd } y$, which may be explicitly written e.g.

$$x = \arcsin(\tanh y),$$

one can solve this for y , getting first $\tanh y = \sin x$ and then

$$y = \text{artanh}(\sin x)$$

(see the area functions). Hence the *inverse Gudermannian* is expressed as

$$\text{gd}^{-1}(x) = \text{aragd } x = \text{artanh}(\sin x) \quad (1)$$

It has other <http://planetmath.org/Equivalent3equivalent> expressions, such as

$$\text{gd}^{-1}(x) = \text{arsinh}(\tan x) = \frac{1}{2} \ln \frac{1 + \sin x}{1 - \sin x} = \int_0^x \frac{dt}{\cos t}. \quad (2)$$

Thus its derivative is

$$\frac{d}{dx} \text{gd}^{-1}(x) = \frac{1}{\cos x}. \quad (3)$$

Cf. the formulae (1)–(3) with the corresponding ones of gd .