

planetmath.org

Math for the people, by the people.

generating function of Legendre polynomials

Canonical name GeneratingFunctionOfLegendrePolynomials

Date of creation 2015-03-08 20:29:06 Last modified on 2015-03-08 20:29:06

Owner pahio (2872) Last modified by pahio (2872)

Numerical id 17

Author pahio (2872)

Entry type Result
Classification msc 33B99
Classification msc 30D10
Classification msc 30B10

Related topic GeneratingFunctionOfHermitePolynomials
Related topic GeneratingFunctionOfLaguerrePolynomials

Related topic VariantOfCauchyIntegralFormula

For finding the generating function

$$F(t) = \sum_{n=0}^{\infty} P_n(z)t^n$$

of the sequence of the Legendre polynomials

$$P_0(z) = 1$$

$$P_1(z) = z$$

$$P_2(z) = \frac{1}{2}(3z^2 - 1)$$

$$P_3(x) = \frac{1}{2}(5z^3 - 3z)$$

$$P_4(z) = \frac{1}{8}(35z^4 - 30z^2 + 3)$$

$$P_5(z) = \frac{1}{8}(63z^5 - 70z^3 + 15z)$$

we have to present $P_n(z)$ as the general coefficient of Taylor series in t, i.e. as the nth derivative of some F(t) in the origin, divided by the factorial n!. The Cauchy integral formula offers the chance to implement that.

Starting from the http://planetmath.org/node/11983Rodrigues formula of Legendre polynomials, we may write

$$P_n(z) = \frac{1}{2^n n!} \frac{d^n}{dz^n} (z^2 - 1)^n = \frac{1}{2^n n!} \frac{n!}{2i\pi} \oint_c \frac{(\zeta^2 - 1)^n}{(\zeta - z)^{n+1}} d\zeta = \frac{1}{2i\pi} \oint_c \left(\frac{1}{2} \frac{\zeta^2 - 1}{\zeta - z}\right)^n \frac{d\zeta}{\zeta - z},$$

where the contour c runs anticlockwise once around the point z. The change of variable

$$\frac{\zeta^2 - 1}{2(\zeta - z)} = \frac{1}{t}, \qquad d\zeta = \frac{zt - 1 - \sqrt{1 - zt + t^2}}{t^2 \sqrt{1 - zt + t^2}} dt$$

gives

$$P_n(z) = -\frac{1}{2i\pi} \oint_{c'} \frac{dt}{t^n t \sqrt{1 - zt + t^2}}$$

where t must go round the origin clockwise, but in

$$P_n(z) = \frac{1}{n!} \cdot \frac{n!}{2i\pi} \oint_{c'} \frac{dt}{\sqrt{1 - zt + t^2} \cdot (t - 0)^{n+1}}$$

anticlockwise. This is, by Cauchy integral formula again,

$$P_n(z) = \frac{1}{n!} \left[\frac{d^n}{dt^n} \frac{1}{\sqrt{1 - zt + t^2}} \right]_{t=0}.$$

This means that

$$F(t) := \frac{1}{\sqrt{1-zt+t^2}}$$

is the searched generating function of the Legendre polynomials:

$$\frac{1}{\sqrt{1-zt+t^2}} = P_0(z) + P_1(z)t + P_2(z)t^2 + P_3(z)t^3 + \dots$$

Cf. the generating function of the Bessel functions.