

Jacobi's identity for ϑ functions

 ${\bf Canonical\ name} \quad {\bf Jacobis Identity For vartheta Functions}$

Date of creation 2013-03-22 14:46:45 Last modified on 2013-03-22 14:46:45

Owner rspuzio (6075) Last modified by rspuzio (6075)

Numerical id 5

Author rspuzio (6075) Entry type Theorem Classification msc 33E05 Jacobi's identities describe how theta functions transform under replacing the period with the negative of its reciprocal. Together with the quasiperiodicity relations, they describe the transformations of theta functions under the modular group.

$$\theta_{1}(z \mid -1/\tau) = -i(-i\tau)^{1/2} e^{\frac{i\tau z^{2}}{\pi}} \theta_{1}(\tau z \mid \tau)$$

$$\theta_{2}(z \mid -1/\tau) = (-i\tau)^{1/2} e^{\frac{i\tau z^{2}}{\pi}} \theta_{4}(\tau z \mid \tau)$$

$$\theta_{3}(z \mid -1/\tau) = (-i\tau)^{1/2} e^{\frac{i\tau z^{2}}{\pi}} \theta_{3}(\tau z \mid \tau)$$

$$\theta_{4}(z \mid -1/\tau) = (-i\tau)^{1/2} e^{\frac{i\tau z^{2}}{\pi}} \theta_{2}(\tau z \mid \tau)$$