



## orthogonality of Legendre polynomials

Canonical name	OrthogonalityOfLegendrePolynomials
Date of creation	2013-03-22 18:55:30
Last modified on	2013-03-22 18:55:30
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	14
Author	pahio (2872)
Entry type	Derivation
Classification	msc 33C45
Related topic	OrthogonalPolynomials
Related topic	OrthogonalityOfChebyshevPolynomials
Related topic	SubstitutionNotation

We start from the first order differential equation

$$(1-x^2)\frac{du}{dx} + 2nxu = 0, \quad (1)$$

where one can <http://planetmath.org/SeparationOfVariables> separate the variables and then get the general solution

$$u = C(1-x^2)^n. \quad (2)$$

Differentiating  $n+1$  times the equation (1) it takes the form

$$(1-x^2)\frac{d^{n+2}u}{dx^{n+2}} - 2x\frac{d^{n+1}u}{dx^{n+1}} + n(n+1)\frac{d^nu}{dx^n} = 0$$

or

$$(1-x^2)\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} + n(n+1)y = 0 \quad (3)$$

where

$$y = \frac{d^nu}{dx^n} = C\frac{d^n}{dx^n}(1-x^2)^n.$$

Especially, the particular solution

$$y = P_n(x) := \frac{1}{2^n n!} \frac{d^n}{dx^n} (1-x^2)^n, \quad (4)$$

which which is the Legendre polynomial of degree  $n$ , has been seen to satisfy the Legendre's differential equation (3).

The equality (4) is <http://planetmath.org/RodriguesFormula> Rodrigues formula. We use it to find the leading coefficient of  $P_n(x)$  and to show the <http://planetmath.org/OrthogonalPolynomials> orthogonality of the Legendre polynomials  $P_0(x)$ ,  $P_1(x)$ ,  $P_2(x)$ , ...

## 0.1 The coefficient of $x^n$

By the binomial theorem,

$$\begin{aligned} P_n(x) &= \frac{1}{2^n n!} \frac{d^n}{dx^n} \sum_{j=0}^n \binom{n}{j} x^{2(n-j)} (-1)^j \\ &= \frac{1}{2^n n!} \sum_{j=0}^n \binom{n}{j} (2n-2j)(2n-2j-1) \cdots (2n-2j-n+1) x^{n-2j} (-1)^j. \end{aligned}$$

From the term with  $j = 0$  we get as the coefficient of  $x^n$  the following:

$$\frac{1}{2^n n!} \binom{n}{0} (2n)(2n-1)(2n-2) \cdots (2n-n+1)(-1)^0 = \frac{1}{2^n n!} \cdot \frac{(2n)!}{(2n-n)!} = \frac{(2n)!}{2^n (n!)^2} \quad (5)$$

## 0.2 Orthogonality

Let  $f_m(x) := a_0 + a_1 x + \dots + a_m x^m$  be any polynomial of degree  $m < n$ . <http://planetmath.org/IntegrationByParts> Integrating by parts  $m$  times we obtain

$$\begin{aligned} \int_{-1}^1 f_m(x) P_n(x) dx &= \frac{1}{2^n n!} \int_{-1}^1 f_m(x) \frac{d^n}{dx^n} (x^2-1)^n dx \\ &= \frac{1}{2^n n!} \int_{-1}^1 f_m(x) \frac{d^{n-1}}{dx^{n-1}} (x^2-1)^n - \frac{1}{2^n n!} \int_{-1}^1 f'_m(x) \frac{d^{n-1}}{dx^{n-1}} (x^2-1)^n dx \\ &\dots \dots \\ &= (-1)^m \frac{a_m m!}{2^n n!} \int_{-1}^1 \frac{d^{n-m}}{dx^{n-m}} (x^2-1)^n dx \\ &= (-1)^m \frac{a_m m!}{2^n n!} \int_{-1}^1 f_m(x) \frac{d^{n-m-1}}{dx^{n-m-1}} (x^2-1)^n = 0, \end{aligned}$$

since  $x = \pm 1$  are zeros of the derivatives  $\frac{d^{n-k}}{dx^{n-k}} (x^2-1)^n$ .

If, on the other hand,  $m = n$ , the calculation gives firstly

$$\int_{-1}^1 f_n(x) P_n(x) dx = 2(-1)^n \frac{a_n}{2^n} \int_0^1 (x^2-1)^n dx = 2(-1)^n \frac{a_n}{2^n} \cdot I_n, \quad (6)$$

where the integral  $I_n$  is gotten from

$$I_n = \int_0^1 x(x^2-1)^n - 2n \int_0^1 x^2(x^2-1)^{n-1} dx = -2n \int_0^1 [(x^2-1)^n + (x^2-1)^{n-1}] dx = -2n I_n - 2n I_{n-1}$$

Thus we infer the recurrence relation

$$I_n = -\frac{2n}{2n+1} I_{n-1}.$$

Using this and  $I_0 = 1$  one easily arrives at

$$I_n = (-1)^n \frac{2 \cdot 4 \cdot 6 \cdots (2n)}{3 \cdot 5 \cdot 7 \cdots (2n+1)} = (-1)^n \frac{[2 \cdot 4 \cdot 6 \cdots (2n)]^2}{(2n+1)!} = (-1)^n \frac{2^{2n}(n!)^2}{(2n+1)!}. \quad (7)$$

If  $f_n(x)$  also is a Legendre polynomial  $P_n(x)$ , we can in (6) by (5) put

$$a_n = \frac{(2n)!}{2^n(n!)^2}$$

and taking into account (7), too, (6) reads

$$\int_{-1}^1 [P_n(x)]^2 dx = \frac{(-1)^n}{2^{n-1}} \cdot \frac{(2n)!}{2^n(n!)^2} \cdot (-1)^n \frac{2^{2n}(n!)^2}{(2n+1)!} = \frac{2}{2n+1}.$$

Our results imply the <http://planetmath.org/Orthonormalorthonormality> condition

$$\int_{-1}^1 P_m(x)P_n(x) dx = \frac{2}{2n+1} \delta_{mn}, \quad (8)$$

where  $\delta_{mn}$  is the Kronecker delta.

## References

- [1] K. KURKI-SUONIO: *Matemaattiset apuneuvot*. Limes r.y., Helsinki (1966).