



Math for the people, by the people.

rectification theorem

Canonical name	RectificationTheorem
Date of creation	2013-03-22 14:57:15
Last modified on	2013-03-22 14:57:15
Owner	Daume (40)
Last modified by	Daume (40)
Numerical id	12
Author	Daume (40)
Entry type	Theorem
Classification	msc 34-00
Classification	msc 34A12
Related topic	ImplicitFunctionTheorem
Related topic	TangentMap

Let U be an open subset of \mathbb{R}^n and let $f \in C^1(U)$ be a continuous differentiable vector field

$$f: U \rightarrow \mathbb{R}^n.$$

If there exists $x_0 \in U$ such that $f(x_0) \neq 0$ then there exists $U_0 \subset U$ an open neighborhood of x_0 such that there exists a diffeomorphism of class C^1

$$F: U_0 \rightarrow V$$

where V is an open subset of \mathbb{R}^n such that

$$[DF(x)]f(x) = e_1$$

for all $x \in U_0$ where $[DF(x)]$ is the Jacobian of the diffeomorphism F evaluated at x and $e_1 = (1, 0, \dots, 0)$ is the first vector of the canonical basis of \mathbb{R}^n . More generally if the vector field f is of class C^r then so is the diffeomorphism F . [?]

References

[AVI] Arnold, V.I.: Ordinary Differential Equations (*translated by R.A. Silverman*). The MIT Press, Cambridge, 1973.