

## planetmath.org

Math for the people, by the people.

## residues of tangent and cotangent

Canonical name ResiduesOfTangentAndCotangent

Date of creation 2013-03-22 18:57:35 Last modified on 2013-03-22 18:57:35

Owner pahio (2872) Last modified by pahio (2872)

Numerical id 6

Author pahio (2872)
Entry type Example
Classification msc 33B10
Classification msc 30D10
Classification msc 30A99
Related topic Residue

Related topic TechniqueForComputingResidues
Related topic ResiduesOfGammaFunction

We will determine the residues of the tangent and the cotangent at their poles, which by the http://planetmath.org/node/9074parent entry are http://planetmath.org/SimplePolesimple.

By the rule in the entry coefficients of Laurent series, in a simple pole z=a of f one has

$$\operatorname{Res}(f; a) = \lim_{z \to a} (z - a) f(z).$$

• We get first

Res(cot; 0) = 
$$\lim_{z \to 0} z \cot z = \lim_{z \to 0} \frac{\cos z}{\frac{\sin z}{z}} = \frac{1}{1} = 1.$$
 (1)

• All the poles of cotangent are  $n\pi$  with  $n \in \mathbb{Z}$ . Since  $\pi$  is the period of cotangent, we could infer that the residues in all poles are the same as (1). We may also calculate (with the change of variable  $z - n\pi = w$ ) directly

$$\operatorname{Res}(\cot; \, n\pi) \; = \; \lim_{z \to n\pi} (z - n\pi) \cot z \; = \; \lim_{w \to 0} w \cot(w + n\pi) \; = \; \lim_{w \to 0} w \cot w \; = \; 1.$$

• In the http://planetmath.org/ComplexTangentAndCotangentparent entry, the complement formula for the tangent function is derived. Using it, we can find the residues of tangent at its poles  $\frac{\pi}{2} + n\pi$ , which are . For example,

$$\operatorname{Res}(\tan; \frac{\pi}{2}) \ = \ \lim_{z \to \frac{\pi}{2}} \left(z - \frac{\pi}{2}\right) \cot \left(\frac{\pi}{2} - z\right) \ = \ \lim_{w \to 0} w \cot (-w) \ = \ -\operatorname{Res}(\cot; \, 0) \ = \ -1.$$

Similarly as above, the residues in other poles are -1.

Consequently, the residues of cotangent are equal to 1 and the residues of tangent equal to -1.