



Math for the people, by the people.

## Riccati equation

Canonical name	RiccatiEquation
Date of creation	2013-03-22 18:05:43
Last modified on	2013-03-22 18:05:43
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	12
Author	pahio (2872)
Entry type	Result
Classification	msc 34A34
Classification	msc 34A05
Synonym	Riccati differential equation
Related topic	BernoulliEquation

The nonlinear differential equation

$$\frac{dy}{dx} = f(x) + g(x)y + h(x)y^2 \quad (1)$$

is called *Riccati equation*. If  $h(x) \equiv 0$ , it is a question of a linear differential equation; if  $f(x) \equiv 0$ , of a Bernoulli equation. There is no general method for integrating explicitly the equation (1), but via the substitution

$$y := -\frac{w'(x)}{h(x)w(x)}$$

one can convert it to a homogeneous linear differential equation with non-constant coefficients.

If one can find a particular solution  $y_0(x)$ , then one can easily verify that the substitution

$$y := y_0(x) + \frac{1}{w(x)} \quad (2)$$

converts (1) to

$$\frac{dw}{dx} + [g(x) + 2h(x)y_0(x)]w + h(x) = 0, \quad (3)$$

which is a linear differential equation of first order with respect to the function  $w = w(x)$ .

**Example.** The Riccati equation

$$\frac{dy}{dx} = 3 + 3x^2y - xy^2 \quad (4)$$

has the particular solution  $y := 3x$ . Solve the equation.

We substitute  $y := 3x + \frac{1}{w(x)}$  to (4), getting

$$\frac{dw}{dx} - 3x^2w - x = 0.$$

For solving this <http://planetmath.org/LinearDifferentialEquationOfFirstOrderfirst> order equation we can put  $w = uv$ ,  $w' = uv' + u'v$ , writing the equation as

$$u \cdot (v' - 3x^3v) + u'v := x, \quad (5)$$

where we choose the value of the expression in parentheses equal to 0:

$$\frac{dv}{dx} - 3x^2v = 0$$

After separation of variables and integrating, we obtain from here a solution  $v = e^{x^3}$ , which is set to the equation (5):

$$\frac{du}{dx} e^{x^3} = x$$

Separating the variables yields

$$du = \frac{x}{e^{x^3}} dx$$

and integrating:

$$u = C + \int x e^{-x^3} dx.$$

Thus we have

$$w = w(x) = uv = e^{x^3} \left[ C + \int x e^{-x^3} dx \right],$$

whence the general solution of the Riccati equation (4) is

$$y = 3x + \frac{e^{-x^3}}{C + \int x e^{-x^3} dx}.$$

It may be proved that if one knows three different solutions of Riccati equation (1), the each other solution may be expresses as a rational function of them.