



planetmath.org

Math for the people, by the people.

duality of Gudermannian and its inverse function

Canonical name	DualityOfGudermannianAndItsInverseFunction
Date of creation	2013-03-22 19:06:41
Last modified on	2013-03-22 19:06:41
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	7
Author	pahio (2872)
Entry type	Topic
Classification	msc 33B10
Classification	msc 26E05
Classification	msc 26A09
Classification	msc 26A48
Related topic	InverseGudermannianFunction
Related topic	IdealInvertingInPruferRing

There are a lot of formulae concerning the Gudermannian function and its inverse function containing a hyperbolic function or a trigonometric function or both, such that if we change functions of one kind to the corresponding functions of the other kind, then the new formula also is true.

Some examples:

$$\operatorname{gd} x = \int_0^x \frac{dt}{\cosh t}, \quad \operatorname{gd}^{-1} x = \int_0^x \frac{dt}{\cos t} \quad (1)$$

$$\frac{d}{dx} \operatorname{gd} x = \frac{1}{\cosh x}, \quad \frac{d}{dx} \operatorname{gd}^{-1} x = \frac{1}{\cos x} \quad (2)$$

$$\tan(\operatorname{gd} x) = \sinh x, \quad \tanh(\operatorname{gd}^{-1} x) = \sin x \quad (3)$$

$$\sin(\operatorname{gd} x) = \tanh x, \quad \sinh(\operatorname{gd}^{-1} x) = \tan x \quad (4)$$

$$\tan \frac{\operatorname{gd} x}{2} = \tanh \frac{x}{2}, \quad \tanh \frac{\operatorname{gd}^{-1} x}{2} = \tan \frac{x}{2} \quad (5)$$

For proving (5) we can check that

$$\frac{d}{dx} [2 \arctan(\tanh \frac{x}{2})] = \frac{1}{\cosh x},$$

and since both the expression in the brackets and the <http://planetmath.org/node/11997> Gudermannian function vanish in the origin, we have

$$\operatorname{gd} x \equiv 2 \arctan(\tanh \frac{x}{2}).$$

This equation implies (5).

The <http://planetmath.org/DualityInMathematics> duality of the formula pairs may be explained by the equality

$$\operatorname{gd} ix = i \operatorname{gd}^{-1} x. \quad (6)$$