



Math for the people, by the people.

spherical mean

Canonical name	SphericalMean
Date of creation	2013-03-22 14:09:04
Last modified on	2013-03-22 14:09:04
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	7
Author	rspuzio (6075)
Entry type	Definition
Classification	msc 35L05
Classification	msc 26E60
Related topic	WaveEquation

Let h be a function (usually real or complex valued) on \mathbb{R}^n ($n \geq 1$). Its *spherical mean* at point x over a sphere of radius r is defined as

$$M_h(x, r) = \frac{1}{A(n-1)} \int_{\|\xi\|=1} h(x + r\xi) dS = \frac{1}{A(n-1, r)} \int_{\|\xi\|=|r|} h(x + \xi) dS,$$

where the integral is over the surface of the unit $n-1$ -sphere. Here $A(n-1)$ is the area of the unit sphere, while $A(n-1, r) = r^{n-1}A(n-1)$ is the <http://planetmath.org/AreaOfTheNSphere> area of a sphere of radius r . In essence, the spherical mean $M_h(x, r)$ is just the average of h over the surface of a sphere of radius r centered at x , as the name suggests.

The spherical mean is defined for both positive and negative r and is independent of its sign. As $r \rightarrow 0$, if h is continuous, $M_h(x, r) \rightarrow h(x)$. If h has two continuous derivatives (is in $C^2(\mathbb{R}^n)$) then the following identity holds:

$$\nabla_x^2 M_h(x, r) = \left(\frac{\partial^2}{\partial r^2} + \frac{n-1}{r} \frac{\partial}{\partial r} \right) M_h(x, r),$$

where ∇^2 is the Laplacian.

Spherical means are used to obtain an explicit general solution for the wave equation in n space and one time dimensions.