



# d'Alembert and D. Bernoulli solutions of wave equation

Canonical name	DAlembertAndDBernoulliSolutionsOfWaveEquation
Date of creation	2013-03-22 18:23:15
Last modified on	2013-03-22 18:23:15
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	11
Author	pahio (2872)
Entry type	Derivation
Classification	msc 35L15
Classification	msc 35L05
Related topic	AdditionFormulasForSineAndCosine
Related topic	SchrodingersWaveEquation

Let's consider the <http://planetmath.org/WaveEquation>d'Alembert's solution

$$u(x, t) := \frac{1}{2}[f(x-ct) + f(x+ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds \quad (1)$$

of the wave equation in one dimension in the special case when the other initial condition is

$$u'_t(x, 0) := g(x) \equiv 0. \quad (2)$$

We shall see that the solution is equivalent with the solution of D. Bernoulli.

We the given function  $f$  to the Fourier sine series on the interval  $[0, p]$ :

$$f(y) = \sum_{n=1}^{\infty} A_n \sin \frac{n\pi y}{p} \quad \text{with} \quad A_n = \frac{2}{p} \int_0^p f(x) \sin \frac{n\pi x}{p} dx \quad (n = 1, 2, \dots)$$

Thus we may write

$$\begin{cases} f(x-ct) = \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{p} - \frac{n\pi ct}{p}\right) = \sum_{n=1}^{\infty} A_n \left(\sin \frac{n\pi x}{p} \cos \frac{n\pi ct}{p} - \cos \frac{n\pi x}{p} \sin \frac{n\pi ct}{p}\right), \\ f(x+ct) = \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{p} + \frac{n\pi ct}{p}\right) = \sum_{n=1}^{\infty} A_n \left(\sin \frac{n\pi x}{p} \cos \frac{n\pi ct}{p} + \cos \frac{n\pi x}{p} \sin \frac{n\pi ct}{p}\right). \end{cases}$$

Adding these equations and dividing by 2 yield

$$u(x, t) = \frac{1}{2}[f(x-ct) + f(x+ct)] = \sum_{n=1}^{\infty} A_n \cos \frac{n\pi ct}{p} \sin \frac{n\pi x}{p}, \quad (3)$$

which indeed is the <http://planetmath.org/SolvingTheWaveEquationByDBernoullisolution> of D. Bernoulli in the case  $g(x) \equiv 0$ .

**Note.** The solution (3) of the wave equation is especially in the special case where one has besides (2) the sine-formed initial condition

$$u(x, 0) := f(x) \equiv \sin \frac{\pi x}{p}. \quad (4)$$

Then  $A_n = 0$  for every  $n$  except 1, and one obtains

$$u(x, t) = \cos \frac{\pi ct}{p} \sin \frac{\pi x}{p}. \quad (5)$$

**Remark.** In the case of quantum systems one has <http://planetmath.org/SchrodingersWaveEquation> wave equation whose solutions are different from the above.