



planetmath.org

Math for the people, by the people.

Ω -stability theorem

Canonical name	OmegastabilityTheorem
Date of creation	2013-03-22 14:30:55
Last modified on	2013-03-22 14:30:55
Owner	Koro (127)
Last modified by	Koro (127)
Numerical id	8
Author	Koro (127)
Entry type	Theorem
Classification	msc 37C75
Synonym	omega-stability theorem
Synonym	Smale's Ω -stability theorem
Defines	Ω -stable
Defines	omega-stable
Defines	Ω -stability
Defines	omega-stability

Let M be a differentiable manifold and let $f: M \rightarrow M$ be a \mathcal{C}^k diffeomorphism. We say that f is \mathcal{C}^k - Ω -stable, if there is a neighborhood \mathcal{U} of f in the \mathcal{C}^k topology of $\text{Diff}^k(M)$ such that for any $g \in \mathcal{U}$, $f|_{\Omega(f)}$ is topologically conjugate to $g|_{\Omega(g)}$.

Ω -stability theorem. If f is Axiom A and satisfies the no-cycles condition, then f is \mathcal{C}^1 - Ω -stable.

Remark. The reciprocal of this theorem is also true (the difficult part is showing that Ω -stability implies Axiom A), but it is unknown whether \mathcal{C}^k - Ω -stability implies Axiom A when $k > 1$. This is known as the \mathcal{C}^k Ω -stability conjecture.