



planetmath.org

Math for the people, by the people.

hyperbolic isomorphism

Canonical name	HyperbolicIsomorphism
Date of creation	2013-03-22 13:39:34
Last modified on	2013-03-22 13:39:34
Owner	Koro (127)
Last modified by	Koro (127)
Numerical id	10
Author	Koro (127)
Entry type	Definition
Classification	msc 37D05
Classification	msc 46B03
Synonym	linear hyperbolic isomorphism

Let X be a Banach space and $T : X \rightarrow X$ a continuous linear isomorphism. We say that T is an *hyperbolic isomorphism* if its spectrum is disjoint with the unit circle, i.e. $\sigma(T) \cap \{z \in \mathbb{C} : |z| = 1\} = \emptyset$.

If this is the case, by the spectral theorem there is a splitting of X into two invariant subspaces, $X = E^s \oplus E^u$ (and therefore, a corresponding splitting of T into two operators $T^s : E^s \rightarrow E^s$ and $T^u : E^u \rightarrow E^u$, i.e. $T = T^s \oplus T^u$), such that $\sigma(T^s) = \sigma(T) \cap \{z : |z| < 1\}$ and $\sigma(T^u) = \sigma(T) \cap \{z : |z| > 1\}$. Also, for any λ greater than the spectral radius of both T^s and T^u there exists an equivalent (box-type) norm $\|\cdot\|_1$ such that

$$\|T^s\|_1 < \lambda \text{ and } \|T^u\|_1 < \lambda$$

and

$$\|x\|_1 = \max\{\|x^u\|_1, \|x^s\|_1\}.$$

In particular, λ can be chosen smaller than 1, so that T^s and T^u are contractions.