



stable manifold

Canonical name	StableManifold
Date of creation	2013-03-22 13:41:07
Last modified on	2013-03-22 13:41:07
Owner	Koro (127)
Last modified by	Koro (127)
Numerical id	11
Author	Koro (127)
Entry type	Definition
Classification	msc 37D10
Classification	msc 37C75
Synonym	stable set
Synonym	unstable set
Synonym	unstable manifold
Related topic	HyperbolicFixedPoint

Let X be a topological space, and $f: X \rightarrow X$ a homeomorphism. If p is a fixed point for f , the **stable and unstable sets** of p are defined by

$$W^s(f, p) = \{q \in X : f^n(q) \xrightarrow[n \rightarrow \infty]{} p\},$$

$$W^u(f, p) = \{q \in X : f^{-n}(q) \xrightarrow[n \rightarrow \infty]{} p\},$$

respectively.

If p is a periodic point of least period k , then it is a fixed point of f^k , and the stable and unstable sets of p are

$$W^s(f, p) = W^s(f^k, p)$$

$$W^u(f, p) = W^u(f^k, p).$$

Given a neighborhood U of p , the **local stable and unstable sets** of p are defined by

$$W_{\text{loc}}^s(f, p, U) = \{q \in U : f^n(q) \in U \text{ for each } n \geq 0\},$$

$$W_{\text{loc}}^u(f, p, U) = W_{\text{loc}}^s(f^{-1}, p, U).$$

If X is metrizable, we can define the stable and unstable sets for any point by

$$W^s(f, p) = \{q \in U : d(f^n(q), f^n(p)) \xrightarrow[n \rightarrow \infty]{} 0\},$$

$$W^u(f, p) = W^s(f^{-1}, p),$$

where d is a metric for X . This definition clearly coincides with the previous one when p is a periodic point.

When K is an invariant subset of X , one usually denotes by $W^s(f, K)$ and $W^u(f, K)$ (or just $W^s(K)$ and $W^u(K)$) the stable and unstable sets of K , defined as the set of points $x \in X$ such that $d(f^n(x), K) \rightarrow 0$ when $x \rightarrow \infty$ or $-\infty$, respectively.

Suppose now that X is a compact smooth manifold, and f is a \mathcal{C}^k diffeomorphism, $k \geq 1$. If p is a hyperbolic periodic point, the stable manifold theorem assures that for some neighborhood U of p , the local stable and unstable sets are \mathcal{C}^k embedded disks, whose tangent spaces at p are E^s and E^u (the stable and unstable spaces of $Df(p)$), respectively; moreover, they vary continuously (in certain sense) in a neighborhood of f in the \mathcal{C}^k topology of $\text{Diff}^k(X)$ (the space of all \mathcal{C}^k diffeomorphisms from X to itself). Finally,

the stable and unstable sets are \mathcal{C}^k injectively immersed disks. This is why they are commonly called **stable and unstable** manifolds. This result is also valid for nonperiodic points, as long as they lie in some hyperbolic set (stable manifold theorem for hyperbolic sets).