



Math for the people, by the people.

divided difference

Canonical name	DividedDifference
Date of creation	2013-03-22 14:40:59
Last modified on	2013-03-22 14:40:59
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	10
Author	rspuzio (6075)
Entry type	Definition
Classification	msc 39A70

Let  $f$  be a real (or complex) function. Given distinct real (or complex) numbers  $x_0, x_1, x_2, \dots$ , the *divided differences* of  $f$  are defined recursively as follows:

$$\Delta^1 f[x_0, x_1] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$$

$$\Delta^{n+1} f[x_0, x_1, \dots, x_{n+1}] = \frac{\Delta^n f[x_1, x_2, \dots, x_{n+1}] - \Delta^n f[x_0, x_2, \dots, x_{n+1}]}{x_1 - x_0}$$

It is also convenient to define the zeroth divided difference of  $f$  to be  $f$  itself:

$$\Delta^0 f[x_0] = f[x_0]$$

Some important properties of divided differences are:

1. Divided differences are invariant under permutations of  $x_0, x_1, x_2, \dots$
2. If  $f$  is a polynomial of order  $m$  and  $m < n$ , then the  $n$ -th divided differences of  $f$  vanish identically
3. If  $f$  is a polynomial of order  $m + n$ , then  $\Delta^n(x, x_1, \dots, x_n)$  is a polynomial in  $x$  of order  $m$ .

Divided differences are useful for interpolating functions when the values are given for unequally spaced values of the argument.

Because of the first property listed above, it does not matter with respect to which two arguments we compute the divided difference when we compute the  $n + 1$ -st divided difference from the  $n$ -th divided difference. For instance, when computing the divided difference table for tabulated values of a function, a convenient choice is the following:

$$\Delta^{n+1} f[x_0, x_1, \dots, x_{n+1}] = \frac{\Delta^n f[x_1, x_2, \dots, x_{n+1}] - \Delta^n f[x_0, x_1, \dots, x_n]}{x_{n+1} - x_0}$$