



planetmath.org

Math for the people, by the people.

solution of equations by divided difference interpolaton

Canonical name	SolutionOfEquationsByDividedDifferenceInterpolaton
Date of creation	2013-03-22 16:49:22
Last modified on	2013-03-22 16:49:22
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	16
Author	rspuzio (6075)
Entry type	Application
Classification	msc 39A70

Divided difference interpolation can be used to obtain approximate solutions to equations and to invert functions numerically. The idea is that, given an equation $f(y) = x$ which we want to solve for y , we first take several numbers y_1, \dots, y_n and compute $x_1 \dots x_n$ as $x_i = f(y_i)$. Then we compute the divided differences of the y_i 's regarded as functions of the x_i 's and form the divided difference series. Substituting x in this series provides an approximation to y .

To illustrate how this works, we will examine the transcendental equation $x + e^{-x} = 2$. We note that $2 + e^{-2} = 2.13533$ and $1.5 + e^{-1.5} = 1.72313$, so there will be a solution between 1.5 and 2, likely closer to 2 than 1.5. Therefore, as our values of the y_i 's, we shall take 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1. We now tabulate $x_i = y_i + e^{-y_i}$ for those values:

y_i	x_i
1.5	1.72313
1.6	1.80190
1.7	1.88268
1.8	1.96530
1.9	2.04957
2.0	2.13533
2.1	2.22246

Next, we form a divided difference table of the y_i 's as a function of the x_i 's:

1.72313	1.50000					
		1.26952				
1.80190	1.60000		-0.19799			
		1.23793		0.12082		
1.88268	1.70000		-0.16873		-0.039609	
		1.21036		0.10789		0.091553
1.96530	1.80000		-0.14201		-0.077347	-0.13457
		1.18666		0.08210		0.024360
2.04957	1.90000		-0.12127		-0.067102	
		1.16604		0.05930		
2.13533	2.00000		-0.10602			
		1.14771				
2.22246	2.10000					

From this table, we form the series

$$\begin{aligned}
& 1.50000 + 1.26952(x - 1.72313) - 0.19799(x - 1.72313)(x - 1.80190) \\
& + 0.12082(x - 1.72313)(x - 1.80190)(x - 1.88268) \\
& - 0.039609(x - 1.72313)(x - 1.80190)(x - 1.88268)(x - 1.96530) \\
& + 0.091553(x - 1.72313)(x - 1.80190)(x - 1.88268)(x - 1.96530)(x - 2.04957) \\
& - 0.13457(x - 1.72313)(x - 1.80190)(x - 1.88268)(x - 1.96530)(x - 2.04957)(x - 2.13533)
\end{aligned}$$

Substituting 2.00000 for x , we obtain 1.84140. Given that

$$1.84140 + e^{-1.84140} < 2 < 1.84141 + e^{-1.84141},$$

this answer is correct to all 5 decimal places.

In the presentation above, we tacitly assumed that there was a solution to our equation and focussed our attention on finding that answer numerically. To complete the treatment we will now show that there indeed exists a unique solution to the equation $x + e^{-x} = 2$ in the interval $(0, \infty)$.

Existence follows from the intermediate value theorem. As noted above,

$$1.5 + e^{-1.5} < 2 < 3 + e^{-2}.$$

Since $x + e^{-x}$ depends continuously on x , it follows that there exists $x \in (1.5, 2)$ such that $x + e^{-x} = 2$.

As for uniqueness, note that the derivative of $x + e^{-x}$ is $1 - e^{-x}$. When $x > 0$, we have $e^{-x} < 1$, or $1 - e^{-x} > 0$. Hence, $x + e^{-x}$ is a strictly increasing function of x , so there can be at most one x such that $x + e^{-x} = 2$.