

planetmath.org

Math for the people, by the people.

divided differences of powers

Canonical name DividedDifferencesOfPowers

Date of creation 2013-03-22 16:47:59 Last modified on 2013-03-22 16:47:59

Owner rspuzio (6075) Last modified by rspuzio (6075)

Numerical id 11

Author rspuzio (6075) Entry type Theorem Classification msc 39A70 In this entry, we will prove the claims about divided differences of polynomials. Because the divided difference is a linear operator, we can focus our attention on powers.

Theorem 1. If $f(x) = x^n$ and $m \le n$, then

$$\Delta^m f[x_0, \dots x_m] = \sum_{k_0 + \dots + k_m = n - m} x_0^{k_0} \dots x_m^{k_m}.$$

If m > n, then $\Delta^m f[x_0, \dots x_m] = 0$.

Proof. We proceed by induction. The formula is trivially true when m = 0. Assume that the formula is true for a certain value of m. Then we have

$$\Delta^{m+1}f[x_0, \dots x_{m+1}] = \frac{\Delta^m f[x_1, x_2 \dots x_m] - \Delta^m f[x_0, x_2 \dots x_m]}{x_1 - x_0}$$
$$= \sum_{k+k_2+\dots+k_m=n-m} \frac{(x_1^k - x_0^k)x_2^{k_2} \dots x_m^{k_m}}{x_1 - x_0}$$

Using the identity for the sum of a geometric series,

$$\frac{x_1^k - x_0^k}{x_1 - x_0} = \sum_{k_0 + k_1 = k - 1} x_0^{k_0} x_1^{k_1},$$

this becomes

$$\Delta^{m+1} f[x_0, \dots x_{m+1}] = \sum_{k+k_2+\dots+k_m=n-m} \sum_{k_0+k_1=k-1} x_0^{k_0} x_1^{k_1} x_2^{k_2} \dots x_{m+1}^{k_{m+1}}$$

$$= \sum_{k_0+k_1+k_2+\dots+k_m=n-(m+1)} x_0^{k_0} x_1^{k_1} x_2^{k_2} \dots x_{m+1}^{k_{m+1}}.$$

Note that when k=0, we have $x_1^k-x_0^k=0$, which is consistent with the formula given above because, in that case, there are no solutions to $k_1+k_2=k$, so the sum is empty and, by convention, equals zero. Likewise, when n=m, then the only solution to $k_0+\cdots+k_m=0$ is $k_0=\cdots=k_m=0$, so the sum only consists of one term, $x_0^0\cdots x_m^0=1$ so $\Delta^n f[x_0,\cdots x_n]=1$, hence taking further differences produces zero.