



planetmath.org

Math for the people, by the people.

midpoint rule

Canonical name	MidpointRule
Date of creation	2013-03-22 15:57:44
Last modified on	2013-03-22 15:57:44
Owner	Wkbj79 (1863)
Last modified by	Wkbj79 (1863)
Numerical id	16
Author	Wkbj79 (1863)
Entry type	Theorem
Classification	msc 41-01
Classification	msc 28-00
Classification	msc 26A42
Related topic	LeftHandRule
Related topic	RightHandRule
Related topic	RiemannSum
Related topic	ExampleOfEstimatingARiemannIntegral

The *midpoint rule* for computing the Riemann integral $\int_a^b f(x) dx$ is

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{j=1}^n f\left(a + \left(j - \frac{1}{2}\right) \left(\frac{b-a}{n}\right)\right) \left(\frac{b-a}{n}\right).$$

If the Riemann integral is considered as a measure of area under a curve, then the expressions $f\left(a + \left(j - \frac{1}{2}\right) \left(\frac{b-a}{n}\right)\right)$ the of the rectangles, and $\frac{b-a}{n}$ is the common of the rectangles.

The Riemann integral can be approximated by using a definite value for n rather than taking a limit. In this case, the partition is $\left\{ \left[a, a + \frac{b-a}{n} \right), \dots, \left[a + \frac{(b-a)(n-1)}{n}, b \right] \right\}$ and the function is evaluated at the midpoints of each of these intervals. Note that this is a special case of a Riemann sum in which the x_j 's are evenly spaced and the c_j 's chosen are the midpoints.

If f is Riemann integrable on $[a, b]$ such that $|f''(x)| \leq M$ for every $x \in [a, b]$, then

$$\left| \int_a^b f(x) dx - \sum_{j=1}^n f\left(a + \left(j - \frac{1}{2}\right) \left(\frac{b-a}{n}\right)\right) \left(\frac{b-a}{n}\right) \right| \leq \frac{M(b-a)^3}{24n^2}.$$