

planetmath.org

Math for the people, by the people.

example of Fourier series

Canonical name ExampleOfFourierSeries

Date of creation 2013-03-22 13:57:13 Last modified on 2013-03-22 13:57:13

Owner alozano (2414) Last modified by alozano (2414)

Numerical id 10

Author alozano (2414) Entry type Example

Classification msc 42A16

Synonym example of Fourier coefficients

Related topic ValueOfTheRiemannZetaFunctionAtS2

Related topic FourierSineAndCosineSeries

Here we present an example of Fourier series:

Example:

Let $f: (-\pi, \pi) \to \mathbb{R}$ be the "identity" function, defined by

$$f(x) = x$$
, for all $x \in (-\pi, \pi)$.

We will compute the Fourier coefficients for this function. Notice that cos(nx) is an even function, while f and sin(nx) are odd functions.

$$a_0^f = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} x dx = 0$$

$$a_n^f = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx = \frac{1}{\pi} \int_{-\pi}^{\pi} x \cos(nx) dx = 0$$

$$b_n^f = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx = \frac{1}{\pi} \int_{-\pi}^{\pi} x \sin(nx) dx = \frac{2}{\pi} \int_{0}^{\pi} x \sin(nx) dx = \frac{2}{\pi} \int_{0}^{\pi} x \sin(nx) dx = \frac{2}{\pi} \left(\left[-\frac{x \cos(nx)}{n} \right]_{0}^{\pi} + \left[\frac{\sin(nx)}{n^2} \right]_{0}^{\pi} = \right) = (-1)^{n+1} \frac{2}{n}$$

Notice that a_0^f , a_n^f are 0 because x and $x \cos(nx)$ are odd functions. Hence the Fourier series for f(x) = x is:

$$f(x) = x = a_0^f + \sum_{n=1}^{\infty} (a_n^f \cos(nx) + b_n^f \sin(nx)) =$$
$$= \sum_{n=1}^{\infty} (-1)^{n+1} \frac{2}{n} \sin(nx), \quad \forall x \in (-\pi, \pi)$$

For an application of this Fourier series, see value of the Riemann zeta function at s=2.