



planetmath.org

Math for the people, by the people.

Fourier series in complex form and Fourier integral

Canonical name	FourierSeriesInComplexFormAndFourierIntegral
Date of creation	2013-03-22 18:02:54
Last modified on	2013-03-22 18:02:54
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	10
Author	pahio (2872)
Entry type	Derivation
Classification	msc 42A38
Classification	msc 42A16
Classification	msc 44A55
Related topic	FourierTransform
Related topic	KalleVaisala
Defines	Fourier integral

0.1 Fourier series in complex form

The Fourier series expansion of a Riemann integrable real function f on the interval $[-p, p]$ is

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi t}{p} + b_n \sin \frac{n\pi t}{p} \right), \quad (1)$$

where the coefficients are

$$a_n = \frac{1}{p} \int_{-p}^p f(x) \cos \frac{n\pi t}{p} dt, \quad b_n = \frac{1}{p} \int_{-p}^p f(x) \sin \frac{n\pi t}{p} dt. \quad (2)$$

If one expresses the cosines and sines via <http://planetmath.org/ComplexSineAndCosineEuler> formulas with <http://planetmath.org/ComplexExponentialFunctionexponential> function, the series (1) attains the form

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{\frac{in\pi t}{p}}. \quad (3)$$

The coefficients c_n could be obtained of a_n and b_n , but they are comfortably derived directly by multiplying the equation (3) by $e^{-\frac{in\pi t}{p}}$ and integrating it from $-p$ to p . One obtains

$$c_n = \frac{1}{2p} \int_{-p}^p f(t) e^{-\frac{in\pi t}{p}} dt \quad (n = 0, \pm 1, \pm 2, \dots). \quad (4)$$

We may say that in (3), $f(t)$ has been dissolved to sum of *harmonics* (elementary waves) $c_n e^{\frac{in\pi t}{p}}$ with amplitudes c_n corresponding the frequencies n .

0.2 Derivation of Fourier integral

For seeing how the expansion (3) changes when $p \rightarrow \infty$, we put first the expressions (4) of c_n to the series (3):

$$f(t) = \sum_{n=-\infty}^{\infty} e^{\frac{in\pi t}{p}} \frac{1}{2p} \int_{-p}^p f(t) e^{-\frac{in\pi t}{p}} dt$$

By denoting $\omega_n := \frac{n\pi}{p}$ and $\Delta_n\omega := \omega_{n+1} - \omega_n = \frac{\pi}{p}$, the last equation takes the form

$$f(t) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{i\omega_n t} \Delta_n\omega \int_{-p}^p f(t) e^{-i\omega_n t} dt.$$

It can be shown that when $p \rightarrow \infty$ and thus $\Delta_n\omega \rightarrow 0$, the limiting form of this equation is

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\omega t} d\omega \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt. \quad (5)$$

Here, $f(t)$ has been represented as a *Fourier integral*. It can be proved that for validity of the expansion (4) it suffices that the function f is piecewise continuous on every finite interval having at most a finite amount of extremum points and that the integral

$$\int_{-\infty}^{\infty} |f(t)| dt$$

converges.

For better to compare to the Fourier series (3) and the coefficients (4), we can write (5) as

$$f(t) = \int_{-\infty}^{\infty} c(\omega) e^{i\omega t} d\omega, \quad (6)$$

where

$$c(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt. \quad (7)$$

0.3 Fourier transform

If we denote $2\pi c(\omega)$ as

$$F(\omega) = \int_{-\infty}^{\infty} e^{-i\omega t} f(t) dt, \quad (8)$$

then by (5),

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\omega t} F(\omega) d\omega. \quad (9)$$

$F(\omega)$ is called the *Fourier transform* of $f(t)$. It is an integral transform and (9) its inverse transform.

N.B. that often one sees both the formula (8) and the formula (9) equipped with the same constant factor $\frac{1}{\sqrt{2\pi}}$ in front of the integral sign.

References

- [1] K. VÄISÄLÄ: *Laplace-muunnos*. Handout Nr. 163. Teknillisen korkeakoulun ylioppilaskunta, Otaniemi, Finland (1968).