



Math for the people, by the people.

uncertainty theorem

Canonical name	UncertaintyTheorem
Date of creation	2013-03-22 18:50:37
Last modified on	2013-03-22 18:50:37
Owner	dh2718 (16929)
Last modified by	dh2718 (16929)
Numerical id	14
Author	dh2718 (16929)
Entry type	Theorem
Classification	msc 42C40
Related topic	UncertaintyPrinciple

The uncertainty principle, as first formulated by Heisenberg, states that the product of the standard deviations of two *conjugated* variables, cannot be less than some minimum. This statement has been generalized to a precise mathematical theorem, in the frame of *wavelet* theory.

1 THE UNCERTAINTY THEOREM

Let $f(t)$ be a real function of the real variable t , satisfying the L^2 condition (see below), and $F(\omega)$ its Fourier transform. The standard deviation Δ_t and Δ_ω of t and ω respectively, satisfy the following inequality:

$$\Delta_t \Delta_\omega \geq \frac{1}{2}$$

For this formula to make sense, Δ_t and Δ_ω must be precisely defined.

2 THE L^2 CONDITION

A real function $f(t)$ of the real variable t will be said to satisfy the L^2 condition if $f(t)$, $tf(t)$ and the derivative $f'(t)$ are all in L^2 .

If $F(\omega)$ is its Fourier transform, $-i\omega F(\omega)$ is the transform of $f'(t)$. All the following functions belong to L^2 :

$$f(t), f'(t), tf(t), F(\omega), \omega F(\omega)$$

The first three functions are just the definition and the two last ones result from Parseval's identity, recalled here in its integral form:

$$\int_{-\infty}^{\infty} \overline{U(\omega)} V(\omega) d\omega = 2\pi \int_{-\infty}^{\infty} \overline{u(t)} v(t) dt$$

$U(\omega)$ and $V(\omega)$ are the Fourier transforms of $u(t)$ and $v(t)$.

3 DEFINITIONS

$f(t)$ and $F(\omega)$ being in L^2 , we may define their finite norms:

$$\|f\|^2 = \int_{-\infty}^{\infty} |f(t)|^2 dt \quad \|F\|^2 = \int_{-\infty}^{\infty} |F(\omega)|^2 d\omega$$

By Parseval's identity, they are related:

$$\| f \|^2 = \frac{1}{2\pi} \| F \|^2$$

We are now able to define the probability distributions $T(t)$ and $\Omega(\omega)$ for the "random" variables t and ω :

$$T(t) = \frac{| f(t) |^2}{\| f \|^2} \quad \Omega(\omega) = \frac{| F(\omega) |^2}{\| F \|^2}$$

Since the L^2 integrals of T and Ω are 1, they are proper probability distributions. The *mean value* t_0 of t is defined the usual way:

$$t_0 = \int_{-\infty}^{\infty} t T(t) dt$$

Note that ω 's mean value is always 0 because $f(t)$ is a real function. Finally, we have the standard deviations for the uncertainty theorem:

$$\Delta_t^2 = \int_{-\infty}^{\infty} T(t)(t - t_0)^2 dt \quad \Delta_\omega^2 = \int_{-\infty}^{\infty} \Omega(\omega)(\omega - 0)^2 d\omega$$

4 PROOF OF THE THEOREM

The heart of the proof is the Cauchy-Schwarz inequality in the L^2 Hilbert space: the product of the norms of two functions $u(t)$ and $v(t)$ is greater than, or equal to, the norm of their scalar product:

$$\int_{-\infty}^{\infty} | u(t) |^2 dt \int_{-\infty}^{\infty} | v(t) |^2 dt \geq \left| \int_{-\infty}^{\infty} \overline{u(t)} v(t) dt \right|^2$$

Equality occurs if, and only if, one of the functions is proportional to the other. For the two functions $u(t) = (t - t_0)f(t)$ and $v(t) = f'(t)$, we have therefore:

$$\int_{-\infty}^{\infty} (t - t_0)^2 f(t)^2 dt \int_{-\infty}^{\infty} f'(t)^2 dt \geq \left| \int_{-\infty}^{\infty} (t - t_0) f(t) f'(t) dt \right|^2$$

The integral at the right hand side can be integrated by parts. Using the definition of $\| f \|^2$:

$$\int_{-\infty}^{\infty} (t - t_0)^2 f(t)^2 dt \int_{-\infty}^{\infty} f'(t)^2 dt \geq \frac{1}{4} \| f \|^4$$

But $\|f\|$ and $\|F\|$ are related by a 2π factor, so:

$$\int_{-\infty}^{\infty} (t - t_0)^2 f(t)^2 dt \int_{-\infty}^{\infty} f'(t)^2 dt \geq \frac{1}{8\pi} \|f\|^2 \|F\|^2$$

Applying Parseval's identity to the second integral of the left hand side, we get:

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} (t - t_0)^2 f(t)^2 dt \int_{-\infty}^{\infty} \omega^2 F(\omega)^2 d\omega \geq \frac{1}{8\pi} \|f\|^2 \|F\|^2$$

We have used the fact that the Fourier transform of $f'(t)$ is $-i\omega F(\omega)$. Now, dividing both sides by the norms, and simplifying by the 2π factor, we get exactly the uncertainty theorem:

$$\int_{-\infty}^{\infty} (t - t_0)^2 f(t)^2 dt \int_{-\infty}^{\infty} \omega^2 F(\omega)^2 d\omega \geq \frac{1}{4}$$

5 THE GAUSSIAN FUNCTION

The Cauchy-Schwarz inequality becomes an equality if, and only if, one of the functions is proportional to the other. In our case, this condition is expressed by $f'(t) = \lambda(t - t_0)f(t)$ where λ is a constant. This differential equation is readily solved: $f(t) = ke^{\lambda(t-t_0)^2}$. $f(t)$ must be in L^2 so that λ must be negative. Defining $\lambda = \frac{-1}{2\sigma^2}$, we get the gaussian function in its traditional form:

$$f(t) = e^{\frac{-(t-t_0)^2}{2\sigma^2}}$$

The constant k has been omitted because it cancels anyway in the probability distributions. The standard deviations are easily computed from their definitions:

$$\Delta_t = \frac{\sigma}{\sqrt{2}} \quad \Delta_\omega = \frac{1}{\sigma\sqrt{2}}$$

Their product is $\frac{1}{2}$, independent of σ . There is no other function with this property.

References

- [1] Roberto Celi *Time-Frequency visualization of helicopter noise*
<http://celi.umd.edu/Jour/NoisePaperColor.pdf>

Despite its frightening title, this paper is mostly theoretical and it is the only place where I saw the uncertainty theorem clearly stated.

- [2] Robi Polikar *The wavelet tutorial*
<http://users.rowan.edu/~polikar/wavelets/wttutorial.html>