



Math for the people, by the people.

## logarithmic convolution

Canonical name	LogarithmicConvolution
Date of creation	2013-03-22 14:28:26
Last modified on	2013-03-22 14:28:26
Owner	swiftset (1337)
Last modified by	swiftset (1337)
Numerical id	4
Author	swiftset (1337)
Entry type	Definition
Classification	msc 44A35
Synonym	scale convolution
Related topic	Convolution

**Definition** The *scale convolution* of two functions  $s(t)$  and  $r(t)$ , also known as their *logarithmic convolution* is defined as the function

$$s *_l r(t) = r *_l s(t) = \int_0^\infty s\left(\frac{t}{a}\right)r(a)\frac{da}{a}$$

when this quantity exists.

**Results** The logarithmic convolution can be related to the ordinary convolution by changing the variable from  $t$  to  $v = \log t$ :

$$\begin{aligned} s *_l r(t) &= \int_0^\infty s\left(\frac{t}{a}\right)r(a)\frac{da}{a} = \int_{-\infty}^\infty s\left(\frac{t}{e^u}\right)r(e^u)du \\ &= \int_{-\infty}^\infty s(e^{\log t - u})r(e^u)du \end{aligned}$$

Define  $f(v) = s(e^v)$  and  $g(v) = r(e^v)$  and let  $v = \log t$ , then

$$s *_l r(v) = f * g(v) = g * f(v) = r *_l s(v).$$