



Math for the people, by the people.

Heaviside formula

Canonical name	HeavisideFormula
Date of creation	2014-03-19 9:14:46
Last modified on	2014-03-19 9:14:46
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	11
Author	pahio (2872)
Entry type	Topic
Classification	msc 44A10
Synonym	Heaviside expansion formula
Synonym	inverse Laplace transform of rational function
Related topic	HyperbolicFunctions
Related topic	ComplexSineAndCosine

Let $P(s)$ and $Q(s)$ be polynomials with the degree of the former less than the degree of the latter.

- If all complex <http://planetmath.org/Zerozeroes> a_1, a_2, \dots, a_n of $Q(s)$ are simple, then

$$\mathcal{L}^{-1} \left\{ \frac{P(s)}{Q(s)} \right\} = \sum_{j=1}^n \frac{P(a_j)}{Q'(a_j)} e^{a_j t}. \quad (1)$$

- If the different zeroes a_1, a_2, \dots, a_n of $Q(s)$ have the multiplicities m_1, m_2, \dots, m_n , respectively, we denote $F_j(s) := (s - a_j)^{m_j} P(s) / Q(s)$; then

$$\mathcal{L}^{-1} \left\{ \frac{P(s)}{Q(s)} \right\} = \sum_{j=1}^n e^{a_j t} \sum_{k=0}^{m_j-1} \frac{F_j^{(k)}(a_j) t^{m_j-1-k}}{k! (m_j-1-k)!}. \quad (2)$$

A special case of the *Heaviside formula* (1) is

$$\mathcal{L}^{-1} \left\{ \frac{Q'(s)}{Q(s)} \right\} = \sum_{j=1}^n e^{a_j t}. \quad (3)$$

Example. Since the zeros of the binomial $s^4 + 4a^4$ are $s = (\pm 1 \pm i)a$, we can calculate by (3) as follows:

$$\mathcal{L}^{-1} \left\{ \frac{s^3}{s^4 + 4a^4} \right\} = \frac{1}{4} \mathcal{L}^{-1} \left\{ \frac{4s^3}{s^4 + 4a^4} \right\} = \frac{1}{4} \sum_{\pm} e^{(\pm 1 \pm i)at} = \frac{e^{at} + e^{-at}}{2} \cdot \frac{e^{iat} + e^{-iat}}{2} = \cosh at \cos at$$

Proof of (1). Without hurting the generality, we can suppose that $Q(s)$ is monic. Therefore

$$Q(s) = (s - a_1)(s - a_2) \cdots (s - a_n).$$

For $j = 1, 2, \dots, n$, denoting

$$Q_j(s) := (s - a_j)Q(s),$$

one has $Q_j(a_j) \neq 0$. We have a partial fraction expansion of the form

$$\frac{P(s)}{Q(s)} = \frac{C_1}{s-a_1} + \frac{C_2}{s-a_2} + \dots + \frac{C_n}{s-a_n} \quad (4)$$

with constants C_j . According to the linearity and the formula 1 of the <http://planetmath.org/LaplaceTransformparent> entry, one gets

$$\mathcal{L}^{-1} \left\{ \frac{P(s)}{Q(s)} \right\} = \sum_{j=1}^n C_j e^{a_j t}. \quad (5)$$

For determining the constants C_j , multiply (3) by $s-a_j$. It yields

$$\frac{P(s)}{Q_j(s)} = C_j + (s-a_j) \sum_{\nu \neq j} \frac{C_\nu}{s-a_\nu}.$$

Setting to this identity $s := a_j$ gives the value

$$C_j = \frac{P(a_j)}{Q_j(a_j)}. \quad (6)$$

But since $Q'(s) = \frac{d}{ds}((s-a_j)Q_j(s)) = Q_j(s) + (s-a_j)Q_j'(s)$, we see that $Q'(a_j) = Q_j(a_j)$; thus the equation (5) may be written

$$C_j = \frac{P(a_j)}{Q'(a_j)}. \quad (7)$$

The values (6) in (4) produce the formula (1).

References

- [1] K. VÄISÄLÄ: *Laplace-muunnos*. Handout Nr. 163. Teknillisen korkeakoulun ylioppilaskunta, Otaniemi, Finland (1968).