



planetmath.org

Math for the people, by the people.

## integration of Laplace transform with respect to parameter

Canonical name	IntegrationOfLaplaceTransformWithRespectToParameter
Date of creation	2013-03-22 18:44:47
Last modified on	2013-03-22 18:44:47
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	8
Author	pahio (2872)
Entry type	Theorem
Classification	msc 44A10
Related topic	TableOfLaplaceTransforms
Related topic	TermwiseDifferentiation
Related topic	MethodsOfEvaluatingImproperIntegrals
Related topic	UsingConvolutionToFindLaplaceTransform
Related topic	RelativeOfCosineIntegral
Related topic	RelativeOfExponentialIntegral

We use the curved  $\curvearrowright$  from the Laplace-transformed functions to the corresponding initial functions.

If

$$f(t, x) \curvearrowright F(s, x),$$

then one can integrate both functions with respect to the parametre  $x$  between the same  $a$  which may be also infinite provided that the integrals converge:

$$\int_a^b f(t, x) dx \curvearrowright \int_a^b F(s, x) dx \quad (1)$$

(1) may be written as

$$\mathcal{L}\left\{\int_a^b f(t, x) dx\right\} = \int_a^b \mathcal{L}\{f(t, x)\} dx. \quad (2)$$

*Proof.* Using the definition of the Laplace transform, we can write

$$\int_a^b f(t, x) dx \curvearrowright \int_0^\infty \left( e^{-st} \int_a^b f(s, x) dx \right) dt = \int_0^\infty \left( \int_a^b e^{-st} f(s, x) dx \right) dt.$$

We change the order of integration in the last double integral and use again the definition, obtaining

$$\int_a^b f(t, x) dx \curvearrowright \int_a^b \left( \int_0^\infty e^{-st} f(s, x) dt \right) dx = \int_a^b F(s, t) dx,$$

Q.E.D.